

Dissolved Inorganic Carbon (DIC) concentrations from CTD from R/V Tangaroa cruise 61TG_3052 in the Southern Ocean in 1999 (SOIREE project)

Website: <https://www.bco-dmo.org/dataset/2840>

Version: 10August2009

Version Date: 2009-08-10

Project

» [Southern Ocean Iron Release Experiment](#) (SOIREE)

Program

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
Bakker, Dorothee C.E.	University of East Anglia (UEA)	Principal Investigator
Watson, Andrew J.	University of East Anglia (UEA)	Co-Principal Investigator
Mackie, Doug	University of Otago	Contact
Gegg, Stephen R.	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

Dataset Description

SOIREE DIC CTD Data

Note: DIC CTD data are final, except for station 1168!!!!

DICrun1 and DICrun2: DIC values using a running mean over the Dickson values for the actual (3x), preceding (1x) and succeeding cell (1x). The DIC values using the running mean for the Dickson values have a better reproducibility for duplicate samples than, if we use the Dickson value for each cell.

Replicate DIC samples from the CTD:

The average difference between duplicates from the CTD

With running mean of Dickson value:

mean = 2.7 umol/kg, stdev = 2.5 umol/kg, n = 18

Methods & Sampling

See [SOIREE Preliminary Voyage Report](#)

Data Processing Description

See [SOIREE Preliminary Voyage Report](#)

BCO-DMO Processing Notes

Generated from original spreadsheet DIC_CTD.xls
provided on the Deep-Sea Research II 48 (2001) accompanying CD-Rom

BCO-DMO Edits

- parameter names modified to conform to BCO-DMO convention
- blank lines removed
- Station Number changed to station
- added 'T' to CTD Station number for compatibility with events in other spreadsheets
- date reformatted to YYYYMMDD
- time reformatted to HHMM
- date, time, Lat, lon added from SOIREE_CTD_at_Bottle_Trip_Data_FINAL.xls
- CTD data recs with no data in any fields deleted from file
- 'NaN' changed to 'nd'

[[table of contents](#) | [back to top](#)]

Data Files

File
DIC_CTD.csv (Comma Separated Values (.csv), 47.93 KB) MD5:f734a81ba765e2d2749a217a1a0429db Primary data file for dataset ID 2840

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
station	Station Id (Text String)	text
date	date	YYYYMMDD
lon	longitude, negative denotes West	decimal degrees
lat	latitude, negative denotes South	decimal degrees
time	time	HHMM
cast	CTD cast number	integer
trip	Niskin	integer
press	CTD Pressure	decibars
temp	CTD Temperature (ITS-90)	degrees celsius
sal	CTD Salinity (PSS-78)	dimensionless
DICrun1	DIC values Run 1	umol/kg
DICrun2	DIC values Run 2	umol/kg

[[table of contents](#) | [back to top](#)]

Deployments

61TG_3052

Website	https://www.bco-dmo.org/deployment/57827
Platform	R/V Tangaroa
Report	http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf
Start Date	1999-01-31
End Date	1999-03-01
Description	Cruise to the Southern Ocean as part of the Fe Sythesis project whose aim was to maintain a coherent patch of iron-enriched seawater for the duration of SOIREE and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period.

[[table of contents](#) | [back to top](#)]

Project Information

Southern Ocean Iron Release Experiment (SOIREE)

Coverage: Southern Ocean

Project in the Southern Ocean aimed at maintaining a coherent patch of iron-enriched seawater for the duration of project and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period of the project.

The Southern Ocean Iron RElease Experiment (SOIREE), was the first in situ iron fertilization experiment performed in the polar waters of the Southern Ocean. SOIREE was an interdisciplinary study involving participants from six countries, and took place in February 1999 south of the Polar Front in the Australasian-Pacific sector of the Southern Ocean.

Approximately 3800 kg of acidified $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ and 165 g of the tracer sulphur hexafluoride (SF_6) were added to a 65-m deep surface mixed layer over an area of $\sim 50 \text{ km}^2$. Initially, mean dissolved iron concentrations were $\sim 2.7 \text{ nM}$, but decreased to ambient levels within days, requiring subsequent additions of 1550-1750 kg of acidified $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ on days 3, 5 and 7 of the experiment.

During the 13-day site occupation, there were iron-mediated increases in phytoplankton growth rates, with marked increases in chlorophyll a (up to $2 \mu\text{g l}^{-1}$) and production rates (up to $1.3 \text{ gCm}^{-2}\text{d}^{-1}$). These resulted in subsequent changes in the pelagic ecosystem structure, and in the cycling of carbon, silica and sulphur, such as a 10% drawdown of surface CO_2 .

The SOIREE bloom persisted for >40 days following our departure from the site, as observed via [SeaWiFS remotely sensed observations of Ocean Colour](#).

BCO-DMO Note:

All original data and metadata provided on a CD-Rom accompanying the Deep-Sea Research II 48 (2001) volume. The CD-Rom contains the main SOIREE datasets and ancillary information including the pre-experiment 'desktop' database study for site-selection, and satellite images of the SOIREE bloom.

© 2001 Elsevier Science Ltd. All rights reserved.

Related files

[SOIREE Preliminary Voyage Report](#)

[SOIREE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

[SOIREE Cruise Track](#)

[[table of contents](#) | [back to top](#)]

Program Information

Iron Synthesis (FeSynth)

Coverage: Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFs (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as

physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO₂ budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

http://www.scor-int.org/Working_Groups/wg131.htm

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

Related file

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[[table of contents](#) | [back to top](#)]