

Uncalibrated data from the ship's Data Acquisition System (DAS) from R/V Tangaroa cruise 61TG_3052 in the Southern Ocean in 1999 (SOIREE project)

Website: <https://www.bco-dmo.org/dataset/2849>

Version: 16July2008

Version Date: 2008-07-16

Project

» [Southern Ocean Iron Release Experiment](#) (SOIREE)

Program

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
Abraham, Edward	New Zealand National Institute of Water and Atmospheric Research (NIWA)	Principal Investigator
Mackie, Doug	University of Otago	Contact
Gegg, Stephen R.	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

Dataset Description

SOIREE Uncalibrated data from the ship's Data Acquisition System (DAS) Underway data collected during SOIREE, sampled at 5 minute intervals
The file also contains shipboard meteorological data.

The data are presented in a raw, uncalibrated form.
Missing data are marked by "nd".

Methods & Sampling

See [SOIREE Preliminary Voyage Report](#)
See comments in dataset description as well

A Chelsea instruments CTD/Fluorometer was mounted in a chamber that was supplied with

flow-through sea-water. From this instrument Temperature and Fluorescence were logged.

A Chelsea Instruments Fasttracka FRRF was used to record phytoplankton photosynthetic

competency (Fv/Fm).

Data Processing Description

See [SOIREE Preliminary Voyage Report](#)
See comments in dataset description as well

BCO-DMO Processing Notes

Generated from original file das_5minute.txt provided on the Deep-Sea Research II 48 (2001) accompanying CD-Rom

BCO-DMO Edits

- Parameter names modified to conform to BCO-DMO convention
- date reformatted to YYYYMMDD
- time reformatted to HHMM
- Note: Some times with 59 seconds truncated when formatted for BCO-DMO
- lat/lon values padded to 4 decimal places
- other data values padded to 2 decimal places
- Note: Wind Speed and Wind Direction appear to have same values

[[table of contents](#) | [back to top](#)]

Data Files

File
DAS.csv (Comma Separated Values (.csv), 541.24 KB) MD5:695a2a5cc695de5867fb0b6dfcb8ef05 Primary data file for dataset ID 2849

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
date	date	YYYYMMDD
lon	longitude, negative denotes West	decimal degrees
lat	latitude, negative denotes South	decimal degrees
time	time	HHMM
SeaTemp	Sea Surface Temperature	degrees celsius
fluor	Fluorescence	volts
Fv_to_Fm_from_FRRF	Ratio of Phytoplankton Photosynthetic Competency from Fast Repetition Rate Fluorometry	dimensionless
WindSpeed	Wind Speed	knots
WindDirection	Wind Direction	degrees
AirTemp	Air Temperature	degrees celsius
BarometricPressure	Barometric Pressure	decibars
RelativeHumidity	Relative Humidity	percentage

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	CTD/Fluorometer
Generic Instrument Name	CTD-FRRfluorometer
Dataset-specific Description	Chelsea instruments CTD-Fluorometer was used to log Temperature and Fluorescence
Generic Instrument Description	A CTD-FRRf fluorometer is an instrument package designed to measure hydrographic information (pressure, temperature and conductivity) and chlorophyll fluorescence. (see more at URL: http://www.chelsea.co.uk/Instruments%20FASTtracka.htm) For example, a Chelsea FASTtracka CTD (conductivity, temperature, and depth) device can be configured with additional sensors to measure fluorescence and photosynthetically active radiation (PAR) in the water column. Fluorescence data collected by the CTD are subsequently calibrated and corrected to give the average concentration of chlorophyll a in the water column. The Chelsea system can be configured as part of a towed package or in a shipboard pumping system. (see more from Chelsea Instruments, Molesey, Surrey, United Kingdom at URL: http://www.chelsea.co.uk/)

Dataset-specific Instrument Name	Fast Repetition Rate Fluorometer
Generic Instrument Name	Fast Repetition Rate Fluorometer
Dataset-specific Description	A Chelsea Instruments Fasttracka FRRF was used to record phytoplankton photosynthetic competency (Fv/Fm)
Generic Instrument Description	An FRRf is used for measuring the fluorescence of a sample of phytoplankton photosynthetic competency (Fv/Fm).

[[table of contents](#) | [back to top](#)]

Deployments

61TG_3052

Website	https://www.bco-dmo.org/deployment/57827
Platform	R/V Tangaroa
Report	http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf
Start Date	1999-01-31
End Date	1999-03-01
Description	Cruise to the Southern Ocean as part of the Fe Sythesis project whose aim was to maintain a coherent patch of iron-enriched seawater for the duration of SOIREE and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period.

[[table of contents](#) | [back to top](#)]

Project Information

Southern Ocean Iron Release Experiment (SOIREE)

Coverage: Southern Ocean

Project in the Southern Ocean aimed at maintaining a coherent patch of iron-enriched seawater for the duration of project and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period of the project.

The Southern Ocean Iron RElease Experiment (SOIREE), was the first in situ iron fertilization experiment performed in the polar waters of the Southern Ocean. SOIREE was an interdisciplinary study involving participants from six countries, and took place in February 1999 south of the Polar Front in the Australasian-Pacific sector of the Southern Ocean.

Approximately 3800 kg of acidified FeSO₄·7H₂O and 165 g of the tracer sulphur hexafluoride (SF₆) were added to a 65-m deep surface mixed layer over an area of ~50 km². Initially, mean dissolved iron concentrations were ~2.7 nM, but decreased to ambient levels within days, requiring subsequent additions of 1550-1750 kg of acidified FeSO₄·7H₂O on days 3, 5 and 7 of the experiment.

During the 13-day site occupation, there were iron-mediated increases in phytoplankton growth rates, with marked increases in chlorophyll a (up to 2 µg l⁻¹) and production rates (up to 1.3 gCm⁻²d⁻¹). These resulted in subsequent changes in the pelagic ecosystem structure, and in the cycling of carbon, silica and sulphur, such as a 10% drawdown of surface CO₂.

The SOIREE bloom persisted for >40 days following our departure from the site, as observed via [SeaWiFS remotely sensed observations of Ocean Colour](#).

BCO-DMO Note:

All original data and metadata provided on a CD-Rom accompanying the Deep-Sea Research II 48 (2001) volume. The CD-Rom contains the main SOIREE datasets and ancillary information including the pre-experiment 'desktop' database study for site-selection, and satellite images of the SOIREE bloom.

© 2001 Elsevier Science Ltd. All rights reserved.

Related files

[SOIREE Preliminary Voyage Report](#)

[SOIREE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

[SOIREE Cruise Track](#)

[[table of contents](#) | [back to top](#)]

Program Information

Iron Synthesis (FeSynth)

Coverage: Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFS (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO₂ budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

http://www.scor-int.org/Working_Groups/wg131.htm

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata

(e.g., description of project/study, general location, PI(s), participants);

2. Level 2, detailed metadata and basic shipboard data and routine ship's operations

(e.g., CTDs, underway measurements, sampling event logs);

3. Level 3, detailed metadata and data from specialized observations

(e.g., discrete observations, experimental results, rate measurements) and

4. Level 4, remaining datasets

(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

Related file

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[[table of contents](#) | [back to top](#)]