

# Nutrient data containing NH<sub>4</sub>, SiO<sub>4</sub>, PO<sub>4</sub>, NO<sub>2</sub>+NO<sub>3</sub> from R/V Atlantis AT16 from the Northwest Atlantic Margin (Carbon Export project)

**Website:** <https://www.bco-dmo.org/dataset/3701>

**Version:** 2012-08-16

## Project

» [Particle Transport and Carbon Export over the Northwest Atlantic Margin](#) (Carbon Export)

## Program

» [Ocean Carbon and Biogeochemistry](#) (OCB)

Contributors	Affiliation	Role
<a href="#">Eglinton, Timothy I.</a>	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
<a href="#">Henderson, Paul B.</a>	Woods Hole Oceanographic Institution (WHOI)	Analyst
<a href="#">Chandler, Cynthia L.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

---

## Table of Contents

- [Dataset Description](#)
  - [Data Processing Description](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

---

## Dataset Description

Bottle nutrient data collected along Line W. Parameters include NH<sub>4</sub>, SiO<sub>4</sub>, PO<sub>4</sub>, NO<sub>2</sub>+NO<sub>3</sub>.

There are some anomalies that the PI wants to track down before making the data public.

## Data Processing Description

Bottle nutrient samples were processed at the Woods Hole Oceanographic Institution Nutrient Analytical Facility, which operates a Lachat Instruments QuickChem 8000 four-channel continuous flow injection system in addition to in-situ chemical analyzers (Ecolab & NAS) to determine dissolved nutrient concentrations in aquatic ecosystems. For additional information see <http://www.whoi.edu/nutrient>

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
station	Station number.	dimensionless
lat	Latitude in decimal degrees, where negative values denote the southern hemisphere.	decimal degrees
lon	Longitude in decimal degrees, where negative values denote western hemisphere.	decimal degrees
depth	depth in meters.	meters
event	Unique identifier for a sampling or processing event.	dimensionless
bottle	Niskin bottle number, originally reported as Niskin.	dimensionless
sample	Unique identifier for sample.	dimensionless
NH4	Ammonium.	micromolar
SiO4	Silicate.	micromolar
PO4	Phosphate.	micromolar
cast	Cast identifier.	dimensionless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Niskin bottle
<b>Generic Instrument Name</b>	Niskin bottle
<b>Generic Instrument Description</b>	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### AT16

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58835">https://www.bco-dmo.org/deployment/58835</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2010-09-10
<b>End Date</b>	2010-10-05
<b>Description</b>	<p>Science Objectives The overall goal of the project is to examine the transport and export of particulate organic carbon (POC) over the northwest Atlantic Ocean margin. Specific objectives are to examine the role of the Deep Western Boundary Current in along-margin transport of POC. It is hypothesized that along- and across-margin transport of POC is important for supply of organic carbon to deep-ocean sediments. Geochemical investigations of organic matter entrained in bottom and intermediate depth suspended particle (nepheloid) layers as well as sinking particles are planned as well as hydrographic surveys in order to place findings within the context of regional physical oceanography. Hydrographic surveys and sampling will occur along 3 across-margin transects in each case extending from the shelf-break to ~ 5000 m water depth (off Nova Scotia, off Cape Cod, and off Cape Hatteras). The assessments of particle dynamics will be coupled to an existing hydrographic observation program along an ocean section between Cape Cod and Bermuda ('Line W'). The 'Line W' program is focused on the physical characteristics of the region, and provides an accompanying framework for the study of carbon dynamics. A series of 3 across-margin transects are designed to capture the sources and movement of particulate matter. The proposed sampling program includes CTD hydrocasts and large-volume (up to 1000L) in-situ filtration with McLane submersible pumps. The science plan also calls for deployment of 2 sediment trap moorings and perhaps some sediment coring (multicorer). The intent is to obtain high vertical and spatial resolution filter samples for detailed analysis of particle abundance and characteristics within intermediate-depth and bottom nepheloid layers. The science implementation plan includes deployment of: Sea-Bird Electronics, Inc. CTD rosette equipped with transmissometer and optical back scatter sensor; Glider Autonomous Underwater Vehicle (AUV) equipped with CTD and optical backscatter sensor; Sea floor-tethered mooring with Deployment of McLane Mark-7 time-series sediment traps; ship's clean seawater intake and McLane submersible pumps (Water Transfer System, WTS). Funding for cruise AT16 is from NSF award number OCE-0851350. Cruise information and original data are available from the NSF R2R data catalog.</p>

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Particle Transport and Carbon Export over the Northwest Atlantic Margin (Carbon Export)

**Coverage:** Nova Scotian continental slope waters; Cape Hatteras margin slope waters

This award is funded under the American Recovery and Reinvestment Act of 2009 (Public Law 111-5).

In this project researchers at the Woods Hole Oceanographic Institution (WHOI) seek to address two primary questions regarding the transport of organic carbon over the New England continental margin. Their first question addresses the relative importance of intermediate nepheloid layers versus benthic nepheloid layers for organic carbon transport. Their second question focuses on assessing the importance of interannual variability on particle transport. The researchers will collect both settling and suspended particles along the NW Atlantic ocean margin, and they will utilize both novel and established tracers of material transport to address their primary research questions. Their assessments of particle dynamics will be coupled to an existing hydrographic observation program along an ocean section between Cape Cod and Bermuda (Line W). This program is focused on the physical characteristics of the region, and provides an accompanying framework for the study of carbon dynamics.

The importance of the work to global carbon cycling lies in the fact that continental margins generally maintain high biological production and are physically dynamic environments. One potential fate of the biologically produced carbon is that it may be exported to the deep ocean, yet the magnitude of that exported carbon is generally not well defined. Over the Northwest Atlantic margin, large-scale currents have the potential to be an important dispersal mechanism for organic carbon to the ocean interior. In a prior research project, the investigators hypothesized that the lateral transport of particulate organic carbon occurs via a bottom nepheloid layer (BNL) maintained by the Deep Western Boundary Current (DWBC). Although their research supports the notion that the DWBC transports a significant amount of organic carbon, their observations also indicate that intermediate nepheloid layers (INLs) emanating from the shelf-slope break and upper slope of the Mid Atlantic Bight may also be important vectors for carbon export.

[ [table of contents](#) | [back to top](#) ]

---

## Program Information

### Ocean Carbon and Biogeochemistry (OCB)

**Website:** <http://us-ocb.org/>

**Coverage:** Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO<sub>2</sub> and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0851350</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0726720</a>

[ [table of contents](#) | [back to top](#) ]