

# Cruise tracks for the CROCKS II and HROCKS projects from R/V Atlantis AT15-44, AT15-59, AT15-68, AT18-10 in the Pacific, off Costa Rica and USA from 2009-2011 (Seep Carbonate Ecology CROCKS II project)

**Website:** <https://www.bco-dmo.org/dataset/3760>

**Version:**

**Version Date:** 2012-10-25

## Project

» [Short-term colonization processes at Costa Rica methane seeps](#) (Seep Carbonate Ecology CROCKS II)

Contributors	Affiliation	Role
<a href="#">Levin, Lisa A.</a>	University of California-San Diego Scripps (UCSD-SIO)	Principal Investigator
<a href="#">Rouse, Gregory</a>	University of California-San Diego Scripps (UCSD-SIO)	Co-Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Dataset Description

Position, date and time data collected daily along the ship track for cruises AT15-44, AT15-59, AT15-68, and AT18-10

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>cruisetracks.csv</b> (Comma Separated Values (.csv), 4.50 MB) MD5:03769928d77638eaadd5f3a24f49aa24
Primary data file for dataset ID 3760

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
cruise_id	cruise identifier; AT=Atlantis	text
year1	year of cruise	YYYY
yrday_gmt	GMT day and decimal time. e.g. as 326.5 for the 326th day of the year or November 22 at 1200 hours (noon).	nd
month_gmt	month UTC time	mm
day_gmt	day UTC time	dd
time_gmt	GMT time	HHMM.SS
ISOdatetime	Date/Time (UTC) ISO formatted	YYYY-mm-ddTHH:MM:SS[time zone; [Z=UTC]
lon	longitude; East is positive	decimal degrees
lat	latitude; North is positive	decimal degrees

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Global Positioning System Receiver
<b>Generic Instrument Name</b>	Global Positioning System Receiver
<b>Generic Instrument Description</b>	The Global Positioning System (GPS) is a U.S. space-based radionavigation system that provides reliable positioning, navigation, and timing services to civilian users on a continuous worldwide basis. The U.S. Air Force develops, maintains, and operates the space and control segments of the NAVSTAR GPS transmitter system. Ships use a variety of receivers (e.g. Trimble and Ashtech) to interpret the GPS signal and determine accurate latitude and longitude.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

**AT15-44**

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58869">https://www.bco-dmo.org/deployment/58869</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2009-02-21
<b>End Date</b>	2009-03-08
<b>Description</b>	Cruise Objective: We will conduct research in exposed carbonate ecosystems on the Costa Rica margin (700-1,400 m), to test hypotheses about the influence of active seepage on carbonate rock animal communities and their successional phases, on microbial activity including anaerobic methane oxidation and sulfide oxidation, on carbon isotopic composition of shelled organisms, and on phylogenetic affinities of animals. To test hypotheses we will sample existing authigenic carbonates from 3 levels of seepage activity: highly active, weak and inactive. Activity level will be defined by presence of /or proximity to bubbles/shimmering water, microbial mat development and megafauna, as well as previous fluid flow and composition measurements made at the Costa Rica study sites. We will sample 5 to 8 locations with each activity level in each study region, controlling for rock size and carbonate configuration when possible. ALVIN: During 3 dives at each of 4 study sites we will conduct bottom surveys and video transects, measure S, T, O <sub>2</sub> , select 4 to 8 highly active, weakly active and inactive sites, photograph organisms and classify rocks in situ, collect rocks of varying sizes with organisms, and sample nearby sediments and biotic substrata (mussels, tube worms) for taxonomic comparisons. The remaining 2 dives at Costa Rica seeps will be used to conduct follow-up survey and sampling of the most promising locations, based on shipboard sample observations. Nighttime operations will consist of CTD casts (a minimum of one each at Mound 11, Mound 13, Jaco Scarp and Mound Quepos), multicoring (adjacent to mounds and at 400 m and 600 m sites in the OMZ), and pre-dive seabeam surveys. Cruise information and original data are available from the NSF R2R data catalog.

#### AT15-59

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58765">https://www.bco-dmo.org/deployment/58765</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2010-01-06
<b>End Date</b>	2010-01-13
<b>Description</b>	Costa Rica seafloor methane seeps 8 deg 55 N 84 depth 990m. Included Alvin dives 4586-4591. The primary goal of the cruise was to recover biological experiments deployed at active and inactive seep areas during Feb./March 2009. We successfully recovered 23/24 experimental units deployed on Mound 12, Costa Rica. One was simply missing on the sea floor. By using a gear elevator each day to maximize sample collection (and reserve ALVIN basket space for experiments) we were able to recover our experiments rapidly. This left us time for exploration of unusual biological communities at Jaco Scar. Cruise information and original data are available from the NSF R2R data catalog. Cruise dates changed Feb 2015 to match WHOI ship schedule and R2R.

#### AT15-68

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58870">https://www.bco-dmo.org/deployment/58870</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2010-07-31
<b>End Date</b>	2010-08-12
<b>Description</b>	Cruise information and original data are available from the NSF R2R data catalog.

#### AT18-10

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58871">https://www.bco-dmo.org/deployment/58871</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2011-08-31
<b>End Date</b>	2011-09-08
<b>Description</b>	Cruise information and original data are available from the NSF R2R data catalog.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Short-term colonization processes at Costa Rica methane seeps (Seep Carbonate Ecology CROCKS II)

**Coverage:** Costa Rica seafloor methane seeps 8 deg 55 N 84 deg 18 W depth 990m

This RAPID project will conduct 5 submersible or ROV dives to collect a series of colonization experiments deployed in March 2009 on Mound 12 off Costa Rica (997 m). These experiments were deployed opportunistically, and to optimize the information that could be obtained, the PIs needed to recover them within a 12 month time frame. Early colonization of rock, wood, shell and tube substrates will be studied. The microbes, foraminiferans and metazoans present after 6-12 mo will be compared to those colonizing similar experiments to be deployed at Hydrate Ridge, where seeps occur within an oxygen minimum zone. The overall project goal is to integrate physical, geological, chemical and biological data to develop a holistic view of the influence of seep-generated carbonate hard-ground ecosystems on margins.

The objectives of the research are to (a) Compare colonizers at seeps off Costa Rica and Hydrate Ridge to assess the importance of different oxygen regimes in the development of anaerobic methane oxidation, sulfide oxidizers and other microbial metabolisms on hard substrates, and to evaluate their roles in driving protozoan and metazoan succession at methane seeps. (b) Deploy a suite of biotic and abiotic substrates to distinguish the specific roles of carbonate substrate from those of other hard substrates (wood, clam and mussel shells, worm tubes) available. (c) Explore the similarity of vent and seep colonization processes by comparing colonization at the Costa Rica seeps, where vent species are common, to the Hydrate Ridge seeps, where they are not. (d) Determine whether there are diagnosable biogeographic isotope or other biomarker signatures from newly recruited Costa Rica microbial, foraminiferal and animal populations at active vs. inactive seeps, and whether these differ from those at Hydrate Ridge.

This research will involve international participation from Costa Rican scientists at the Univ. of Costa Rica.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0825791</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0826254</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0939559</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0825436</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0939232</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0939557</a>

[ [table of contents](#) | [back to top](#) ]