

Counts of *Henricia* sp. adults and recruits in quadrats, north-central California coast: Andrew Molean State Park to Manchester State Park from 2005-2014 (CHIPS project)

Website: <https://www.bco-dmo.org/dataset/528964>

Version: 4

Version Date: 2015-07-15

Project

» [Ecological & genetic recovery from a massive invertebrate die-off along the central coast of California \(CHIPS\)](#)

Contributors	Affiliation	Role
Gaylord, Brian	University of California - Davis: Bodega Marine Laboratory (UC Davis-BML)	Principal Investigator
Dawson, Michael N.	University of California-Merced (UC Merced)	Co-Principal Investigator
Grosberg, Richard K.	University of California-Davis (UC Davis)	Co-Principal Investigator
Jurgens, Laura J.	University of California - Davis: Bodega Marine Laboratory (UC Davis-BML)	Student, Contact
Schiebelhut, Lauren	University of California-Merced (UC Merced)	Student, Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Twenty accessible rocky intertidal sites were selected to span the 'kill-zone' and peripheral locations (see list of sites in Jurgens et al. submitted). Two rocky intertidal areas were sampled per site, usually one on either side of the point of entry onto the beach and separated by approximately 100 m. We used quadrats (1m² [4 * 0.25 m²] or 0.0625 m²) and swaths (2 m wide transects) to estimate juvenile and adult abundance for each target species.

This data set includes counts of *Henricia* sp. (Pacific blood star) in quadrats along the California coast from 36.3 N to 39.2 N.

Access Restriction: Access to these data are restricted until publication of the associated manuscript, which is currently in review at PLOS ONE Progress Series. [2015-02-05]

Version 4 (2015-07-15) replaces version 3 (2014-10-03). Values for the sampled area were corrected.

Related Datasets:

[Purple urchin density](#)
[Purple urchin density by depth](#)
[Seastar and chiton counts - quadrats and swaths](#)
[CHIPS02 - seastar chiton urchin counts in quadrats](#)

Methods & Sampling

Quadrats for *C. stelleri*, *Henricia sp.*, *Leptasterias sp.*, *P. ochraceus*:

We exhaustively searched 32-40 one-meter square quadrats per site (i.e. 16-20 per each of 2 areas), recording GPS waypoint, time, percent cover of major habitat types, and abundances and sizes of target species for each quadrat. Quadrat locations were selected by first finding one of the target habitat types - surf grass, low-zone red algae, coralline turf, cobble or boulder field, urchin pools with pits either empty or occupied, or mussel bed - selecting a starting point haphazardly, and then using a random numbers table to choose specific quadrat locations. Some sites had fewer than 32 quadrats or quadrats smaller than one-square meter due to the limited time in which we could work in suitable habitat. For mussel habitat one 0.25 m x 0.25 m quadrat was used. For Jurgens et al. (submitted) we calculated the number of each target species summed across all quadrats surveyed within each of the two areas within sites.

Swaths for *C. stelleri*, *Henricia sp.*, *Leptasterias sp.*, *P. ochraceus*:


Timed, GPS-tracked, 2 m wide swath transects were nested within each of the two areas within all sites. From a distance, an approximate starting point and orientation (with landmarks) for the starting transect was selected. The GPS was set to auto-record a trackpoint every 6 seconds. Transects ran from the most shoreward to the most seaward possible suitable habitat at approximately 10 m intervals along shore, particularly targeting the low zone when the tide was the lowest, with as many transects being done as permitted by the tide. The start and end of each transect was GPS waypointed and recorded. All target species were searched for across the 2 m wide swath, with intertidal zones and habitat type recorded. A waypoint was recorded for each individual target species, with any additional species found within a meter being counted and recorded for the same waypoint. When the field of view was clear some swaths were extended to 4 meters wide. For Jurgens et al. (submitted) we calculated the number of each target species summed across all swaths within each of the two areas within sites.

Data Processing Description

To estimate swath length we first summed the Euclidean distances between consecutive GPS trackpoints.

Given the GPS had an error of +/- 3 meters we smoothed the GPS tracks by averaging across two trackpoints and then by three. We selected a window size of two with which to estimate our total transect length to adjust for the noise in our swath transects but to minimize oversmoothing. (see example figure below). We identified and removed outlying trackpoints that led to a Euclidean distance ≥ 8 meters (likely due to temporarily poor satellite signal). The distance was then recalculated between the trackpoints immediately preceding and following the removed trackpoint. Total transect search area was calculated by multiplying the total adjusted transect length by swath width. Total transect search time was estimated using field notes and GPS waypoints and trackpoints, removing 6-second intervals where < 0.05 m were traveled (suggesting this was time during which notes were being written or tissues samples were being collected).

Swathfigure: Sample GPS tracks for two transects at Iversen Point, Mendocino County. The blue line is the original unsmoothed track. The red line represents a window size of two, averaging latitudes and longitudes across each pair of consecutive trackpoints. The green line represents a window size of 3, averaging across 3 trackpoints.



BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date, reference information
- renamed parameters to BCO-DMO standard
- sorted by latitude, high to low, and date

[[table of contents](#) | [back to top](#)]

Data Files

File
henricia.csv (Comma Separated Values (.csv), 2.15 KB) MD5:05ec3d32964debc8a97aed6a547b80c4 Primary data file for dataset ID 528964

[[table of contents](#) | [back to top](#)]

Related Datasets

IsRelatedTo

Dawson, M. N., Gaylord, B., Grosberg, R. K. (2022) **Comparison of recruitment dynamics in five intertidal marine invertebrates following mass mortality along the northeastern Pacific coastline in 2005 (CHIPS project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2015-08-06 doi:10.26008/1912/bco-dmo.562467.1 [[view at BCO-DMO](#)]

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
site	survey site name	unitless
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
loss_zone	The site is inside or outside the region across which intertidal <i>S. purpuratus</i> were extirpated. Unknown means that there was no pre-HAB urchin population.	unitless
date_survey	date of survey	YYYY-MM-DD
area	Total square meters surveyed at Site; Year or time-period may be specified.	m ²
count_Henricia	number of <i>Henricia</i> adults and recruits found within one quadrat	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	GPS
Generic Instrument Name	Global Positioning System Receiver
Generic Instrument Description	The Global Positioning System (GPS) is a U.S. space-based radionavigation system that provides reliable positioning, navigation, and timing services to civilian users on a continuous worldwide basis. The U.S. Air Force develops, maintains, and operates the space and control segments of the NAVSTAR GPS transmitter system. Ships use a variety of receivers (e.g. Trimble and Ashtech) to interpret the GPS signal and determine accurate latitude and longitude.

[[table of contents](#) | [back to top](#)]

Deployments

CalifCoast_Gaylord_CHIPS

Website	https://www.bco-dmo.org/deployment/528985
Platform	shoreside Calif_shore
Start Date	2005-01-01
End Date	2014-05-15
Description	Various intertidal invertebrate studies.

Project Information

Ecological & genetic recovery from a massive invertebrate die-off along the central coast of California (CHIPS)

Coverage: Northeastern Pacific (northern California) 38-39 N

This project is a Collaborative Research project funded by an NSF RAPID grant.

Description from NSF award abstract:

The potentially important role that larval dispersal may play in determining gene flow, distributions, and population structure of marine invertebrates remains unclear despite many hundreds of descriptive comparisons of pelagic duration and population genetic structure. This lack of clarity suggests many factors may influence population genetic structure and their interactions may be complex. Difficulties studying these factors include (under normal circumstances) distinguishing local from exogenous recruitment and therefore the true distribution of dispersal distances. For example, experiments that normally could be undertaken to explore this issue are very small scale relative to the distances that many marine taxa may disperse.

In August 2011, a large-scale natural removal experiment was initiated along a 100 km stretch of the central California coast. The PIs propose to use this rare opportunity to clarify the effects of dispersal and species interactions on marine population genetic variation and community structure. They propose to study three species that suffered very high rates of mortality: an ecosystem engineer (*Strongylocentrotus purpuratus*, ~100% mortality), a keystone species (*Pisaster ochraceus*, ~10-70% mortality), and one of its competitors (*Leptasterias* sp., ~100% mortality). Their objectives during this first year following the natural large-scale die-off are to:

- (1) quantify the abundance and distribution of the target species at sites across the impacted range and reference sites to the south and north,
- (2) develop and use genetic markers to identify the sources and dispersal distances of new recruits of *P. ochraceus*, *Leptasterias* sp., and *S. purpuratus* that recolonize the impacted range, and
- (3) describe changes in abundance of these three species and their prey and competitors at sites throughout the impacted range.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1243970
NSF Division of Ocean Sciences (NSF OCE)	OCE-1243958