

# Underway Data from R/V Melville, R/V Roger Revelle cruises MV1101, RR1202 in the Southern Ocean (30-60S); 2011-2012 (Great Calcite Belt project)

Website: <https://www.bco-dmo.org/dataset/560142>

Version: 08 June 2015

Version Date: 2015-06-08

## Project

» [The Great Southern Coccolithophore Belt](#) (Great Calcite Belt)

Contributors	Affiliation	Role
<a href="#">Balch, William M.</a>	Bigelow Laboratory for Ocean Sciences	Principal Investigator, Contact
<a href="#">Gegg, Stephen R.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Dataset Description

Along track temperature, Salinity, backscatter, Chlorophyll Fluorescence, and normalized water leaving radiance (nLw).

On the bow of the vessel was a Satlantic SeaWiFS Aircraft Simulator (MicroSAS) system, used to estimate water-leaving radiance from the ship, analogous to the nLw derived by the SeaWiFS and MODIS satellite sensors, but free from atmospheric error (hence, it can provide data below clouds).

The system consisted of a down-looking radiance sensor and a sky-viewing radiance sensor, both mounted on a steerable holder on the bow. A downwelling irradiance sensor was mounted at the top of the ship's meteorological mast, on the bow, far from any potentially shading structures. These data were used to estimate normalized water-leaving radiance as a function of wavelength. The radiance detector was set to view the water at 40deg from nadir as recommended by Mueller et al. [2003b]. The water radiance sensor was able to view over an azimuth range of ~180deg across the ship's heading with no viewing of the ship's wake. The direction of the sensor was adjusted to view the water 90-120deg from the sun's azimuth, to minimize sun glint. This was continually adjusted as the time and ship's gyro heading were used to calculate the sun's position using an astronomical solar position subroutine interfaced with a stepping motor which was attached to the radiometer mount (designed and fabricated at Bigelow Laboratory for Ocean Sciences). Protocols for operation and calibration were performed according to Mueller [Mueller et al., 2003a; Mueller et al., 2003b; Mueller et al., 2003c]. Before 1000h and after 1400h, data quality was poorer as the solar zenith angle was too low. Post-cruise, the 10Hz data were filtered to remove as much residual white cap and glint as possible (we accept the lowest 5% of the data). Reflectance plaque measurements were made several times at local apparent noon on sunny days to verify the radiometer calibrations.

Within an hour of local apparent noon each day, a Satlantic OCP sensor was deployed off the stern of the vessel after the ship oriented so that the sun was off the stern. The ship would secure the starboard Z-drive, and use port Z-drive and bow thruster to move the ship ahead at about 25cm s<sup>-1</sup>. The OCP was then trailed aft and brought to the surface ~100m aft of the ship, then allowed to sink to 100m as downwelling spectral irradiance and upwelling spectral radiance were recorded continuously along with temperature and salinity. This procedure ensured there were no ship shadow effects in the radiometry.

Instruments include a WETLabs wetstar fluorometer, a WETLabs ECOTriplet and a SeaBird microTSG. Radiometry was done using a Satlantic 7 channel microSAS system with Es, Lt and Li sensors.

Chl data is based on inter calibrating surface discrete Chlorophyll measure with the temporally closest fluorescence measurement and applying the regression results to all fluorescence data.

Data have been corrected for instrument biofouling and drift based on weekly purewater calibrations of the system. Radiometric data has been processed using standard Satlantic processing software and has been checked with periodic plaque measurements using a 2% spectralon standard.

Lw is calculated from Lt and Lsky and is "what Lt would be if the sensor were looking straight down". Since our sensors are mounted at 40o, based on various NASA protocols, we need to do that conversion.

Lwn adds Es to the mix. Es is used to normalize Lw. Nlw is related to Rrs, Remote Sensing Reflectance

### Techniques used are as described in:

Balch WM, Drapeau DT, Bowler BC, Booth ES, Windecker LA, Ashe A (2008) Space-time variability of carbon standing stocks and fixation rates in the Gulf of Maine, along the GNATS transect between Portland, ME, USA, and Yarmouth, Nova Scotia, Canada. J Plankton Res 30:119-139

## Methods & Sampling

Instruments include a WETLabs wetstar fluorometer, a WETLabs ECOTriplet and a SeaBird microTSG. Radiometry was done using a Satlantic 7 channel microSAS system with Es, Lt and Li sensors.

Chl data is based on inter calibrating surface discrete Chlorophyll measure with the temporally closest fluorescence measurement and applying the regression results to all fluorescence data.

### Techniques used are as described in:

Balch WM, Drapeau DT, Bowler BC, Booth ES, Windecker LA, Ashe A (2008) Space-time variability of carbon standing stocks and fixation rates in the Gulf of Maine, along the GNATS transect between Portland, ME, USA, and Yarmouth, Nova Scotia, Canada. J Plankton Res 30:119-139

## Data Processing Description

### Techniques used are as described in:

Balch WM, Drapeau DT, Bowler BC, Booth ES, Windecker LA, Ashe A (2008) Space-time variability of carbon standing stocks and fixation rates in the Gulf of Maine, along the GNATS transect between Portland, ME, USA, and Yarmouth, Nova Scotia, Canada. J Plankton Res 30:119-139

Data have been corrected for instrument biofouling and drift based on weekly purewater calibrations of the system. Radiometric data has been processed using standard Satlantic processing software and has been checked with periodic plaque measurements using a 2% spectralon standard.

## BCO-DMO Processing Notes

- Generated from original files "mv1101-merged-SAS-flow.out" and "rr1202-merged-SAS-flow.out" contributed by Bruce Bowler

- Date reformatted to YYYYMMDD
- Time reformatted to HHMMSS
- Parameter names edited to conform to BCO-DMO naming convention found at [Choosing Parameter Name](#)

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>Underway_Data.csv</b> (Comma Separated Values (.csv), 3.45 MB) <small>MDS:b84cb63ea471f21eebfa9e8f93f942</small>
Primary data file for dataset ID 560142

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
CruiseId	Official UNOLS cruise id	text
Measurement_Depth	Measurement Depth	meters
ISO_DateTime_UTC	ISO DateTime UTC	YYYY-MM-DDTHH:MM:SS[.xx]Z
date	Date (UTC)	YYYYMMDD
time	Time (UTC)	HHMMSS
lat	Latitude Position (South is negative)	decimal degrees
lon	Longitude Position (West is negative)	decimal degrees
Wt	Water temperature	degreesC
sal	Salinity	PSU
chl	Chlorophyll Fluoresence	mg/m <sup>3</sup>
bbp532	Backscatter@532nm	1/m
es412	irradiance at 412nm	uW/cm <sup>2</sup> /nm
es443	irradiance at 443nm	uW/cm <sup>2</sup> /nm
es490	irradiance at 490nm	uW/cm <sup>2</sup> /nm
es510	irradiance at 510nm	uW/cm <sup>2</sup> /nm
es531	irradiance at 531nm	uW/cm <sup>2</sup> /nm
es555	irradiance at 555nm	uW/cm <sup>2</sup> /nm
es670	irradiance at 670nm	uW/cm <sup>2</sup> /nm
Lt412	water radiance at 412nm	uW/cm <sup>2</sup> /nm/sr
Lt443	water radiance at 443nm	uW/cm <sup>2</sup> /nm/sr
Lt490	water radiance at 490nm	uW/cm <sup>2</sup> /nm/sr
Lt510	water radiance at 510nm	uW/cm <sup>2</sup> /nm/sr
Lt531	water radiance at 531nm	uW/cm <sup>2</sup> /nm/sr
Lt555	water radiance at 555nm	uW/cm <sup>2</sup> /nm/sr

Lt670	water radiance at 670nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky412	sky radiance at 412nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky443	sky radiance at 443nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky490	sky radiance at 490nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky510	sky radiance at 510nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky531	sky radiance at 531nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky555	sky radiance at 555nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lsky670	sky radiance at 670nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw412	water leaving radiance at 412nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw443	water leaving radiance at 443nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw490	water leaving radiance at 490nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw510	water leaving radiance at 510nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw531	water leaving radiance at 531nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw555	water leaving radiance at 555nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lw670	water leaving radiance at 670nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn412	normalized water leaving radiance at 412nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn443	normalized water leaving radiance at 443nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn490	normalized water leaving radiance at 490nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn510	normalized water leaving radiance at 510nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn531	normalized water leaving radiance at 531nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn555	normalized water leaving radiance at 555nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
Lwn670	normalized water leaving radiance at 670nm	$\mu\text{W}/\text{cm}^2/\text{nm}/\text{sr}$
relaz	relative azimuth	degrees

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Fluorometer
<b>Generic Instrument Name</b>	Fluorometer
<b>Dataset-specific Description</b>	WETLabs wetstar fluorometer Product Brochure
<b>Generic Instrument Description</b>	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.



feature occurs every year during austral summer and appears to be within the high-nutrient, low chlorophyll region of the Southern Ocean. The area of the Great Belt is ~88 million km<sup>2</sup>, 26% of the global ocean. Evidence from several cruises into the Great Belt region of the Atlantic, Indian and Pacific sectors has verified elevated concentrations of coccolithophores; previous work in the Atlantic sector verified high optical scattering from PIC. The few ship observations we have are entirely consistent with the satellite views. In this project, the investigators will systematically study the coccolithophores of the Great Belt guided by the following science goals: (a) identify the coccolithophore species within this belt; (b) measure the abundance of coccolithophores and associated PIC; (c) measure coccolithophore calcification rates; (d) elucidate factors that may limit coccolithophore latitudinal range (e.g. stratification, temperature, macronutrients, trace metals, grazing); (e) demonstrate whether the variability in PIC relates to shallow export flux; (f) define how variability in PIC production relates to the pCO<sub>2</sub>, total alkalinity and dissolved inorganic carbon budgets; and (g) examine the impact of short-term ocean acidification on coccolithophore growth and calcite dissolution.

The research will involve cruises along the 50 S parallel to sample the Great Belt, during the austral summer. The investigators will use a combination of underway surface sampling (primarily optical and hydrographic) and vertical station profiles (using CTD/rosette and large volume submersible pumps) to address hypotheses related to the above goals. The cruise track will elucidate both zonal and meridional variability in the Great Belt. Controlled carboy incubation experiments will examine the impact of ocean acidification at various future scenarios on coccolithophore growth and dissolution. Dilution experiments will address grazing-related mortality and dissolution questions. Controlled metal-addition incubations will focus on potential iron, zinc and cobalt limitation of the coccolithophores or competition from diatoms related to silica availability. The proposed field observations and metal-addition experiments will provide important information on the current status of the Great Belt in the context of global biogeochemistry. The ocean acidification experiments to be undertaken are more forward-looking in terms of the fate of the Southern Ocean coccolithophores in a future acidified ocean.

Broader impacts: The globally significant size of the Great Belt indicates that it likely plays a major role in global biogeochemistry and climate change feedbacks. Thus, the investigators expect this work to have broad, transformative impacts in biological and chemical oceanography. Ocean acidification from the burning of fossil fuels is predicted to lower the pH of the surface ocean by 0.3 units in the next century and up to 0.7 units - a 5-fold increase in the proton concentration by the year 2300. A major goal of this study is to examine the effects of ocean acidification on coccolithophores in a region of low calcite saturation (i.e., one of the first regions expected to become sub-saturating for calcite). The results of these experiments will therefore be highly relevant to our basic understanding of the marine carbon cycle. Related to career development and Criterion II activities, the project includes field experience on two cruises for NSF REU undergraduates from Maine universities or colleges, providing funds for them to attend a scientific meeting. Participation of undergraduate students from Maine colleges builds capacity in our rural coastal state and helps thwart the serious issue of 'brain drain', in which the best students are leaving Maine to seek opportunity in wealthier, more populated states. A teacher will also participate on the cruises (via the NSF-sponsored ARMADA program). This teacher will develop learning modules for students about such topics as coccolithophores, calcification, export production, metal-plankton interactions, ocean acidification and climate change.

#### **PUBLICATIONS PRODUCED AS A RESULT OF THIS RESEARCH**

Balch, WM; Drapeau, DT; Bowler, BC; Lyczkowski, E; Booth, ES; Alley, D. "The contribution of coccolithophores to the optical and inorganic carbon budgets during the Southern Ocean Gas Exchange Experiment: New evidence in support of the "Great Calcite Belt" hypothesis," *JOURNAL OF GEOPHYSICAL RESEARCH-OCEANS*, v.116, 2011. View record at Web of Science

Poulton, AJ; Young, JR; Bates, NR; Balch, WM. "Biometry of detached *Emiliania huxleyi* coccoliths along the Patagonian Shelf," *MARINE ECOLOGY-PROGRESS SERIES*, v.443, 2011, p. 1. View record at Web of Science

#### **BOOKS/ONE TIME PROCEEDING**

Brown, Michael S, W. Balch, S. Craig, B. Bowler, D. Drapeau, J. Grant. "Optical closure within a Patagonian Shelf coccolithophore bloom", 06/01/2011-05/31/2012, 2012, "ACCESS'12. Atlantic Canada Coastal & Estuarine Science Society. Dalhousie University, Halifax, Nova Scotia. 10-13 May, 2012."

[ [table of contents](#) | [back to top](#) ]

---

#### **Funding**

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0961660</a>

[ [table of contents](#) | [back to top](#) ]