Metadata from sediment samples collected at Paleochori Bay, Milos Island, Greece in 2012 (Hydrothermal Autotrophic Carbon Fixation project)

Website: https://www.bco-dmo.org/dataset/650175
Data Type: Cruise Results
Version:
Version Date: 2016-06-28

Project
» Autotrophic carbon fixation at a shallow-water hydrothermal system: Constraining microbial activity, isotopic and geochemical regimes (Hydrothermal Autotrophic Carbon Fixation)

Program
» Center for Dark Energy Biosphere Investigations (C-DEBI)

Contributors	Affiliation	Role
Vetriani, Costantino	Rutgers University (Rutgers IMCS)	Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
  - Methods & Sampling
- Data Files
- Parameters
- Instruments
- Deployments
- Project Information
- Program Information
- Funding

Dataset Description

List of samples collected at the shallow-water hydrothermal vent sites at Paleochori Bay, Milos Island, Greece, and includes the sample description, collection distance from the vent, type and amount of nucleic acids.

Related Datasets:
GenBank accessions
Microbial isolates

Methods & Sampling

Sequence data (bacterial ssrRNA) was obtained using the Ion Torrent platform from total RNA extracted from 32 sediment and microbial biofilm samples characterized along a transect. Sequence data are currently being generated and analyzed, they will be deposited in GenBank prior to publication and will be made available to the scientific community.

[table of contents | back to top]

Data Files

File
sample_log.csv (Comma Separated Values (.csv), 3.76 KB) MD5:e21c5be16c2b7cde2a5c3941405a2951 Primary data file for dataset ID 650175

[table of contents | back to top]

Parameters

Parameter	Description	Units
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
sample_description	type of sample: from sediment of microbial mat	unitless
sample_id	sample identification	unitless
dist_vent_cm	distance from vent center	cm
sampling_instr	instrument used to collect sample	unitless
preservation	sample preservative	unitless
core_depth	depth within the sediment core	cm
date_collected	date sampled	unitless
nucleic_acid	nucleic acid extracted	unitless
nucleic_acid_conc_ng_ul	nucleic acid concentration	nanograms/microliter
cDNA_conc_ng_ul	cDNA concentration	nanograms/microliter

[table of contents | back to top]

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	Automated DNA Sequencer
Dataset-specific Description	Ion Torrent platform
Generic Instrument Description	A DNA sequencer is an instrument that determines the order of deoxynucleotides in deoxyribonucleic acid sequences.

Dataset-specific Instrument Name	
Generic Instrument Name	Push Corer
Dataset-specific Description	push cores were collected by SCUBA divers
Generic Instrument Description	Capable of being performed in numerous environments, push coring is just as it sounds. Push coring is simply pushing the core barrel (often an aluminum or polycarbonate tube) into the sediment by hand. A push core is useful in that it causes very little disturbance to the more delicate upper layers of a sub-aqueous sediment. Description obtained from: http://web.whoi.edu/coastal-group/about/how-we-work/field-methods/coring/

[table of contents | back to top]

Deployments

Milos_vents_2012

Website	https://www.bco-dmo.org/deployment/474391
Platform	shoreside Milos_vents
Start Date	2012-05-21
End Date	2012-05-30
Description	Vent fluids and sediment cores were were collected at shallow-water hydrothermal vent sites at Paleochori Bay, Milos Island, Greece, and analyzed for their chemical composition (anion, cation, trace elements/metals).

[table of contents | back to top]

Project Information

Autotrophic carbon fixation at a shallow-water hydrothermal system: Constraining microbial activity, isotopic and geochemical regimes (Hydrothermal Autotrophic Carbon Fixation)

Coverage: Shallow-water hydrothermal vents, Paleochori Bay, Milos Island, Greece

In this project we studied the shallow-water hydrothermal vent sites at Milos Island (Greece) to better understand the extent of autotrophic carbon fixation and its chemical and isotopic signature along environmental (redox/thermal) gradients. This was a 12-day long expedition (May 18 to 30, 2012) to sample vent fluids, gases and retrieve sediment cores at Paleochori Bay by using SCUBA diving at 8-10 m depth. In

addition to the submarine vent sites, two subaerial locations of venting were identified at 36o 40' 28"N - 24o 31' 14" E and 36o 40' 25" N - 24o 30' 44" E. Both the subaerial and submarine sites are located on the same fracture zone that likely controls the hydrothermal circulation of evolved meteoritic water and seawater within the magmatic zone of Milos Island. To this end, the geochemistry of the fluids and gases emitted from subaerial sites provide important information towards identifying the linkage between the subaerial and submarine magmatic activity and provide insights on the metabolic functions (e.g. H2 oxidation, Fe(III) reduction, C and S cycling) of the subsurface microbial community.

Abstract:
Currently, there is only limited information on the identity and activity of the microorganisms carrying out CO2-fixation in situ, despite the fact that these organisms form the basis of their respective ecosystems. Representatives that are able to grow autotrophically are known to exist in almost all major groups of prokaryotes, and these organisms play essential roles in ecosystems by providing a continuous supply of organic carbon for heterotrophs. Microorganisms present in extreme environments utilize CO2- fixation pathways other than the Calvin-Benson-Bassham (CBB) cycle. At present, five alternative autotrophic CO2 fixation pathways are known. Different carbon fixation pathways result in distinct isotopic signatures of the produced biomass due to the isotopic discrimination between light (12C) and heavy (13C) carbon by the carboxylating enzymes. Thus, inferences about the carbon fixation pathway predominantly utilized by the microbial community can also be made based on the stable carbon isotopic composition of the organic matter, in extant systems as well as in the geological record. However, at present little is known about the systematics and extents of fractionation during carbon fixation by prokaryotic organisms, and to our knowledge no studies exist that have systematically studied the relationship between the operation of different carbon fixation pathways and how this is reflected in the stable carbon isotopic composition in a natural system. This is a 2-year interdisciplinary, international research program that employs a powerful combination of cutting-edge research tools aiming to improve our understanding of autotrophic carbon fixation and its chemical and isotopic signature along environmental gradients in a natural hydrothermal system. The following hypotheses are addressed:

1. The diversity of microorganisms present along a thermal and redox gradient, and rates of CO2 fixation, will reflect adaptation to in situ temperatures and geochemical conditions

2. Microorganisms utilizing the CBB cycle for autotrophic CO2-fixation will represent a smaller percentage of the chemolithoautotrophic community at higher temperatures, where microorganisms utilizing alternative CO2-fixation pathways dominate

3. Isotopic values of biomass and specific biomarker molecules will vary along a thermal and redox gradient from zones characterized by a higher hydrothermal fluid flux and thus higher temperatures to the surrounding, cooler areas, corresponding to the physiology of the microorganisms utilizing different pathways for carbon fixation

The PIs will use a multidisciplinary approach to delineate the relative contribution of the different carbon fixation pathways along an environmental gradient by combining metagenomic analyses coupled with: 1) an assessment of the frequency and the expression of specific key genes involved in carbon fixation, and 2) with the measurement of carbon fixation rates. These data will be integrated with the determination of stable C isotopic composition of biomass, DIC, and specific hydrocarbons/lipids. Due to its easy accessibility, well-established environmental gradients, and extensive background information, the shallow-water vents off Milos (Greece) will be used as a natural laboratory to perform these studies.

Intellectual Merit. The data generated in this study will allow constraints on the relationship between autotrophic carbon fixation and the resulting isotopic signatures of biomass and specific biomarker molecules (e.g. CH4, C2+ alkanes, lipids) in a natural system. This has implications for assessing the importance of carbon fixation in extant ecosystems, and it will also provide a tool to improve the interpretation of isotopic values in the geological record.

[table of contents | back to top]

Program Information

Center for Dark Energy Biosphere Investigations (C-DEBI)

Website: http://www.darkenergybiosphere.org

Coverage: Global

The mission of the Center for Dark Energy Biosphere Investigations (C-DEBI) is to explore life beneath the seafloor and make transformative discoveries that advance science, benefit society, and inspire people of all ages and origins.

C-DEBI provides a framework for a large, multi-disciplinary group of scientists to pursue fundamental questions about life deep in the sub-surface environment of Earth. The fundamental science questions of C-DEBI involve exploration and discovery, uncovering the processes that constrain the sub-surface biosphere below the oceans, and implications to the Earth system. What type of life exists in this deep biosphere, how much, and how is it distributed and dispersed? What are the physical-chemical conditions that promote or limit life? What are the important oxidation-reduction processes and are they unique or important to humankind? How does this biosphere influence global energy and material cycles, particularly the carbon cycle? Finally, can we discern how such life evolved in geological settings beneath the ocean floor, and how this might relate to ideas about the origin of life on our planet?

C-DEBI's scientific goals are pursued with a combination of approaches:
(1) coordinate, integrate, support, and extend the research associated with four major programs—Juan de Fuca Ridge flank (JdF), South Pacific Gyre (SPG), North Pond (NP), and Dorado Outcrop (DO)—and other field sites;
(2) make substantial investments of resources to support field, laboratory, analytical, and modeling studies of the deep subseafloor ecosystems;
(3) facilitate and encourage synthesis and thematic understanding of submarine microbiological processes, through funding of scientific and technical activities, coordination and hosting of meetings and workshops, and support of (mostly junior) researchers and graduate students; and
(4) entrain, educate, inspire, and mentor an interdisciplinary community of researchers and educators, with an emphasis on undergraduate and graduate students and early-career scientists.

Note: Katrina Edwards was a former PI of C-DEBI; James Cowen is a former co-PI.

Data Management:
C-DEBI is committed to ensuring all the data generated are publically available and deposited in a data repository for long-term storage as stated in their Data Management Plan (PDF) and in compliance with the NSF Ocean Sciences Sample and Data Policy. The data types and products resulting from C-DEBI-supported research include a wide variety of geophysical, geological, geochemical, and biological information, in addition to education and outreach materials, technical documents, and samples. All data and information generated by C-DEBI-supported research projects are required to be made publically available either following publication of research results or within two (2) years of data generation.

To ensure preservation and dissemination of the diverse data-types generated, C-DEBI researchers are working with BCO-DMO Data Managers make data publicly available online. The partnership with BCO-DMO helps ensure that the C-DEBI data are discoverable and available for reuse. Some C-DEBI data is better served by specialized repositories (NCBI's GenBank for sequence data, for example) and, in those cases, BCO-DMO provides dataset documentation (metadata) that includes links to those external repositories.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1124141

[table of contents | back to top]