

# Focal lionfish behaviors at different densities of local lionfish and prey fish in Eleuthera, Bahamas from July to August in 2012

**Website:** <https://www.bco-dmo.org/dataset/653689>

**Data Type:** experimental

**Version:** 1

**Version Date:** 2016-08-10

## Project

» [Mechanisms and Consequences of Fish Biodiversity Loss on Atlantic Coral Reefs Caused by Invasive Pacific Lionfish](#) (BiodiversityLossEffects\_lionfish)

Contributors	Affiliation	Role
<a href="#">Hixon, Mark</a>	University of Hawai'i (UH)	Principal Investigator
<a href="#">Benkwitt, Cassandra E.</a>	Oregon State University (OSU)	Contact
<a href="#">Ake, Hannah</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Focal lionfish behaviors at different densities of local lionfish and prey fish in Eleuthera, Bahamas from July to August in 2012

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Temporal Extent:** 2012-07-16 - 2012-08-30

## Dataset Description

This was an observational field study on natural patch reefs with varying lionfish densities to determine if lionfish behavior and movements change at different local lionfish and prey fish densities. Each reef was visited at three times of day: dawn, midday, and dusk, and two focal lionfish were observed during each visit at each reef. Behaviors of all lionfish were recorded the moment they were first sighted ("Initial behaviors"). One or two focal lionfish were observed for 10 minute intervals and all activity was recorded ("Focal behaviors").

The number of lionfish on each reef that arrived at or departed from each reef to/from the surrounding seagrass habitat was recorded throughout each visit ("Movement"). Censuses of the native prey fish populations were conducted at the end of every visit ("Native fish surveys").

**Related Manuscripts:** Benkwitt, C.E. (2016)

**Species key for all individuals identified in this dataset - DOI:** [10.1575/1912/bco-dmo.655195.2](https://doi.org/10.1575/1912/bco-dmo.655195.2)

## Methods & Sampling

This was an observational field study conducted from June - August 2012 to determine whether lionfish behavior and movements change at different local lionfish and prey fish densities. The study was conducted on sixteen reefs in Rock Sound, Eleuthera, The Bahamas. All reefs were at least 300 m from any reef on which lionfish removals had occurred, and were selected to encompass a range of natural lionfish densities and reef sizes.

A pair of divers visited each reef at three times of day: within 2 hours of sunrise ('dawn'), greater than 3 hours from sunrise or sunset ('midday'), and within 2 hours of sunset ('dusk'). Upon arriving at a reef, observers counted the number of lionfish present by conducting lionfish-focused searches. For each lionfish, observers recorded the size (total length, visually estimated to the nearest cm), behavior, and location the moment it was sighted. Behaviors were categorized as resting (sitting on the substrate, not moving), hovering (in the water column oriented parallel to the bottom, but not moving), swimming (actively moving), or hunting (oriented head down with pectoral fins flared). Location was categorized as the microhabitat on which lionfish were observed (e.g. under a ledge, on top of the reef, in the surrounding seagrass) and later divided into two major categories: sheltering (hidden under structure) or exposed (on top of reef or in surrounding area). Then, 10-minute focal observations were conducted on two randomly-selected lionfish or a single lionfish when there was only one individual present per reef. During focal observations, a trained observer recorded the behavior of lionfish at 30-second intervals for 10 minutes using the same categories as above. The observers also noted any strikes at prey, successful kills, and obviously aggressive interactions (chases, posturing) between lionfish or between lionfish and other species. Throughout the entire visit to each reef, divers noted the time when any lionfish departed from or arrived at the reef and its behavior. A lionfish was defined as departing from the reef if it traveled at least 10 m from the reef. A lionfish was considered arriving at a reef if it swam in from the surrounding areas and had not been previously observed at that reef during that observation period. At the conclusion of the focal observations, the divers re-counted the number of lionfish present while conducting a survey of resident native fishes. Divers recorded the abundance and body size (TL) of all fish 1 - 15 cm TL, native mesopredators that are ecologically similar to lionfish (e.g. *Cephalopholis cruentata* [graysby grouper]), and top predators (e.g. *Epinephelus striatus* [Nassau grouper]) on and within 1 m of the reef.

## Data Processing Description

All data were entered by one person, and then subsequently checked to ensure accuracy. Note that several reefs included in the data file were not included in any analyses because they were too close to other reefs, and/or too small, and/or were not visited at all three time periods: CEI 83 Sat, CEI 97 Oct, CEI Biscuit, CEI Biscuit-2, CEI Biscuit-3, CEI Friendly, CEI Friendly Dat, CEI Friendly Sat 3, and CEI Nicola Sat.

### DMO Notes:

- reformatted column names to comply with BCO-DMO standards
- replaced all spaces with "\_"
- replaced all blank cells with "nd"

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>behaviorMovement_focalBehaviors.csv</b> (Comma Separated Values (.csv), 11.39 KB) MD5:62d4271c311f1cacee5616163448fdef
Primary data file for dataset ID 653689

[ [table of contents](#) | [back to top](#) ]

## Related Publications

Benkwitt, C. (2016). Invasive lionfish increase activity and foraging movements at greater local densities. Marine Ecology Progress Series, 558, 255–266. doi:[10.3354/meps11760](https://doi.org/10.3354/meps11760)  
*General*

## Parameters

Parameter	Description	Units
date	Date that reef was surveyed; mm/dd/yy	unitless
reef	Reef id where lionfish were observed	unitless
time_period	Time of observation; classified as either dawn midday or dusk	unitless
time_start	Time that focal observation began; HH:MM	unitless
time_end	Time that focal observation ended; HH:MM	unitless
PTVO_size	Total length of lionfish	centimeters
specific_location	Location of lionfish within the reef	unitless
prop_rest	Proportion of time focal lionfish spent resting (out of 10 minute observation period)	dimensionless
prop_hover	Proportion of time focal lionfish spent hovering (out of 10 minute observation period)	dimensionless
prop_swim	Proportion of time focal lionfish spent swimming (out of 10 minute observation period)	dimensionless
prop_hunt	Proportion of time focal lionfish spent hunting (out of 10 minute observation period)	dimensionless
departed	Y = yes: lionfish departed from reef during observation; N = no: lionfish did not depart from reef during observation	unitless
returned	If lionfish departed from reef: Y = yes lionfish returned to reef during observation; N = no lionfish did not return to reef during observation	unitless
number_strikes	Number of times lionfish struck at prey (attempts) during observation	count
strike_species	Species that lionfish struck at	unitless

number_blows	Number of times lionfish used the blowing technique during observation	count
blow_species	Species that lionfish blew at	unitless
number_kills	Number of times lionfish successfully captured prey during observation	count
kill_species	Species that lionfish captured	unitless
notes	Notes on observations	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### Eleuthera\_Reef\_Surveys\_2012

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/59028">https://www.bco-dmo.org/deployment/59028</a>
<b>Platform</b>	Cape_Eleuthera_Reefs
<b>Start Date</b>	2012-07-03
<b>End Date</b>	2012-08-28
<b>Description</b>	Reefs were surveyed near the Cape Eleuthera Institute, Eleuthera Bahamas during the summer of 2012 as part of the project "Ecological Release and Resistance at Sea: Invasion of Atlantic Coral Reefs by Pacific Lionfish" and "Mechanisms and Consequences of Fish Biodiversity Loss on Atlantic Coral Reefs Caused by Invasive Pacific Lionfish" (NSF OCE-0851162 & OCE-1233027).

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Mechanisms and Consequences of Fish Biodiversity Loss on Atlantic Coral Reefs Caused by Invasive Pacific Lionfish (BiodiversityLossEffects\_lionfish)

**Website:** <http://hixon.science.oregonstate.edu/content/highlight-lionfish-invasion>

**Coverage:** Three Bahamian sites: 24.8318, -076.3299; 23.8562, -076.2250; 23.7727, -076.1071; Caribbean Netherlands: 12.1599, -068.2820

The Pacific red lionfish (*Pterois volitans*), a popular aquarium fish, was introduced to the Atlantic Ocean in the vicinity of Florida in the late 20th century. Voraciously consuming small native coral-reef fishes, including the juveniles of fisheries and ecologically important species, the invader has undergone a population explosion that now ranges from the U.S. southeastern seaboard to the Gulf of Mexico and across the greater Caribbean region. The PI's past research determined that invasive lionfish (1) have escaped their natural enemies in the Pacific (lionfish are much less abundant in their native range); (2) are not yet controlled by Atlantic predators, competitors, or parasites; (3) have strong negative effects on populations of native Atlantic fishes; and (4) locally reduce the diversity (number of species) of native fishes. The lionfish invasion has been recognized as one of the major conservation threats worldwide.

The Bahamas support the highest abundances of invasive lionfish globally. This system thus provides an

unprecedented opportunity to understand the direct and indirect effects of a major invader on a diverse community, as well as the underlying causative mechanisms. The PI will focus on five related questions: (1) How does long-term predation by lionfish alter the structure of native reef-fish communities? (2) How does lionfish predation destabilize native prey population dynamics, possibly causing local extinctions? (3) Is there a lionfish-herbivore-seaweed trophic cascade on invaded reefs? (4) How do lionfish modify cleaning mutualisms on invaded reefs? (5) Are lionfish reaching densities where natural population limits are evident?

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1233027</a>

[ [table of contents](#) | [back to top](#) ]