

Dates of sea ice movement and sea ice distance in McMurdo Sound, Antarctica from MODIS and SSM/I imagery between 1978-2015

Website: <https://www.bco-dmo.org/dataset/674819>

Data Type: Other Field Results

Version: 2

Version Date: 2020-02-20

Project

» [Food web dynamics in an intact ecosystem: the role of top predators in McMurdo Sound](#) (McMurdo Predator Prey)

Contributors	Affiliation	Role
Kim, Stacy	Moss Landing Marine Laboratories (MLML)	Chief Scientist, Principal Investigator, Contact
Ainley, David G.	H.T. Harvey & Associates	Co-Principal Investigator
Ballard, Grant	Point Blue Conservation Science	Co-Principal Investigator
Daly, Kendra L.	University of South Florida (USF)	Co-Principal Investigator
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

This dataset contains dates relevant to sea ice movement including dates of initial ice retreat, final ice retreat, ice minimum, general ice minimum, ice minimum for the west of McMurdo Sound, ice minimum for the east of McMurdo Sound, and refreeze start. It also includes the minimum distance to the ice edge from McMurdo Station. Dates were derived from Moderate-resolution Imaging Spectroradiometers (MODIS) collected between 2003 and 2015 and Scanning Multichannel Microwave Radiometer and Special Sensor Microwave Imager-family passive microwave sensors (SSM/I) imagery collected between 1978 and 2015.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:-76 E:166 S:-78.5 W:163

Temporal Extent: 1978-01-01 - 2015-01-01

Dataset Description

This dataset contains dates relevant to sea ice movement including dates of initial ice retreat, final ice retreat, ice minimum, general ice minimum, ice minimum for the west of McMurdo Sound, ice minimum for the east of

McMurdo Sound, and refreeze start. It also includes the minimum distance to the ice edge from McMurdo Station.

Dates were derived from Moderate-resolution Imaging Spectroradiometers (MODIS) collected between 2003 and 2015 and Scanning Multichannel Microwave Radiometer and Special Sensor Microwave Imager-family passive microwave sensors (SSM/I) imagery collected between 1978 and 2015.

Methods & Sampling

MODIS:

Dates of fast ice movement were determined from visible-wavelength scenes from the MODIS aboard the Aqua and Terra satellites. MODIS images are available only for the months when there is sunlight, generally October to April, and image resolution was 250 m. Figure A (see Supplemental Documents) shows the division of McMurdo Sound into six zones; fast ice areas were generated manually from clear-sky images by drawing polygons in GIS software, and pack ice was excluded from analysis. The fast ice in MODIS scenes was sometimes obscured by clouds, so for days with missing imagery we interpolated linearly between valid data. Areas of persistent multiyear ice or glaciers that were not considered in the analysis (white in the figure). Linear distance between McMurdo Station and the nearest open water on the day the ice reached the minimum extent for the season was also measured.

SSM/I:

In the SSM/I record, images were available for all months of the year, and image resolution was 25 km. Figure B (see Supplemental Documents) shows the total of seven pixels in the 25 km polar stereographic projection that encompassed McMurdo Sound; for each pixel, daily or bi-daily fractional sea ice cover was extracted. SSM/I passive microwave sea ice observations do not distinguish between fast ice and pack ice. There are small amounts of land and ice shelf contamination in pixels 1, 2, 5 and 7.

Data Processing Description

The value "notobs" indicates the phenomena was not observed for two possible reasons; either the phenomena did not occur that season or imagery was not available (e.g. MODIS visual imagery was only collected during daylight months). The value 'nd' indicates no data (e.g. SSM/I imagery was too coarse to distinguish east and west side minimums).

SSM/I:

Data were processed using the bootstrap algorithm (National Snow and Ice Data Center, Comiso, 2000, Cavalieri et al. 2015) for the 1978/79-2014/15 seasons.

Comiso, J (2000, updated 2015) Bootstrap Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS. Version 2, 1979-2015. Boulder, Colorado USA: NASA DAAC at the National Snow and Ice Data Center.

Cavalieri DJ, Parkinson CL, Gloersen P, and Zwally HJ (2015) Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, Version 1.1 Southern Hemisphere, 1978-2015. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. (doi:10.5067/8GQ8LZQVL0VL) Accessed 11 Dec 2015.

BCO-DMO Data Manager Processing Notes:

- * added a conventional header with dataset name, PI name, version date
- * modified parameter names to conform with BCO-DMO naming conventions
- * blank values indicating no observed phenomena replaced with no data value 'nd'
- * added approximate latitude and longitude of McMurdo Station

Data version 2 (2020-02-20)

- * Data from data version 1 (2017-01-31) updated. Column values for "edge_dist_min, ice_min_west_year, ice_min_west_month" switched with "refreeze_begin_year, refreeze_begin_month, refreeze_begin_day" values where image_source was SSMI.

Data Files

File
IceEdge.csv (Comma Separated Values (.csv), 5.27 KB) MD5:c915ccf065f600927ba06067976a6e06 Primary data file for dataset ID 674819

Supplemental Files

File
Map of sea ice and MODIS/SSM/I survey areas in McMurdo Sound filename: MODIS.jpg (JPEG Image (.jpg), 128.41 KB) MD5:c378921f0ae7b87aec54b54c64a88d26 Figure A shows the division of McMurdo Sound into six zones; fast ice areas were generated manually from clear-sky images by drawing polygons in GIS software, and pack ice was excluded from analysis. The fast ice in MODIS scenes was sometimes obscured by clouds, so for days with missing imagery we interpolated linearly between valid data. Areas of persistent multiyear ice or glaciers that were not considered in the analysis (white in the figure). Figure B shows the total of seven pixels in the 25 km polar stereographic projection that encompassed McMurdo Sound; for each pixel, daily or bi-daily fractional sea ice cover was extracted. SSM/I passive microwave sea ice observations do not distinguish between fast ice and pack ice. There are small amounts of land and ice shelf contamination in pixels 1, 2, 5 and 7.

Parameters

Parameter	Description	Units
lat_approx	Approximate latitude of sampling near McMurdo Station; Antarctica; south is negative	decimal degrees
lon_approx	Approximate longitude of sampling near McMurdo Station; Antarctica; west is negative	decimal degrees
image_source	Image source (MODIS or SSM/I)	unitless
austral_year	Austral Year (start year/end year) in format yyyy/yy	unitless
retreat_initial_year	Year of initial ice retreat	unitless
retreat_initial_month	Month of initial ice retreat	unitless
retreat_initial_day	Day of initial ice retreat	unitless

retreat_final_year	Year of final ice retreat	unitless
retreat_final_month	Month of final ice retreat	unitless
retreat_final_day	Day of final ice retreat	unitless
ice_min_year	Year ice reached minimum in McMurdo Sound	unitless
ice_min_month	Month ice reached minimum in McMurdo Sound	unitless
ice_min_day	Day ice reached minimum in McMurdo Sound	unitless
edge_dist_min	Linear distance between McMurdo Station and the nearest open water on the day the ice reached the minimum extent for the season.	kilometers
ice_min_west_year	Year ice reached minimum on west side of McMurdo Sound	unitless
ice_min_west_month	Month ice reached minimum on west side of McMurdo Sound	unitless
ice_min_west_day	Day ice reached minimum on west side of McMurdo Sound	unitless
ice_min_east_year	Year ice reached minimum on east side of McMurdo Sound	unitless
ice_min_east_month	Month ice reached minimum on east side of McMurdo Sound	unitless
ice_min_east_day	Day ice reached minimum on east side of McMurdo Sound	unitless
refreeze_begin_year	Year ice began to refreeze	unitless
refreeze_begin_month	Month ice began to refreeze	unitless
refreeze_begin_day	Day ice began to refreeze	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Scanning Multichannel Microwave Radiometer
Generic Instrument Name	Radiometer
Dataset-specific Description	Scanning Multichannel Microwave Radiometer
Generic Instrument Description	Radiometer is a generic term for a range of instruments used to measure electromagnetic radiation (radiance and irradiance) in the atmosphere or the water column. For example, this instrument category includes free-fall spectral radiometer (SPMR/SMSR System, Satlantic, Inc), profiling or deck cosine PAR units (PUV-500 and 510, Biospherical Instruments, Inc). This is a generic term used when specific type, make and model were not specified.

Dataset-specific Instrument Name	MODIS
Generic Instrument Name	Radiometer
Dataset-specific Description	Moderate-resolution Imaging Spectroradiometers (MODIS)
Generic Instrument Description	Radiometer is a generic term for a range of instruments used to measure electromagnetic radiation (radiance and irradiance) in the atmosphere or the water column. For example, this instrument category includes free-fall spectral radiometer (SPMR/SMSR System, Satlantic, Inc), profiling or deck cosine PAR units (PUV-500 and 510, Biospherical Instruments, Inc). This is a generic term used when specific type, make and model were not specified.

Dataset-specific Instrument Name	SSM/I
Generic Instrument Name	Special Sensor Microwave Imager
Dataset-specific Description	Special Sensor Microwave Imager-family passive microwave sensors (SSM/I)
Generic Instrument Description	<p>The Special Sensor Microwave Imager (SSM/I) is a passive microwave radiometer that measures atmospheric, oceanic and terrestrial microwave brightness temperatures at 19.35, 22.2, 37.0, and 85.5 GHz, from which ocean surface wind speed, atmospheric water vapour, cloud liquid water and rain rate can be derived. The SSM/I consists of an offset parabolic reflector of dimensions 24 x 26 inches, fed by a corrugated, broad-band, seven-port horn antenna. The reflector and feed are mounted on a drum that contains the radiometers, digital data subsystem, mechanical scanning subsystem, and power subsystem. The reflector-feed-drum assembly is rotated about the axis of the drum by a coaxially mounted bearing and power transfer assembly (BAPTA). All data, commands, timing and telemetry signals, and power pass through the BAPTA on slip ring connectors to the rotating assembly. The absolute brightness temperature of the scene incident upon the antenna is received and spatially filtered by the antenna to produce an effective input signal or antenna temperature at the input of the feed horn antenna. The SSM/I has seven channels, four frequencies, and is orthogonally polarised. It rotates continuously about an axis parallel to the local spacecraft at 31.6 rpm and measures the upwelling scene brightness temperature over an angular range of 102.4 degrees at nadir. The active scene measurements lie +51.2 to -51.2 degrees about the forward (F10, F11) or aft (F8) direction. This results in a swath width of approximately 1400 kilometres. During each 1.9 second scan, 128 discrete uniformly spaced radiometric samples are taken at the two 85 GHz channels and, on alternate scans, 64 discrete samples are taken at the remaining lower frequency channels. The antenna beam intersects the Earth's surface at an incidence angle of 53.1 degrees. Flown on F08, F10, F11, F13, F14, F15, F16, F17, F18 F19.</p> <p>https://nsidc.org/data/docs/daac/ssmi_instrument.gd.html</p>

[[table of contents](#) | [back to top](#)]

Deployments

TERRA_McMurdo_observations

Website	https://www.bco-dmo.org/deployment/679890
Platform	TERRA
Start Date	2003-01-09
End Date	2015-02-10

AQUA_McMurdo_observations

Website	https://www.bco-dmo.org/deployment/679894
Platform	AQUA
Start Date	2003-02-10
End Date	2015-02-10

[[table of contents](#) | [back to top](#)]

Project Information

Food web dynamics in an intact ecosystem: the role of top predators in McMurdo Sound (McMurdo Predator Prey)

Website: <https://scini-penguin.mlml.calstate.edu/pauls-wordpress-test-site/>


Coverage: McMurdo Sound, Antarctica

Extracted from the NSF award abstract:

The research project investigates the importance of top down forcing on pelagic food webs. The relatively pristine Ross Sea includes large populations of upper-level predators such as minke and killer whales, Adélie and Emperor penguins, and Antarctic toothfish. This project focuses on food web interactions of Adélie penguins, minke whales, and the fish-eating Ross Sea killer whales, all of which exert foraging pressure on their main prey, crystal krill (*Euphausia cyrstorophias*) and silver fish (*Pleuragramma antarcticum*) in McMurdo Sound.

The investigators used a video- and acoustic-capable ROV, and standard biological and environmental sensors to quantify the abundance and distribution of phytoplankton, sea ice biota, prey, and relevant habitat data. The sampling area included 37 stations across an 30 x 15 km section of McMurdo Sound, stratified by distance from the ice edge as a proxy for air-breathing predator access. This study will be among the first to assess top-down forcing in the Ross Sea ecosystem and will form the basis for multidisciplinary studies in the future.

Map sampling stations



[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Polar Programs (NSF PLR)	PLR-0944747
NSF Division of Polar Programs (NSF PLR)	PLR-0944511
NSF Division of Polar Programs (NSF PLR)	PLR-0944694

[[table of contents](#) | [back to top](#)]