

Pleuromamma xiphias mitochondrial genetic data (Atlantic; mtDNA) for analyses of population structure across the Atlantic Ocean

Website: <https://www.bco-dmo.org/dataset/699440>

Data Type: Cruise Results

Version: 1

Version Date: 2017-05-04

Project

» [Basin-scale genetics of marine zooplankton](#) (Plankton Population Genetics)

Contributors	Affiliation	Role
Goetze, Erica	University of Hawai'i at Mānoa (SOEST)	Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:39.6470388 E:-22.4662575 S:-40.0732082 W:-42.3712545

Temporal Extent: 2012-10-16 - 2012-11-17

Dataset Description

Specimens were collected by bulk plankton tows on the Atlantic Meridional Transect Cruise 22 (AMT22), on RRS James Cook from Oct-Nov 2012.

Data files associated with the publication:

Goetze, E., Hüdepohl, P., Chang, C., Iacchei, M., Van Woudenberg, L., Peijnenburg, K. T. C. A. (2016) Ecological dispersal barrier across the equatorial Atlantic in a migratory planktonic copepod. Progress in Oceanography – AMT special issue. doi:[10.1016/j.pocean.2016.07.001](https://doi.org/10.1016/j.pocean.2016.07.001)

The data files included here were used in all population genetic analyses, which only included stations with sample sizes N > 40. Data files include:

- (1) a README file describing the other files;
- (2) A fasta file with all unique haplotypes (Goetze_et_al_2016_all_haplotypes.fasta);
- (3) An infile formatted for use in Arlequin (<http://cmpg.unibe.ch/software/arlequin35/>); (Goetze_et_al_2016_plxi_mtCOI.arp);
- (4) A population report file that describes the mtCOI haplotype composition of each sample (Goetze_et_al_2016_pop_report.html).

Note: In addition to these files, mtCOI sequences representing unique haplotypes from this study are available under GenBank accession numbers KT429028 - KT429159.

Refer to related dataset ["AMT22 Population Samples"](#) for information on population samples included in genetic analyses.

Methods & Sampling

Refer to the following publication for complete methodology details:

Goetze, E., Hüdepohl, P., Chang, C., Iacchei, M., Van Woudenberg, L., Peijnenburg, K. T. C. A. (2016) Ecological dispersal barrier across the equatorial Atlantic in a migratory planktonic copepod. *Progress in Oceanography* – AMT special issue. doi: [10.1016/j.pocean.2016.07.001](https://doi.org/10.1016/j.pocean.2016.07.001)

In summary (excerpted from above):

Bulk plankton samples were collected on Atlantic Meridional Transect Cruise 22 (AMT22) between 10/13/2012 and 11/19/2012. Oblique tows were conducted with bongo nets (200 µm, 333 µm), towed between on average 324 m depth and the sea surface. A General Oceanics flowmeter (2030RC) mounted in the mouth of the 200 µm net was used to measure seawater filtered during the tow. Plankton from the 200 µm mesh net was bulk preserved immediately in 100% ethyl alcohol, the alcohol was changed to fresh within 12–24 h of collection, and samples were stored at -20 °C. Plankton from the 333 µm mesh net was sorted live at sea, and *Pleuromamma xiphius* specimens were preserved immediately in RNALater (Ambion), followed by cryopreservation in liquid nitrogen, and long-term storage at -80 °C.

Specimens included in the genetic analyses in this study were collected at 18 stations, located between 39° 38.82'N and 40° 4.39'S latitude. The majority of genetic analyses focused on stations with sufficient sample size for population-level inference ($N > 42$), including AMT22-09 through AMT22-29 and AMT22-45 through AMT22-68. Specimens used in genetic analyses were primarily RNALater-preserved, but some specimens were included from ethanol-preserved samples to achieve the minimum target sample size of 45 individuals per station. When a major genetic break across the equatorial region was identified, we included samples from stations AMT22-31 through AMT22-43 to assess the genetic composition of populations across this region. Population-level analyses were not conducted on these latter stations due to small sample sizes and low abundance in this region, but sequences from these animals were included in the haplotype network. DNA was extracted from individual *P. xiphius* adults using the DNeasy Blood & Tissue kit (Qiagen), following the manufacturer's protocol, with the exception of longer elution incubation times (Goetze, 2011). The second of two elutions for each individual was used in this study. Polymerase Chain Reaction (PCR) amplification of a 681-bp fragment of mtCOI was conducted with primers and PCR and sequencing protocols as described in (Goetze, 2011). Forward and reverse sequences from each individual were aligned and checked for errors in Geneious (v7.1.8, Biomatters). Consensus sequences for all individuals were aligned using MUSCLE (Edgar, 2004), and unique mtCOI haplotypes were identified using FABox (<http://users-birc.au.dk/biopv/php/fabox/>). MtCOI sequences representing unique haplotypes are available under GenBank accession numbers KT429028–KT429159. A minimum spanning haplotype network was inferred for all mtCOI sequences using Population Analysis with Reticulate Trees (PopART; <http://popart.otago.ac.nz>), in order to investigate geographic patterns in the distribution of haplotypes across the Atlantic.

Population subdivision across samples was investigated using several approaches. Refer to publication cited above.

[[table of contents](#) | [back to top](#)]

Data Files

File	
Goetze_et_al_2016_all_haplotypes.fasta	(FASTA, 72.86 KB) MD5:b2c088f99506cab0107bb118f914dff1
A fasta file with all haplotypes	
Goetze_et_al_2016_pxi_mtCOI.arp	(Octet Stream, 122.93 KB) MD5:1fe9082aaff284e648f21ab9a6720420
An arlequin infile	
Goetze_et_al_2016_pop_report.html	(HTML, 9.87 KB) MD5:b23e542fba5e068f446316eb3011bd41
A population report file that describes the mtCOI haplotype composition of each sample	

[[table of contents](#) | [back to top](#)]

Supplemental Files

File	
README.txt	(Plain Text, 1.39 KB) MD5:f40cff261f65582c8820313207f48a8c
Readme file describing the other files	

[[table of contents](#) | [back to top](#)]

Related Publications

Goetze, E., Hüdepohl, P. T., Chang, C., Van Woudenberg, L., Iacchei, M., & Peijnenburg, K. T. C. A. (2017). Ecological dispersal barrier across the equatorial Atlantic in a migratory planktonic copepod. Progress in Oceanography, 158, 203–212. doi:[10.1016/j.pocean.2016.07.001](https://doi.org/10.1016/j.pocean.2016.07.001)
Results

[[table of contents](#) | [back to top](#)]

Related Datasets

Different Version

Goetze, E., Hüdepohl, P. T., Chang, C., Van Woudenberg, L., Iacchei, M., & Peijnenburg, K. T. C. A. (2017). Data from: Ecological dispersal barrier across the equatorial Atlantic in a migratory planktonic copepod [Data set]. Dryad Digital Repository. <https://doi.org/10.5061/dryad.s058r>

[[table of contents](#) | [back to top](#)]

Parameters

Parameters for this dataset have not yet been identified

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	bongo net
Generic Instrument Name	Bongo Net
Dataset-specific Description	Oblique tows were conducted with bongo nets (200 um, 333 um), towed between on average 324 m depth and the sea surface.
Generic Instrument Description	A Bongo Net consists of paired plankton nets, typically with a 60 cm diameter mouth opening and varying mesh sizes, 10 to 1000 micron. The Bongo Frame was designed by the National Marine Fisheries Service for use in the MARMAP program. It consists of two cylindrical collars connected with a yoke so that replicate samples are collected at the same time. Variations in models are designed for either vertical hauls (OI-2500 = NMFS Pairovet-Style, MARMAP Bongo, CalVET) or both oblique and vertical hauls (Aquatic Research). The OI-1200 has an opening and closing mechanism that allows discrete "known-depth" sampling. This model is large enough to filter water at the rate of 47.5 m ³ /minute when towing at a speed of two knots. More information: Ocean Instruments, Aquatic Research, Sea-Gear

Dataset-specific Instrument Name	PCR
Generic Instrument Name	Thermal Cycler
Dataset-specific Description	Polymerase Chain Reaction (PCR) amplification of a 681-bp fragment of mtCOI was conducted with primers and PCR and sequencing protocols as described in (Goetze, 2011).
Generic Instrument Description	A thermal cycler or "thermocycler" is a general term for a type of laboratory apparatus, commonly used for performing polymerase chain reaction (PCR), that is capable of repeatedly altering and maintaining specific temperatures for defined periods of time. The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. They can also be used to facilitate other temperature-sensitive reactions, including restriction enzyme digestion or rapid diagnostics. (adapted from http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html)

[[table of contents](#) | [back to top](#)]

Deployments

JC079

Website	https://www.bco-dmo.org/deployment/540458
Platform	RRS James Cook
Report	http://dmoserv3.whoi.edu/data_docs/Goetze/AMT22_cruise/jc079.pdf
Start Date	2012-10-10
End Date	2012-11-24
Description	The AMT22 cruise set sail from Southampton in the UK on 10 October 2012 and arrived in Punta Arenas, Chile on 24 November 2012. The final cruise report and other cruise information, including all science components, can be found online at the Atlantic Meridional Transect webpage (http://www.amt-uk.org/Cruises), or through the British Oceanographic Data Centre (BODC) (http://www.bodc.ac.uk/projects/uk/amt/). Zooplankton ecology data from the project "Does habitat specialization drive population genetic structure of oceanic zooplankton?" (NSF OCE-1029478) were collected on this cruise.

[[table of contents](#) | [back to top](#)]

Project Information

Basin-scale genetics of marine zooplankton (Plankton Population Genetics)

Coverage: Atlantic Ocean, 46 N - 46 S

Description from NSF award abstract:

Marine zooplankton show strong ecological responses to climate change, but little is known about their capacity for evolutionary response. Many authors have assumed that the evolutionary potential of zooplankton is limited. However, recent studies provide circumstantial evidence for the idea that selection is a dominant evolutionary force acting on these species, and that genetic isolation can be achieved at regional spatial scales in pelagic habitats. This RAPID project will take advantage of a unique opportunity for basin-scale transect sampling through participation in the Atlantic Meridional Transect (AMT) cruise in 2014. The cruise will traverse more than 90 degrees of latitude in the Atlantic Ocean and include boreal-temperate, subtropical and tropical waters. Zooplankton samples will be collected along the transect, and mitochondrial and microsatellite markers will be used to identify the geographic location of strong genetic breaks within three copepod species. Bayesian and coalescent analytical techniques will test if these regions act as dispersal barriers. The physiological condition of animals collected in distinct ocean habitats will be assessed by measurements of egg production (at sea) as well as body size (condition index), dry weight, and carbon and nitrogen content. The PI will test the prediction that ocean regions that serve as dispersal barriers for marine holoplankton are areas of poor-quality habitat for the target species, and that this is a dominant mechanism driving population genetic structure in oceanic zooplankton.

Note: This project is funded by an NSF RAPID award. This RAPID grant supported the shiptime costs, and all the sampling reported in the [AMT24 zooplankton ecology cruise report \(PDF\)](#).

Online science outreach blog at: <https://atlanticplankton.wordpress.com>

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1338959
NSF Division of Ocean Sciences (NSF OCE)	OCE-1029478

[[table of contents](#) | [back to top](#)]