

FASTA file of protein identifications from R/V Kilo Moana KM1128 in the North Pacific during 2011 (MetZyme project)

Website: <https://www.bco-dmo.org/dataset/708816>

Data Type: Cruise Results

Version: 1

Version Date: 2017-07-14

Project

» [Connecting Trace Elements and Metalloenzymes Across Marine Biogeochemical Gradients \(GPc03\)](#)
(MetZyme)

Program

» [U.S. GEOTRACES](#) (U.S. GEOTRACES)

Contributors	Affiliation	Role
Saito, Mak A.	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
Ake, Hannah	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

FASTA file of protein identifications from the KM1128 in 2011. There is one entry for every protein ID.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [BCO-DMO Processing Description](#)
 - [Data Files](#)
 - [Related Publications](#)
 - [Related Datasets](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Location: Tropical North Pacific

Spatial Extent: N:21.4574 E:-151.3409 S:-17.0523 W:-177.7081

Temporal Extent: 2011-10 - 2018-10

Dataset Description

These data are part of the Ocean Protein Portal "MetZyme 0.2" dataset version 5 (<https://proteinportal.whoi.edu/>; Saito et al., 2020).

Methods & Sampling

Ocean metaproteomes were sampled and extracted as described in Saito et al., 2014. The protocols are reproduced here:

Samples were collected during the KM1128 METZYME research expedition (Metals and Enzymes in the Pacific) on the R/V Kilo Moana October 1-25, 2011 from Oahu, Hawaii to Apia, Samoa, with Carl Lamborg and Mak Saito as Chief Scientists. Microbial biomass for protein analyses was collected on vertical profiles using in situ high volume particle filtration pumps with a focus on the North Pacific and Equatorial regions. Protein samples were collected by a suite of 4 L/min and 8 L/min McLane Pumps (WTS-LV; McLane Research Laboratories Inc., Falmouth MA, USA) outfitted with custom Mini-MULVS multiple filter head systems. The 0.2-3.0 micron size fraction was collected on 142 mm filters (Supor, Pall Corp.) for analyses used in this study. The volume of water was pumped until a minimal flow rate was achieved or the allotted cast time period expired, typically ~300-500 L. Filters were sectioned immediately after pump retrieval, and protein samples (1/4 filter) were stored in RNeasy lysis reagent (Qiagen, Life Technologies), which has been shown to be an effective preservative for cyanobacterial biomass (Saito et al., 2011), frozen at -80C, transported back to the laboratory on dry ice, and stored at -80C until analysis.

Total microbial protein (0.2-3.0µm fraction) was extracted using detergent-based methods, described below. Total protein showed enhanced concentrations in the photic zone, particularly in the Equatorial and South Pacific portions of the transect (Fig. S1F). For protein extraction, samples were thawed and the filter and RNeasy lysis reagent (Qiagen Life Technologies) were separated. The removed preservative was spin-concentrated by a 5K MWCO membrane (Sartorius Stedim Biotech 6 mL, 5 K MWCO Vivaspins units; Goettingen, Germany), and rinsed with 0.1M Tris buffer to recover, desalt, and concentrate any suspended material. The sample filter was unfolded and placed in a larger tube to which 1% SDS extraction buffer (1% SDS, 0.1M Tris/HCL pH 7.5, 10mM EDTA) and the rinsed/desalted RNeasy lysis fraction was added back. Each sample was incubated at room temperature for 15 minutes, heated at 95C for 10 minutes, and shaken at room temperature for 350 rpm for 1 h. The protein extract was decanted and placed in a new tube and centrifuged for 30 min x 3220 g at room temperature. The supernatants were removed and filtered through a 5 µm low protein binding syringe filter (Fisher Scientific), and the filter rinsed with extraction buffer. The extracts were concentrated by 5 kD membrane centrifugation to a small volume, washed with extraction buffer, and concentrated again. Each sample was precipitated with cold 50% methanol (MeOH) 50% acetone 0.5 mM HCL for 3 days at -20C, centrifuged at 14100 x g (14500 rpm) for 30 min at 4C, decanted and dried by vacuum concentration (Thermo Savant Speedvac) for 10 min or until dry. Pellets were resuspended in 1% SDS extraction buffer and left at room temperature (RT) for 1 h to redissolve. Total protein was quantified (Bio-Rad DC protein assay, Hercules, CA) with BSA as a standard.

Extracted proteins were purified from SDS detergent, reduced, alkylated, and trypsin digested, while embedded within a polyacrylamide tube gel, using a modified protocol from a previously published method (38). The tube gel approach allowed all proteins including membrane proteins to be solubilized by detergent and purified while immobilized in the gel matrix. A gel premix was made by combining 1 M Tris HCL (pH 7.5) and 40% Bis-acrylamide L 29:1 (Acros Organics) at a ratio of 1:3. The premix (103 µl) was combined with an extracted protein sample (usually 25 µg-200 µg), TE, 7 µl 1% APS and 3 µl of TEMED (Acros Organics) to a final volume of 200 µl. After 1 h of polymerization at RT, 200 µl of gel fix solution (50% ETOH, 10% acetic acid in LC/MS grade water) was added to the top of the gel and incubated at RT for 20 minutes. Liquid was then removed and the tube gel was transferred into a new 1.5 mL microtube.

BCO-DMO Processing Description

Version 1:

- * Originally submitted file added to BCO-DMO dataset. No processing performed.
- * Related datasets with peptide and protein spectral counts were iterated to version 5 at both BCO-DMO and the Ocean Protein Portal (OPP). However, v5 at the OPP "MetZyme 0.2" used this original (v1) fasta, it was not changed between OPP "MetZyme 0.2" versions 1-5.

[[table of contents](#) | [back to top](#)]

Data Files

File	
Metzyme_Fusion_AllStations_precursors.fasta	(FASTA, 7.29 MB)
MD5:e5962c00ace4b07d126ec71328940130	
FASTA file of protein identifications. This is the primary data file for dataset 708816.	

Related Publications

Saito, M. A., Bertrand, E. M., Duffy, M. E., Gaylord, D. A., Held, N. A., Hervey, W. J., Hettich, R. L., Jagtap, P. D., Janech, M. G., Kinkade, D. B., Leary, D. H., McIlvin, M. R., Moore, E. K., Morris, R. M., Neely, B. A., Nunn, B. L., Saunders, J. K., Shepherd, A. I., Symmonds, N. I., & Walsh, D. A. (2019). Progress and Challenges in Ocean Metaproteomics and Proposed Best Practices for Data Sharing. *Journal of Proteome Research*, 18(4), 1461–1476. <https://doi.org/10.1021/acs.jproteome.8b00761>
Methods

Saito, M. A., Saunders, J. K., Chagnon, M., Gaylord, D. A., Shepherd, A., Held, N. A., Dupont, C., Symmonds, N., York, A., Charron, M., & Kinkade, D. B. (2020). Development of an Ocean Protein Portal for Interactive Discovery and Education. *Journal of Proteome Research*, 20(1), 326–336.
<https://doi.org/10.1021/acs.jproteome.0c00382>
Related Research

Related Datasets

IsRelatedTo

Saito, M. (2019) **Station locations for the global metaproteomic dataset from R/V Kilo Moana cruise KM1128, the METZYME expedition in the tropical North Pacific in 2011.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 3) Version Date 2019-04-12
doi:10.1575/1912/bco-dmo.730891.3 [view at BCO-DMO]
Relationship Description: Station locations and sampling information for these data.

Saito, M. A. (2022) **Peptides and their spectral counts from KM1128 the METZYME expedition on R/V Kilo Moana in the tropical North Pacific in 2011.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 5) Version Date 2022-06-03 <http://lod.bco-dmo.org/id/dataset/731314> [view at BCO-DMO]
Relationship Description: Related peptides and their spectral counts. This dataset is part of the same "MetZyme 0.2" dataset at the Ocean Protein Portal (<https://proteinportal.whoi.edu/>).

Saito, M. A. (2022) **Total spectral count of proteins from R/V Kilo Moana cruise KM1128 for the METZYME expedition in the tropical North Pacific in 2011.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 5) Version Date 2022-06-06 <http://lod.bco-dmo.org/id/dataset/730722> [view at BCO-DMO]
Relationship Description: This dataset is part of the same "MetZyme 0.2" dataset at the Ocean Protein Portal (<https://proteinportal.whoi.edu/>).

IsPartOf

Protein Portal (2024). Ocean Protein Portal. Available from <https://proteinportal.whoi.edu/>

Parameters

Parameter	Description	Units
sequence	Precursors from all METZYME stations	unitless
scaffold_id	Links to the scaffold_id in the protein spectral counts dataset	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	5 kD membrane centrifuge
Generic Instrument Name	Centrifuge
Generic Instrument Description	A machine with a rapidly rotating container that applies centrifugal force to its contents, typically to separate fluids of different densities (e.g., cream from milk) or liquids from solids.

Dataset-specific Instrument Name	
Generic Instrument Name	McLane Pump
Generic Instrument Description	McLane pumps sample large volumes of seawater at depth. They are attached to a wire and lowered to different depths in the ocean. As the water is pumped through the filter, particles suspended in the ocean are collected on the filters. The pumps are then retrieved and the contents of the filters are analyzed in a lab.

[[table of contents](#) | [back to top](#)]

Deployments

KM1128

Website	https://www.bco-dmo.org/deployment/59053
Platform	R/V Kilo Moana
Start Date	2011-10-01
End Date	2011-10-25
Description	This is a MetZyme project cruise. The original cruise data are available from the NSF R2R data catalog.

[[table of contents](#) | [back to top](#)]

Project Information

Connecting Trace Elements and Metalloenzymes Across Marine Biogeochemical Gradients (GPc03) (MetZyme)

Coverage: Tropical North Pacific along 150 degrees West from 18 degrees North to the equator

MetZyme project researchers will determine the role of enzymatic activity in the cycling of trace metals. Specifically the research will address the following questions: (1) degradation of sinking particulate organic

material in the Tropical North Pacific can be influenced by the ability of microbes to synthesize zinc proteases, which in turn is controlled by the abundance or availability of zinc, and (2) methylation of mercury is controlled, in part, by the activity of cobalt-containing enzymes, and therefore the supply of labile cobalt to the corrinoid-containing enzymes or co-factors responsible for methylation. To attain their goal, they will collect dissolved and particulate samples for trace metals and metalloenzymes from three stations along a biogeochemical gradient in the Tropical North Pacific (along 150 degrees West from 18 degrees North to the equator). Sinking particles from metal clean sediment traps will also be obtained. The samples will also be used to carry out shipboard incubation experiments using amendments of metals, metal-chelators, B12, and proteases to examine the sensitivity and metal limitation of heterotrophic, enzymatic degradation of organic matter within the oceanic "Twilight Zone" (100-500 m). This study will result in a novel metaproteomic/metalloenzyme datasets that should provide insights into the biogeochemical cycling of metals, as well as co-limitation of primary productivity and controls on the export of carbon from the photic zone. In addition to the final data being contributed to BCO-DMO, an online metaproteomic data server will be created so the community has access to the raw data files generated by this research.

[[table of contents](#) | [back to top](#)]

Program Information

U.S. GEOTRACES (U.S. GEOTRACES)

Website: <http://www.geotraces.org/>

Coverage: Global

GEOTRACES is a [SCOR](#) sponsored program; and funding for program infrastructure development is provided by the [U.S. National Science Foundation](#).

GEOTRACES gained momentum following a special symposium, S02: Biogeochemical cycling of trace elements and isotopes in the ocean and applications to constrain contemporary marine processes (GEOSECS II), at a 2003 Goldschmidt meeting convened in Japan. The GEOSECS II acronym referred to the Geochemical Ocean Section Studies To determine full water column distributions of selected trace elements and isotopes, including their concentration, chemical speciation, and physical form, along a sufficient number of sections in each ocean basin to establish the principal relationships between these distributions and with more traditional hydrographic parameters;

- * To evaluate the sources, sinks, and internal cycling of these species and thereby characterize more completely the physical, chemical and biological processes regulating their distributions, and the sensitivity of these processes to global change; and

- * To understand the processes that control the concentrations of geochemical species used for proxies of the past environment, both in the water column and in the substrates that reflect the water column.

GEOTRACES will be global in scope, consisting of ocean sections complemented by regional process studies. Sections and process studies will combine fieldwork, laboratory experiments and modelling. Beyond realizing the scientific objectives identified above, a natural outcome of this work will be to build a community of marine scientists who understand the processes regulating trace element cycles sufficiently well to exploit this knowledge reliably in future interdisciplinary studies.

Expand "Projects" below for information about and data resulting from individual US GEOTRACES research projects.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1260233
Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)	GBMF3782

[[table of contents](#) | [back to top](#)]