

Current velocity at station E at an oyster reef in Wassaw Sound, Georgia in November of 2014 (SensoryNCE project)

Website: <https://www.bco-dmo.org/dataset/709379>

Data Type: Other Field Results

Version: 2

Version Date: 2018-07-12

Project

» [The role of the sensory environment and predator chemical signal properties in determining NCE strength in cascading interactions on oyster reefs](#) (SensoryNCE)

Contributors	Affiliation	Role
Weissburg, Marc	Georgia Institute of Technology (GA Tech)	Principal Investigator
Kubanek, Julia	Georgia Institute of Technology (GA Tech)	Co-Principal Investigator
Webster, Donald	Georgia Institute of Technology (GA Tech)	Co-Principal Investigator, Contact
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: Lat:31.9199 Lon:-81.1392

Temporal Extent: 2014-11-03 - 2014-11-06

Dataset Description

This dataset contains velocity measurements at an intertidal and subtidal oyster reefs in Wassaw Sound, Georgia from November 3rd to 6th, 2014.

These data were published in Young et al., 2016 (see Fig. 3)

In addition to the "Get data" button above which provides various options to download the data in ASCII format, the originally submitted data is available in MATLAB .mat format ([FinalADV07_11032014.mat](#), 346 KB)

A subset of these data corresponding to Figure 3 in Young et al., 2016 can be downloaded as ([ADV_E_Velocity_11032014.mat](#), 203 KB)

Methods & Sampling

The equipment used is an Acoustic Doppler Velocimeter (ADV, Nortek Vector – Nortek AS, Rud, Norway). ADVs measure the fluid velocity at essentially a point in the flow. The ADV measurements have a resolution of 1 mm/s and were collected continuously at 1 Hz.

Sampling Location Map:




Image courtesy of David Young

Data Processing Description

Data collected during low tide (determined by instances of the PT at station D being dry) were discarded. Following the recommendations of Chanson et al. (2008) and Wilson et al. (2013), data for which the average correlation coefficient of the three beams was less than 70% for 45 consecutive samples, or for which the average over 300 samples of the mean correlation coefficient of the three beams dropped below 70%, were discarded. This filtering predominately removed data collected while the instrument was not submerged, and less than 2% of the submerged instrument data were removed due to filtering.

BCO-DMO Data Manager Processing Notes:

- * added a conventional header with dataset name, PI name, version date
- * data extracted from MATLAB .mat files to ASCII flat files to serve in the BCO-DMO system
- * modified parameter names to conform with BCO-DMO naming conventions
- * added approximate lat and lon of sampling location to data
- * data rounded to three decimal places
- * Time format changed from matlab datenum to Date and Time (yyyy-mm-dd,HH:MM:SS)
- * Data version 2 (2018-07-12) replaces data version 1 (2017-07-20) - improved metadata, added timestamp in UTC in addition to local date/time

[[table of contents](#) | [back to top](#)]

Data Files

File**ADV_E.csv**(Comma Separated Values (.csv), 1.64 MB)

MD5:c27840a5be1d94c043cf5447ced22d27

Primary data file for dataset ID 709379

[\[table of contents \]](#) | [\[back to top \]](#)

Related Publications

Young, D. L., Bruder, B. L., Haas, K. A., & Webster, D. R. (2016). The hydrodynamics of surface tidal flow exchange in saltmarshes. *Estuarine, Coastal and Shelf Science*, 172, 128-137. doi:[10.1016/j.ecss.2016.02.006](https://doi.org/10.1016/j.ecss.2016.02.006)

*Results**Methods*[\[table of contents \]](#) | [\[back to top \]](#)

Parameters

Parameter	Description	Units
date	Date (local EST) in format yyyy-mm-dd	unitless
time	Time (local EST) in format HH:MM	unitless
velocity_mag	Velocity magnitude	centimeters per second(cm/s)
velocity_dir	Velocity direction; angle relative to east	radians
lat_approx	Approximate latitude	decimal degrees
lon_approx	Approximate longitude	decimal degrees
ISO_DateTime_UTC	Timestamp (UTC) in standard ISO 8601:2004(E) format YYYY-mm-ddTHH:MM:SSZ	unitless

[\[table of contents \]](#) | [\[back to top \]](#)

Instruments

Dataset-specific Instrument Name	ADV, Nortek Vector
Generic Instrument Name	Acoustic Doppler Velocimeter
Dataset-specific Description	Acoustic Doppler Velocimeter (ADV, Nortek Vector - Nortek AS, Rud, Norway).
Generic Instrument Description	ADV is the acronym for acoustic doppler velocimeter. The ADV is a remote-sensing, three-dimensional velocity sensor. Its operation is based on the Doppler shift effect. The sensor can be deployed either as a moored instrument or attached to a still structure near the seabed. Reference: G. Voulgaris and J. H. Trowbridge, 1998. Evaluation of the Acoustic Doppler Velocimeter (ADV) for Turbulence Measurements. J. Atmos. Oceanic Technol., 15, 272-289. doi: http://dx.doi.org/10.1175/1520-0426(1998)0152.0.CO;2

[[table of contents](#) | [back to top](#)]

Deployments

SensoryNCE 2014

Website	https://www.bco-dmo.org/deployment/709363
Platform	Wassaw_Sound_GA
Start Date	2014-11-03
End Date	2014-11-06
Description	Wassaw Sound, Georgia

[[table of contents](#) | [back to top](#)]

Project Information

The role of the sensory environment and predator chemical signal properties in determining NCE strength in cascading interactions on oyster reefs (SensoryNCE)

Coverage: Intertidal and subtidal oyster reefs in Wassaw Sound, Georgia, US

Extracted from the NSF award abstract:

In this project, the investigators will examine the ability of top blue crab predators to indirectly benefit the abundance of basal oyster prey by reducing the density (consumptive effects, CEs) and suppressing foraging (non-consumptive effects, NCEs) of intermediate mud crab predators. These NCEs are mediated by chemical perception of aversive cues in blue crab urine and produce a behaviorally mediated trophic cascade. Through a series of manipulative experiments, the investigators will examine how the strength of this behaviorally-mediated trophic cascade is modulated and factors that influence perceptible range such as predator diet and intake rate, and the flow environment. The investigators will also determine the chemical identity, concentration and release rate of chemical cues.

Identifying the quantitative and molecular aspects of aversive cues, and linking them to behavioral responses that produce trophic cascades establishes the chemical basis of risk perception by prey and how this translates into cascading ecological effects. The use of perceptible range as a framework for evaluating the effects of both chemistry and environment provides an integrated view of processes affecting chemically-mediated NCEs. The use of a water borne predator-prey signaling system to test ideas on the strength of

NCEs should have broad applications.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1234449

[[table of contents](#) | [back to top](#)]