

Carbonate chemistry and light data from bottle samples

Website: <https://www.bco-dmo.org/dataset/719535>

Data Type: Other Field Results

Version:

Version Date: 2017-11-17

Project

» [Bermuda ocean Acidification and CO₂ reef iNvestigation](#) (BEACON)

Program

» [Ocean Carbon and Biogeochemistry](#) (OCB)

Contributors	Affiliation	Role
Andersson, Andreas	University of California-San Diego Scripps (UCSD-SIO)	Principal Investigator, Contact
Bates, Nicholas	Bermuda Institute of Ocean Sciences (BIOS)	Co-Principal Investigator
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:32.457333 E:-64.797469 S:32.400777 W:-64.834759

Dataset Description

Surface seawater samples were collected approximately monthly and analyzed for carbonate chemistry and nutrients.

These data were utilized in the following publication:

Courtney, T. A., Lebrato, M., Bates, N. R., Collins, A., de Putron, S. J., Garley, R., ... & Andersson, A. J. (2017). Environmental controls on modern scleractinian coral and reef-scale calcification. *Science advances*, 3(11), e1701356. doi: [10.1126/sciadv.1701356](https://doi.org/10.1126/sciadv.1701356)

Methods & Sampling

At each sample site, a 5 liter Niskin bottle was deployed at 0.5-1.0 m depth according to best practices. Samples for DIC and TA were collected in 200 mL Kimax glass sample bottles, and immediately fixed using 100 µL saturated solution of HgCl₂. TA samples were analyzed via closed cell potentiometric titrations using a VINDTA 3S system and DIC was analyzed using coulometric methods on a VINDTA 3c or an infrared based analysis on an AIRICA system. Accuracy and precision of TA and DIC measurements were evaluated by

certified reference materials provided by the laboratory of Prof. A. Dickson at Scripps Institution of Oceanography yielding typical precision and accuracy of $\pm 1-2 \mu\text{mol/kg}$. A YSI Handheld Multiparameter Instrument was used to measure in situ temperature ($\pm 0.15^\circ\text{C}$), salinity ($\pm 1\%$), dissolved oxygen ($\pm 2\%$), and percent saturation dissolved oxygen ($\pm 2\%$). Nutrient samples were filtered upon sampling using a $0.4 \mu\text{M}$ filter and frozen in opaque plastic bottles until processing at Woods Hole Oceanographic Institution Nutrient Analytical Facility using agency methods of analysis for ammonium (detection limit $0.034 \mu\text{mol/L}$, method G-172-96), nitrite+nitrate (detection limit $0.010 \mu\text{mol/L}$, method G-177-96), silicate (detection limit $0.016 \mu\text{mol/L}$, method G-177-96), and phosphate (detection limit $0.025 \mu\text{mol/L}$, method G-297-03).

Sampling locations: Bermuda Platform: Hog Reef (32.45733, -64.83476) and Crescent Reef (32.40078, -64.79747).

Data Processing Description

TAcorr and DICcorr represents instrument TA value and instrument DIC value calibrated as fixed offsets to certified reference material from the laboratory of A. Dickson at Scripps Institution of Oceanography.

BCO-DMO Data Manager Processing Notes:

- * added a conventional header with dataset name, PI name, version date
- * modified parameter names to conform with BCO-DMO naming conventions
- * NaN values changed to nd for "no data"
- * Added Lat,Lon for Hog and Cres reefs into the data

[[table of contents](#) | [back to top](#)]

Data Files

File
BottleSamples.csv (Comma Separated Values (.csv), 11.24 KB) MD5:38c166f3ea7fefdea7d2bb702867a350 Primary data file for dataset ID 719535

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Date	Day of sampling in formay DD-MM-YY	unitless
Time	Time of sampling in format HH:mm	unitless
Site	Site name of coral in situ growth (Hog = Hog Reef, Bermuda or Cres = Crescent Reef, Bermuda)	unitless
Lat	Latitude of site	decimal degrees
Lon	Longitude of site	decimal degrees
Temp	Sea surface temperature	degrees Celsius
Sal_YSI	Sea surface salinity measured by YSI instrument	parts per thousand (ppt)
Sal_Salinometer	Sea surface salinity by Autosal Salinometer	Practical Salinity Units (PSU)
DO_mg_l	Sea surface dissolved oxygen	milligrams per liter (mg/L)
DO_percent	Sea surface dissolved oxygen	percent saturation (%)
uM_NH4	Sea surface ammonium	micromolar ($\mu\text{mol/L}$)
uM_SiO4	Sea surface silicate	micromolar ($\mu\text{mol/L}$)
uM_PO4	Sea surface phosphate	micromolar ($\mu\text{mol/L}$)
uM_NO2_NO3	Sea surface nitrite + nitrate	micromolar ($\mu\text{mol/L}$)
TACorr	Sea surface total alkalinity (corrected value, see processing description)	micromoles per liter per kilogram ($\mu\text{mol/kg}$)
DICcorr	Sea surface dissolved inorganic carbon (corrected value, see processing description)	micromoles per liter per kilogram ($\mu\text{mol/kg}$)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Autosal Salinometer 8400B (Salinity)
Generic Instrument Name	Autosal salinometer
Dataset-specific Description	Samples for salinity were collected in glass bottles and later analyzed using an autosalinometer (Guildline Instruments)
Generic Instrument Description	The salinometer is an instrument for measuring the salinity of a water sample.

Dataset-specific Instrument Name	AIRICA (Marianda Inc.)
Generic Instrument Name	Inorganic Carbon Analyzer
Dataset-specific Description	The Automated Infra Red Inorganic Carbon Analyzer (AIRICA) utilizes infrared detection of CO ₂ gas purged from an acidified seawater sample. A high-precision syringe pump extracts the seawater sample, acidifies the sample with phosphoric acid, and analyzes the gas released with an infrared light analyzer (LICOR). The CO ₂ signal is integrated for each sample to quantify the total inorganic carbon for a given aliquot of seawater analyzed. Three aliquots and peak integrations are performed for each seawater sample and averaged to determine the dissolved inorganic carbon for each sample. Precision was typically $\pm 1\text{--}2\text{ }\mu\text{mol/kg}$ for TA. Please see http://marianda.com/index.php?site=products&subsite=airica for a complete instrument description.
Generic Instrument Description	Instruments measuring carbonate in sediments and inorganic carbon (including DIC) in the water column.

Dataset-specific Instrument Name	VINDTA 3c (Marianda Inc.)
Generic Instrument Name	MARIANDA VINDTA 3C total inorganic carbon and titration alkalinity analyser
Dataset-specific Description	DIC was analyzed using coulometric methods on a VINDTA 3c or an infrared-based analysis on an AIRICA system. The Versatile INstrument for the Determination of Total inorganic carbon and titration Alkalinity (VINDTA) 3C is a laboratory alkalinity titration system combined with an extraction unit for coulometric determination of total dissolved inorganic carbon. The instrument analyzes alkalinity and dissolved inorganic carbon content of a sample in parallel. The sample transport is performed with peristaltic pumps and acid is added to the sample using a membrane pump. No pressurizing system is required and only one gas supply (nitrogen or dry and CO ₂ -free air) is necessary. The system uses a Metrohm Titrino 719S, an ORION-Ross pH electrode and a Metrohm reference electrode. The burette, the pipette and the analysis cell have a water jacket around them. Precision is typically $\pm 1\text{ }\mu\text{mol/kg}$ for TA. DIC is analyzed with a UIC coulometer. Please see http://marianda.com/index.php?site=products&subsite=vindta3c for a complete instrument description.
Generic Instrument Description	The Versatile INstrument for the Determination of Total inorganic carbon and titration Alkalinity (VINDTA) 3C is a laboratory alkalinity titration system combined with an extraction unit for coulometric titration, which simultaneously determines the alkalinity and dissolved inorganic carbon content of a sample. The sample transport is performed with peristaltic pumps and acid is added to the sample using a membrane pump. No pressurizing system is required and only one gas supply (nitrogen or dry and CO ₂ -free air) is necessary. The system uses a Metrohm Titrino 719S, an ORION-Ross pH electrode and a Metrohm reference electrode. The burette, the pipette and the analysis cell have a water jacket around them. Precision is typically $\pm 1\text{ }\mu\text{mol/kg}$ for TA and/or DIC in open ocean water.

Dataset-specific Instrument Name	Niskin Bottle
Generic Instrument Name	Niskin bottle
Dataset-specific Description	From 2007 to 2012, surface seawater samples from the Bermuda coral reef platform were collected once a month at 0.5–1-m depth using a 5-L Niskin bottle.
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset-specific Instrument Name	MARIANDA VINDTA 3S
Generic Instrument Name	Titration
Dataset-specific Description	Total alkalinity (TA) samples were analyzed via closed cell potentiometric titrations using a VINDTA 3S system. VINDTA (Versatile INstrument for the Determination of Titration Alkalinity). The system uses a Metrohm Titrino 719S, an ORION-Ross pH electrode and a Metrohm reference electrode. The burette, the pipette and the analysis cell have a water jacket around them. Precision is typically +/- 1 umol/kg for TA. Please see for a complete instrument description.
Generic Instrument Description	Titration is an instrument that incrementally add quantified aliquots of a reagent to a sample until the end-point of a chemical reaction is reached.

Dataset-specific Instrument Name	YSI 556 Handheld Multiparameter Instrument
Generic Instrument Name	Water Quality Multiprobe
Dataset-specific Description	A YSI 556 Handheld Multiparameter Instrument was used to measure in situ temperature (accuracy $\pm 0.15^{\circ}\text{C}$), salinity (accuracy $\pm 1\%$), DO _{mg} (accuracy $\pm 2\%$), and DO _% (accuracy $\pm 2\%$).
Generic Instrument Description	An instrument which measures multiple water quality parameters based on the sensor configuration.

[[table of contents](#) | [back to top](#)]

Deployments

BIOS_BEACON

Website	https://www.bco-dmo.org/deployment/626096
Platform	BIOS_Small_Boat
Start Date	2007-06-15
End Date	2012-09-18
Description	Sample collection platforms for the BEACON Project. The samples were collected from a small boat (27 ft Twin Vee or 26 ft Seadance)

[[table of contents](#) | [back to top](#)]

Project Information

Bermuda ocean Acidification and CO₂ reef iNvestigation (BEACON)

Website: <https://scripps.ucsd.edu/research/projects/bermuda-ocean-acidification-and-coral-reef-investigation-beacon>

Coverage: Bermuda

NSF abstract:

Ocean acidification owing to anthropogenic emission of CO₂ is a significant and imminent threat to marine calcifying organisms and ecosystems such as corals and coral reefs. As a result of future ocean acidification, i.e., increasing seawater CO₂, and decreasing pH, carbonate ion concentration [CO₃], and carbonate saturation state, it is likely that marine calcifiers will have difficulty growing their shells and skeletons of calcium carbonate (CaCO₃) at their present rates. Dissolution of carbonate sediments and structures are also likely to increase, and could ultimately exceed calcification and CaCO₃ production, leading to a transition from net accumulation to a net loss in carbonate material of individual coral colonies, coral communities and coral reef ecosystems. Because of Bermuda's relatively high-latitude location (32° N), the annual average surface seawater [CO₃] is lower in Bermuda than regions closer to the tropics. As a consequence, the Bermuda coral reef is likely to experience critical [CO₃] values and net dissolution before its tropical counterparts as a result of continued ocean acidification. Furthermore, a natural gradient in [CO₃] exists along the Bermuda reef with environmental parameters such as, light, temperature, and nutrients being near identical. This gradient allows for unique cross-comparisons of calcification of individual calcifiers and calcifying communities under different [CO₃] in a natural environment.

In this study, researchers at the Bermuda Institute of Ocean Science (BIOS) will launch the BEACON project to further our understanding of the consequences of ocean acidification to the process of calcification and CaCO₃ production at three different spatial scales including (1) individual coral colonies, (2) local reef communities, and (3) regional coral reef ecosystems. They will conduct (1) in situ and in vitro experiments to assess growth and evaluate net calcification of individual coral colonies of three different species common to Bermuda and the Caribbean exposed to different [CO₃] under both natural and controlled experimental conditions; (2) diel and quasi-lagrangian calcification experiments to evaluate net calcification of local reef communities and in moving water masses along the natural [CO₃] gradient existent on the Bermuda platform; and (3) time series data collected across the Bermuda platform and offshore, to evaluate net calcification and CaCO₃ production of the Bermuda coral reef ecosystem and platform over seasonal and annual cycles.

Broader impacts: This project will provide fundamental data on the consequences of ocean acidification to coral reefs on different temporal and spatial scales. Combined, knowledge at each of the scales will contribute to an improved understanding of this problem in a broader context, i.e., the effect on coral reefs as a global entity and role in the global carbon cycle during past, present and future seawater chemical conditions. As the meaning implies, the research team envisions BEACON to serve as a guiding light to assist researchers and policymakers in framing future strategies and making decisions regarding the management of coral reefs and CO₂ emission policies in order to establish CO₂ stabilization targets. Scientific understanding and research products from the project will be specifically used in the BIOS explorer program, CoE POGO ocean acidification module, and the coral reef ecology class taught at BIOS. It will also contribute to developing the research and technical skills of one graduate student and one research technician at BIOS, and will include the involvement

of interns and NSF REU fellowship students each year of the project.

Based on the findings of the BEACON project, and especially the results published in Andersson et al. (Nature Climate Change, 4, 56-61, 2014) and Yeakel et al. (PNAS, 112, 14512-14517, 2015), BEACON II (<https://www.bco-dmo.org/project/737955>) aims to assess the links between offshore and reef biogeochemistry by continuing and expanding on the physical and chemical measurements on the Bermuda coral reef and in the surrounding Sargasso Sea.

[[table of contents](#) | [back to top](#)]

Program Information

Ocean Carbon and Biogeochemistry (OCB)

Website: <http://us-ocb.org/>

Coverage: Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO₂ and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0928406

[[table of contents](#) | [back to top](#)]