

Temperature and salinity from 4 field sites in Bogue Sound, North Carolina from 2014 to 2015.

Website: <https://www.bco-dmo.org/dataset/720188>

Data Type: Other Field Results

Version: 1

Version Date: 2017-12-14

Project

» [Microbial Regulation of Greenhouse Gas N2O Emission from Intertidal Oyster Reefs](#) (Oyster Reef N2O Emission)

Contributors	Affiliation	Role
Piehler, Michael F.	University of North Carolina at Chapel Hill (UNC-Chapel Hill-IMS)	Principal Investigator
Ake, Hannah	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Temperature and salinity from 4 field sites in Bogue Sound, North Carolina from 2014 to 2015.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: Lat:34.710129 Lon:-76.867882

Temporal Extent: 2014 - 2015

Dataset Description

Temperature and salinity from 4 field sites in Bogue Sound, located in coastal North Carolina.

Methods & Sampling

Water samples collected on-site. The temperature was measured with a mercury thermometer and salinity measured with a refractometer. The 4 field sites are sites of oyster reef and salt marsh restoration. Identified in data as UNC Institute of Marine Sciences (IMS), Carrot Island (Carrot), NOAA Beaufort (NOAA), and Army Marsh (Army). All sites located in Bogue Sound near Morehead City, NC.

Data Processing Description

BCO-DMO Data Processing Description:

- data sorted by season, site

Data Files

File
salinity_temp.csv (Comma Separated Values (.csv), 313 bytes) MD5:9f5811e80ee76aa8ba596ed4906062a4
Primary data file for dataset ID 720188

Parameters

Parameter	Description	Units
Season	Season data were collected	unitless
Site	Site where data were collected	unitless
Salinity	Salinity	PSU
Temperature	Water temperature	Celsius

Instruments

Dataset-specific Instrument Name	Mercury thermometer
Generic Instrument Name	digital thermometer
Dataset-specific Description	Used to measure water temperature
Generic Instrument Description	An instrument that measures temperature digitally.

Dataset-specific Instrument Name	Refractometer
Generic Instrument Name	Refractometer
Dataset-specific Description	Used to measure salinity
Generic Instrument Description	A refractometer is a laboratory or field device for the measurement of an index of refraction (refractometry). The index of refraction is calculated from Snell's law and can be calculated from the composition of the material using the Gladstone-Dale relation. In optics the refractive index (or index of refraction) n of a substance (optical medium) is a dimensionless number that describes how light, or any other radiation, propagates through that medium.

Project Information

Microbial Regulation of Greenhouse Gas N₂O Emission from Intertidal Oyster Reefs (Oyster Reef N₂O Emission)

Extracted from the NSF award abstract:

Oyster reefs are biogeochemical hot spots and prominent estuarine habitats that provide disproportionate ecological function. Suspension-feeding eastern oysters, *Crassostrea virginica*, are capable of improving water quality and diminishing eutrophication by filtering nutrients and particles from the water and depositing them in the sediments. Remineralization of these deposits may enhance sedimentary denitrification that facilitates nitrogen removal in tidal estuaries. However, the scientific underpinning of oyster reef function has been challenged in various studies. In addition, recent studies of filter feeding invertebrates reported the production of nitrous oxide (N₂O), a greenhouse gas, as an end product of incomplete denitrification by gut microbes. *C. virginica* could be another source of N₂O flux from intertidal habitats. Preliminary work indicated substantial N₂O production from individual oysters. The estimated N₂O production from high density oyster reefs may exceed the N₂O flux measured from some estuaries. With the new discovery of N₂O emission and uncertainty regarding eutrophication control, the ecological value of oyster reef restoration may become equivocal.

This project will quantify N₂O fluxes to understand the factors controlling N₂O emission from oyster reefs. Sedimentary N processes will be examined to develop an oyster reef N model to estimate N₂O emission from tidal creek estuaries relative to other N cycling processes. The PIs hypothesize that intertidal oyster reefs are a substantial source of N₂O emission from estuarine ecosystems and the magnitude of emission may be linked to water quality. If substantial N₂O flux from oyster reefs is validated, ecological benefits of oyster reef restoration should be reevaluated. This interdisciplinary research team includes a microbial ecologist, a biogeochemist, an ecologist and an ecosystem modeler. They will utilize stable isotope and molecular microbiological techniques to quantify oyster N₂O production, elucidate microbial sources of N₂O emission from oysters and sediments, and estimate seasonal variation of N₂O fluxes from oyster reefs. Measurements from this study will be integrated into a coupled oyster bioenergetics-sediment biogeochemistry model to compare system level rates of N cycling on oyster reefs as a function of oyster density and water quality. Modeling results will be used to assess the relative trade-offs of oyster restoration associated with N cycling. They expect to deliver the following end products: 1) estimation of annual N₂O flux from oyster reefs as an additional source of greenhouse gases from estuaries, 2) a better understanding of the environmental and microbial factors influencing N₂O and N₂ fluxes in tidal estuaries, 3) transformative knowledge for the effect of oyster restoration on water quality enhancement and ecosystem function, 4) direct guidance for oyster restoration projects whose goals include water quality enhancement, and 5) a modeling tool for use in research and restoration planning.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1233327