

# The effects of vermetids and sediments on coral growth in Moorea, French Polynesia from June to August 2012 (Vermetids\_Corals project)

**Website:** <https://www.bco-dmo.org/dataset/726215>

**Data Type:** experimental

**Version:** 2017-10-05

## Project

» [Spatial patterns of coral-vermetid interactions: short-term effects and long-term consequences](#)

(Vermetids\_Corals)

Contributors	Affiliation	Role
<a href="#">Zill, Julie</a>	University of Hawai'i (UH)	Principal Investigator
<a href="#">Gil, Michael A.</a>	University of California-Davis (UC Davis)	Co-Principal Investigator
<a href="#">Osenberg, Craig</a>	University of Georgia (UGA)	Co-Principal Investigator, Contact
<a href="#">Biddle, Mathew</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:-17.47279 E:-149.78277 S:-17.48365 W:-149.84698

**Temporal Extent:** 2012-06 - 2012-08

## Dataset Description

A manipulative experiment to determine the growth rate of corals in the presence and absence of sediment and vermetids.

### Related Datasets:

- Zill et al 2017 Coral Growth: <https://www.bco-dmo.org/dataset/726215> (Current page)
- Zill et al 2017 Sediment Removal Rate: <https://www.bco-dmo.org/dataset/726232>

## Methods & Sampling

In June-August 2012, we conducted a 54-day field experiment on the north shore of Moorea, French Polynesia. We manipulated the presence of the vermetid *Ceraesignum maximum* (formerly *Dendropoma maximum*) and high sedimentation, yielding four fully-crossed treatments. In response to which, we measured the growth of the reef-building coral massive *Porites* (a species complex mostly composed of *P. lobata* and *P. lutea*, see

Veron 1995).

This study was performed in a shallow lagoonal backreef area, where vermetid densities can be quite high (up to >100 per m<sup>2</sup>: Smalley 1984; Shima et al. 2010). We selected 36 large (approximately 1 meter in diameter), mostly-dead colonies of massive *Porites* to serve as deployment sites for our experimental corals. These colonies occurred at a similar depth along a 150-m transect that was roughly parallel to the reef crest and perpendicular to prevailing water currents. Because differences in water flow could affect vermetid net casting efficacy and sediment removal rates, we quantified relative flow variation among sites using calibrated clod cards (Jokiel & Morrissey 1993). The flat top of each site was home to an average of  $5 \pm 1$  (SD) vermetids in a 400 square centimeter area, ensuring that mucus nets would consistently cover the experimental corals receiving the “+vermetid” treatments. We randomly assigned the four experimental treatments to our sites, from half of which we removed all vermetids to create the “-vermetid” treatments. Our experimental corals were 36 juvenile massive *Porites* colonies that we collected from a nearby location. Each coral was attached to a small plastic base using marine epoxy, and buoyantly weighed (following Davies 1989). They were then assigned to nine blocks of four corals based on similarity in morphology, and outplanted to each deployment site using u-nails.

After 54 days, corals were retrieved, sediments were removed, the corals’ final buoyant mass (Davies 1989) and surface area (using the aluminum foil method of Marsh 1970) were quantified to calculate the per-tissue-area change in coral mass.

## Data Processing Description

### BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Zilletal_2017_Coralgrowth.csv</b> (Comma Separated Values (.csv), 692 bytes) MD5:f3fbefa7f3a45db1bd018b8cc45bec5c
Primary data file for dataset ID 726215

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Davies, P.S. (1989). Short-term growth measurements of corals using an accurate buoyant weighing technique. *Marine Biology*, 101(3), 389–395. doi:10.1007/bf00428135 <https://doi.org/10.1007/BF00428135>  
*Methods*

Jokiel, P., & Morrissey, J. (1993). Water motion on coral reefs: evaluation of the “clod card” technique . *Marine Ecology Progress Series*, 93, 175–181. doi:[10.3354/meps093175](https://doi.org/10.3354/meps093175)  
*Methods*

Shima, J. S., Osenberg, C. W., & Stier, A. C. (2010). The vermetid gastropod *Dendropoma maximum* reduces coral growth and survival. *Biology Letters*, 6(6), 815–818. doi:[10.1098/rsbl.2010.0291](https://doi.org/10.1098/rsbl.2010.0291)  
*Methods*

Smally, T. L. (1984). Possible effects of intraspecific competition on the population structure of a solitary vermetid mollusc. *Marine ecology progress series*. Oldendorf, 14(2), 139-144.  
*Methods*

Zill, J. A., Gil, M. A., & Osenberg, C. W. (2017). When environmental factors become stressors: interactive effects of vermetid gastropods and sedimentation on corals. *Biology Letters*, 13(3), 20160957. doi:[10.1098/rsbl.2016.0957](https://doi.org/10.1098/rsbl.2016.0957)

## Parameters

Parameter	Description	Units
Unit	the name of individual coral; the letter corresponds to the treatment received (P=Control; Y=Vermetids only; O=sediment only; R= both vermetids and sediment)	unitless
Block	the block number that the experimental coral belonged to which was based on morphology	unitless
Vermetids	presence [1] or absence [0] of vermetids in the vicinity of the coral	unitless
Sediment	whether the high sediment treatment was applied [1] or not applied [0]	unitless
GrowthRate	how much the coral grew on average per day over the duration of the experiment. Mass was calculated using the buoyant mass method (Davies 1989), and then converted to growth rate by dividing mass by surface for each coral and length of experiment (54 days).	milligrams per centimeter squared per day (mg/cm <sup>2</sup> day)

## Deployments

### Osenberg et al Moorea

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/644752">https://www.bco-dmo.org/deployment/644752</a>
<b>Platform</b>	Osenberg et al Moorea
<b>Start Date</b>	2003-05-19
<b>End Date</b>	2015-07-12

## Project Information

### Spatial patterns of coral-vermetid interactions: short-term effects and long-term consequences (Vermetids\_Corals)

**Coverage:** Moorea, French Polynesia (-17.48 degrees S, -149.82 degrees W)

#### Description from NSF abstract:

Ecological surprises are most likely to be manifest in diverse communities where many interactions remain uninvestigated. Coral reefs harbor much of the world's biodiversity, and recent studies by the investigators suggest that one overlooked, but potentially important, biological interaction involves vermetid gastropods. Vermetid gastropods are nonmobile, tube-building snails that feed via an extensive mucus net. Vermetids

reduce coral growth by up to 80%, and coral survival by as much as 60%. Because effects vary among coral taxa, vermetids may substantially alter the structure of coral communities as well as the community of fishes and invertebrates that inhabit the coral reef.

The investigators will conduct a suite of experimental and observational studies that: 1) quantify the effects of four species of vermetids across coral species to assess if species effects and responses are concordant or idiosyncratic; 2) use meta-analysis to compare effects of vermetids relative to other coral stressors and determine the factors that influence variation in coral responses; 3) determine the role of coral commensals that inhabit the branching coral, *Pocillopora*, and evaluate how the development of the commensal assemblage modifies the deleterious effects of vermetids; 4) determine how vermetid mucus nets affect the local environment of corals and evaluate several hypotheses about proposed mechanisms; and 5) assess the long-term implications of vermetids on coral communities and the fishes and invertebrates that depend on the coral.

**Note:** The Principal Investigator, Dr. Craig W. Osenberg, was at the University of Florida at the time the NSF award was granted. Dr. Osenberg moved to the University of Georgia during the summer of 2014 ([current contact information](#)).

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1130359</a>

[ [table of contents](#) | [back to top](#) ]