

Water column nitrate+nitrite d15N measurements from R/V L'Atalante in the southwest Pacific Ocean between New Caledonia and Tahiti from February to March 2015

Website: <https://www.bco-dmo.org/dataset/733237>

Data Type: Cruise Results

Version: 2018-04-11

Project

» [Quantifying nitrogen fixation along unique geochemical gradients in the southwest Pacific Ocean](#) (SW Pac N2 fixation)

Contributors	Affiliation	Role
Knapp, Angela N.	Florida State University (FSU)	Principal Investigator, Contact
Bonnet, Sophie	Mediterranean Institute of Oceanography (MIO)	International Collaborator
Moutin, Thierry	Mediterranean Institute of Oceanography (MIO)	International Collaborator
Grosso, Oliver	Mediterranean Institute of Oceanography (MIO)	Scientist
Leblond, Nathalie	Villefranche Oceanographic Laboratory (LOV)	Scientist
McCabe, Kelly M.	Florida State University (FSU)	Student
Biddle, Mathew	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Constraining the rates and spatial distribution of di-nitrogen (N₂) fixation fluxes to the ocean informs our understanding of the environmental sensitivities of N₂ fixation as well as the timescale over which the fluxes of nitrogen (N) to and from the ocean may respond to each other. Here we quantify rates of N₂ fixation as well as its contribution to export production along a zonal transect in the Western Tropical South Pacific (WTSP) Ocean using N isotope ("d15N") budgets. Comparing measurements of water column nitrate+nitrite d15N with the d15N of sinking particulate N at a western, central, and eastern station, these d15N budgets indicate high, modest, and low rates of N₂ fixation at the respective stations. The results also imply that N₂ fixation supports exceptionally high, i.e., >50%, of export production at the western and central stations, which are also proximal to the largest iron sources. These geochemically-based rates of N₂ fixation are equal to or greater than those previously reported in the tropical North Atlantic, indicating that the WTSP Ocean has the capacity to support globally significant rates of N₂ fixation, which may compensate for N removal in the oxygen deficient zones of the eastern tropical Pacific.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:-18.179 E:-172.7813 S:-19.9832 W:168.0118

Temporal Extent: 2015-02-23 - 2015-03-30

Dataset Description

This data set includes water column nitrate+nitrite d15N measurements. These measurements were used together with measurements of the d15N of particulate nitrogen collected in floating sediment traps that were deployed for several days to calculate the relative contribution of subsurface nitrate and nitrogen from N2 fixation for supporting export production (“d15N budgets”). The results suggest that N2 fixation supported a majority of export at Long Duration (LD) stations A and B, and a minor fraction of export at LD C. The results at LD stations A and B are unique compared to other d15N budgets from the oligotrophic regions, whereas the results from LD C are similar to prior reports from the eastern tropical South Pacific as well as the North Pacific near Hawaii. Additionally, these data are compared with other metrics of N2 fixation made on the same cruise.

Methods & Sampling

NO3-+NO2- d15N analysis was by the “denitrifier method” and followed the methods described by Sigman et al., 2001, Casciotti et al., 2002, McIlvin and Casciotti, 2011, and Weigand et al., 2016. Briefly, NO3-+NO2- was quantitatively reduced to N2O by *Pseudomonas aureofaciens* and *Pseudomonas chlororaphis*, which was then cryogenically focused and analyzed on an isotope ratio mass spectrometer. A volume of sample was added to each bacterial vial to achieve a final quantity of 10 or 20 nmols N2O, which was then purged from the vial using a helium carrier gas. The d15N of N2O in samples was calibrated with the international isotopic reference materials described in the Processing Description.

Data Processing Description

Nitrate+nitrite concentration was measured by others using colorimetric methods, and can be found at: <http://www.obs-vlfr.fr/proof/php/outpace/outpace.php>

The average precision of nitrate+nitrite d15N measurements was <0.2 per mil, but with the standard deviation for duplicate analyses of each sample reported here. Samples were calibrated with IAEA N3 and USGS 34 NO3-d15N isotopic reference materials as described in McIlvin and Casciotti, 2011.

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date.
- modified parameter names to conform with BCO-DMO naming conventions.
- re-formatted date from dd Month yyyy (day month year) to yyymmdd.
- appended date, latitude, and longitude to each data entry.
- If an observation did not exist, 'nd' (no data) was inserted.

[[table of contents](#) | [back to top](#)]

Data Files

File
OUTPACE.csv (Comma Separated Values (.csv), 5.24 KB) MD5:2b45eb3c7b6fdb0b23e874a5726846d7 Primary data file for dataset ID 733237

[[table of contents](#) | [back to top](#)]

Related Publications

Casciotti, K. L., Sigman, D. M., Hastings, M. G., Böhlke, J. K., & Hilkert, A. (2002). Measurement of the Oxygen Isotopic Composition of Nitrate in Seawater and Freshwater Using the Denitrifier Method. *Analytical Chemistry*, 74(19), 4905–4912. doi:[10.1021/ac020113w](https://doi.org/10.1021/ac020113w)

General

McIlvin, M. R., & Casciotti, K. L. (2011). Technical Updates to the Bacterial Method for Nitrate Isotopic Analyses. Analytical Chemistry, 83(5), 1850–1856. doi:[10.1021/ac1028984](https://doi.org/10.1021/ac1028984)

General

Sigman, D. M., Casciotti, K. L., Andreani, M., Barford, C., Galanter, M., & Böhlke, J. K. (2001). A Bacterial Method for the Nitrogen Isotopic Analysis of Nitrate in Seawater and Freshwater. Analytical Chemistry, 73(17), 4145–4153. doi:[10.1021/ac010088e](https://doi.org/10.1021/ac010088e)

General

Weigand, M. A., Foriel, J., Barnett, B., Oleynik, S., & Sigman, D. M. (2016). Updates to instrumentation and protocols for isotopic analysis of nitrate by the denitrifier method. Rapid Communications in Mass Spectrometry, 30(12), 1365–1383. doi:[10.1002/rcm.7570](https://doi.org/10.1002/rcm.7570)

Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Cast	cast number	unitless
Depth	depth of observation	meters
Sigma_theta	potential density	kilograms per meter cubed (kg/m3)
NO3_NO2	nitrate+nitrite concentration ([NO3-+NO2-])	micromole (uM)
NO3_NO2_d15N	nitrate+nitrite d15N (NO3-+NO2- d15N)	per mil
NO3_NO2_d15N_1_SD	nitrate+nitrite d15N standard deviation	per mil
Date	Date of observation in four digit year; 2 digit month; 2 digit day (yyyymmdd) format	unitless
Longitude	Longitude with west negative.	decimal degrees
Latitude	Latitude with south negative.	decimal degrees
Station	name of the station	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Thermo Finnigan Delta V isotope ratio mass spectrometer.
Generic Instrument Name	Isotope-ratio Mass Spectrometer
Dataset-specific Description	Nitrate+nitrite d15N was measured using a Thermo Finnigan Delta V isotope ratio mass spectrometer.
Generic Instrument Description	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

[[table of contents](#) | [back to top](#)]

Deployments

OUTPACE

Website	https://www.bco-dmo.org/deployment/663891
Platform	R/V L'Atalante
Start Date	2015-02-18
End Date	2015-04-03
Description	Oligotrophy to UTRa-oligotrophy PACific Experiment (OUTPACE) cruise DOI: http://dx.doi.org/10.17600/15000900 For more information see the cruise website: https://outpace.mio.univ-amu.fr/?lang=en South west Pacific between New Caledonia (166°28' E; 22°14' S) and Tahiti (149°36' W; 17°34' S) 0-2000 m

[[table of contents](#) | [back to top](#)]

Project Information

Quantifying nitrogen fixation along unique geochemical gradients in the southwest Pacific Ocean (SW Pac N2 fixation)

Website: <http://scope.soest.hawaii.edu/data/lava/lava.html>

Coverage: Southwest Pacific Ocean between New Caledonia and Tahiti along ~18 deg S

NSF Award Abstract:

The availability of nitrogen in the surface ocean plays a critical role regulating rates of primary productivity in the ocean, and thus through modification of the carbon cycle, nitrogen has the capacity to influence climate. The dominant source of biologically available nitrogen to the ocean is through a process known as di-nitrogen (N₂) fixation, which involves the reduction of N₂ gas dissolved in seawater to ammonium by microbes referred to as diazotrophs. While significant progress has been made identifying a diversity of marine diazotrophs in recent years using molecular tools, quantifying global rates of N₂ fixation, and identifying which ocean basin supports the highest fluxes, has remained a vexing question. This research will quantify rates of N₂ fixation as well as its importance for supporting production in the southwest Pacific Ocean. Results from this research will shed light on the sensitivities of N₂ fixation (temperature, iron concentrations) as well as the extent of spatial and temporal coupling of nitrogen sources and sinks in the ocean. The work will be carried out by an early career scientist, and involve mentoring of young women, middle school girls and minorities, training of undergraduate and graduate researchers, and international collaborations.

Identifying the spatial distribution of the largest di-nitrogen (N₂) fixation fluxes to the ocean remains a critical goal of chemical oceanography. The spatial distribution can inform our understanding of the environmental sensitivities of N₂ fixation and the capacity for the dominant marine nitrogen (N) source and sink processes to respond to each other and thus influence the global carbon cycle and climate. In addition to temperature, two factors are at the heart of the current debate over what influences the spatial distribution of N₂ fixation in the ocean: 1) the presence of adequate iron to meet the needs of N₂ fixing microbes, and, 2) the absolute concentrations as well as ratios of surface ocean nitrate and phosphate concentrations that are low relative to the "Redfield" ratio, which are thought to favor N₂ fixing microbes. This project will test the effects of gradients in atmospheric dust deposition on N₂ fixation rates when surface waters have relatively constant but favorable nitrate to phosphate concentrations. The work will be carried out in the southwest Pacific, a region highlighted by new modeling work for its unique geochemical characteristics that are expected to favor significant N₂ fixation fluxes. Nitrate+nitrite d¹⁵N as well as total dissolved nitrogen (TDN) concentration and d¹⁵N will be measured in water column samples collected on a French cruise and sediment traps were deployed to capture the sinking particulate N flux. The results will be compared with published work to evaluate which ocean regions support the largest N₂ fixation fluxes.

More information:

This project was part of the Oligotrophy to UTRa-oligotrophy PACific Experiment (OUTPACE) cruise in the Southwest Pacific between New Caledonia (166°28' E; 22°14' S) and Tahiti (149°36' W; 17°34' S) 0-2000 m

* OUTPACE cruise (doi: <http://dx.doi.org/10.17600/15000900>)

* OUTPACE website: <https://outpace.mio.univ-amu.fr/?lang=en>

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1537314

[[table of contents](#) | [back to top](#)]