

Counts of picocyanobacteria from SMILE cruise (SKQ201703S) at the North Pacific Subtropical Front in March 2017

Website: <https://www.bco-dmo.org/dataset/736702>

Data Type: Cruise Results

Version: 1

Version Date: 2018-05-16

Project

» [Microbial ecology of coexisting ecotypes: Are all Prochlorococcus equal?](#) (ProEco)

Program

» [Ocean Time-series Sites](#) (Ocean Time-series)

Contributors	Affiliation	Role
Thompson, Anne W.	Portland State University (PSU)	Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

This dataset includes counts of picocyanobacteria from SMILE cruise (SKQ201703S) at the North Pacific Subtropical Front in March 2017.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:35.18034 E:-139.241 S:25.98495 W:-147.0968

Temporal Extent: 2017-03-06 - 2017-03-27

Dataset Description

Counts of prochlorococcus, synechococcus, and picoeukaryotes from SMILE cruise (SKQ201703S) at the North Pacific Subtropical Front in March 2017.

Methods & Sampling

Water was sampled from the CTD into 15mL sterile polypropylene tubes. 2mL samples were fixed with 0.125% glutaraldehyde then flash frozen in liquid nitrogen and stored until analysis.

Cell counts were determined by flow cytometry using a BD Biosciences Influx high speed cell sorter. A 488nm laser was used in addition to chlorophyll and phycoerythrin filters, forward scatter relative to 1um beads, and side scattered light.

Data Processing Description

The program BD Software was used to collect flow cytometry data and generate distinct files for analysis. The program Flow Jo (version 7.6.5) was used to process flow cytometry files.

BCO-DMO Processing:

- modified parameter names to conform with BCO-DMO naming conventions;
- created date field with month and year (original dataset only had column for day of march);
- changed lon values from positive degrees west to negative degrees east.

[[table of contents](#) | [back to top](#)]

Data Files

File
SMILE_counts.csv (Comma Separated Values (.csv), 42.75 KB) MD5:e43b55e5f65ba380d26d2aa83090e15a
Primary data file for dataset ID 736702

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
castNum	Cast number	unitless
date	Year, month, and day of sampling (formatted as yyyy-mm-dd)	unitless
day_of_March	Day of the month	unitless
depth	Sample depth	meters
latitude	Latitude; positive values = North	decimal degree
longitude	Longitude; negative values = West	decimal degree
temperature	Temperature	degrees Celsius
salinity	Salinity	psu
oxygen	Oxygen	micromoles per kilogram (umol/kg)
PAR	PAR	unknown
prochlorococcus_per_mL	Prochlorococcus counts per mL	cells per milliliter
synechococcus_per_mL	Synechococcus counts per mL	cells per milliliter
picoeukaryotes_per_mL	Picoeukaryote counts per mL	cells per milliliter

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	CTD - profiler
Generic Instrument Description	The Conductivity, Temperature, Depth (CTD) unit is an integrated instrument package designed to measure the conductivity, temperature, and pressure (depth) of the water column. The instrument is lowered via cable through the water column. It permits scientists to observe the physical properties in real-time via a conducting cable, which is typically connected to a CTD to a deck unit and computer on a ship. The CTD is often configured with additional optional sensors including fluorometers, transmissometers and/or radiometers. It is often combined with a Rosette of water sampling bottles (e.g. Niskin, GO-FLO) for collecting discrete water samples during the cast. This term applies to profiling CTDs. For fixed CTDs, see https://www.bco-dmo.org/instrument/869934 .

Dataset-specific Instrument Name	BD Biosciences Influx high speed cell sorter
Generic Instrument Name	Flow Cytometer
Generic Instrument Description	Flow cytometers (FC or FCM) are automated instruments that quantitate properties of single cells, one cell at a time. They can measure cell size, cell granularity, the amounts of cell components such as total DNA, newly synthesized DNA, gene expression as the amount messenger RNA for a particular gene, amounts of specific surface receptors, amounts of intracellular proteins, or transient signalling events in living cells. (from: http://www.bio.umass.edu/micro/immunology/facs542/facswhat.htm)

[[table of contents](#) | [back to top](#)]

Deployments

SKQ201703S

Website	https://www.bco-dmo.org/deployment/736151
Platform	R/V Sikuliaq
Start Date	2017-03-03
End Date	2017-04-04
Description	Additional cruise information can be found at the Rolling Deck to Repository (R2R): http://www.rvdata.us/catalog/SKQ201703S

[[table of contents](#) | [back to top](#)]

Project Information

Microbial ecology of coexisting ecotypes: Are all Prochlorococcus equal? (ProEco)

Coverage: North Pacific Ocean, Station ALOHA

Description from the NSF award abstract:

Prochlorococcus is a photosynthetic organism that is tremendously abundant in the ocean and influences biogeochemical cycles on global scales. This project aims to link Prochlorococcus community structure to primary productivity in situ. The twelve known Prochlorococcus ecotypes exhibit extensive diversity. It is thought that this diversity allows the Prochlorococcus "collective" to maintain numerical dominance across gradients in light, nutrients, and temperature that accompany changes in depth, season, and latitude. A large gap in our understanding lies in whether we should assess the ecosystem value of Prochlorococcus by its abundance or by its community structure or both. Ecosystem models assign all ecotypes the same role. However, genomic and physiological evidence from cultivated isolates and wild populations suggests tentatively that distinct genotypes may contribute differently to the ecosystem through variation in light and nutrient physiologies and interactions with other microorganisms. The consequences of these molecular-level differences to primary productivity in situ are unknown. This project tests whether absolute abundance, or community structure, determines the contributions of Prochlorococcus to biogeochemical dynamics by measuring the contributions of different ecotypes to primary productivity. The results of this project will inform ecosystem models towards better representation of how shifts in climate and Prochlorococcus diversity will affect global nutrient cycles, trophic cascades, and interactions with other bacteria, viruses, and grazers. The insights and approaches delineated by this work will be generally applicable to the ecology of abundant microbial populations in the open ocean such as pigmented and non-pigmented eukaryotes, heterotrophic bacteria, and other cyanobacterial lineages. A basic understanding of differences between coexisting ecotypes will provide inroads into understanding mechanisms of cooperation, competition, and collaboration among ecotypes in all microbial ecosystems. The investigators will build a teaching module to expose high school students to microbial oceanography, big data, and systems biology through virtual ocean exploration. The primary objective will be to impress upon students the importance of an "invisible forest" of microorganisms in the ocean. Students will examine the distribution patterns of abundant microbial groups in the context of oceanographic data from large publically available databases. High school teachers and student interns, a graduate student, the investigators, and an educational specialist will design, implement, and test the module for classrooms nationwide. This effort will follow a successful education model (Systems Education Experience - SEE) developed previously.

The investigators will address an overarching hypothesis that Prochlorococcus ecotypes vary in their contribution to the ecosystem as primary producers. More specifically, the investigators hypothesize that patterns of cell division and carbon fixation vary between coexisting ecotypes, and these differences are a function of genome content, gene expression, environmental conditions, and community composition. The technical approach will involve two field-based experiments will be applied to three different depths, at the oceanographic Station ALOHA, that differ in Prochlorococcus community composition. Experiment 1 will examine whether coexisting ecotypes vary in cell division, using 16S rRNA sequencing to quantify ecotype abundance in G1, S, and G2 cells. Experiment 2 will examine how carbon fixation varies between coexisting ecotypes using RNA-stable isotope probing and 16S rRNA sequencing of RNA enriched in ¹³C after incubation with ¹³C-bicarbonate. These experiments will be performed with Prochlorococcus communities under native in situ conditions and shifts in conditions to mimic light and temperature of other depths. In both experiments, the temporal gene expression of a selected set of carbon fixation and cell division genes will be examined to link gene expression patterns to primary productivity. All data will be related to the oceanographic environment including its physical, chemical, and biological features.

[[table of contents](#) | [back to top](#)]

Program Information

Ocean Time-series Sites (Ocean Time-series)

Coverage: Bermuda, Cariaco Basin, Hawaii

Program description text taken from Chapter 1: Introduction from the **Global Intercomparability in a Changing Ocean: An International Time-Series Methods Workshop** report published following the workshop held November 28-30, 2012 at the Bermuda Institute of Ocean Sciences. The full report is available from the workshop Web site hosted by US OCB: <http://www.whoi.edu/website/TS-workshop/home>

Decades of research have demonstrated that the ocean varies across a range of time scales, with

anthropogenic forcing contributing an added layer of complexity. In a growing effort to distinguish between natural and human-induced earth system variability, sustained ocean time-series measurements have taken on a renewed importance. Shipboard biogeochemical time-series represent one of the most valuable tools scientists have to characterize and quantify ocean carbon fluxes and biogeochemical processes and their links to changing climate (Karl, 2010; Chavez et al., 2011; Church et al., 2013). They provide the oceanographic community with the long, temporally resolved datasets needed to characterize ocean climate, biogeochemistry, and ecosystem change.

The temporal scale of shifts in marine ecosystem variations in response to climate change are on the order of several decades. The long-term, consistent and comprehensive monitoring programs conducted by time-series sites are essential to understand large-scale atmosphere-ocean interactions that occur on interannual to decadal time scales. Ocean time-series represent one of the most valuable tools scientists have to characterize and quantify ocean carbon fluxes and biogeochemical processes and their links to changing climate.

Launched in the late 1980s, the US JGOFS (Joint Global Ocean Flux Study; <http://usjgofs.whoi.edu>) research program initiated two time-series measurement programs at Hawaii and Bermuda (HOT and BATS, respectively) to measure key oceanographic measurements in oligotrophic waters. Begun in 1995 as part of the US JGOFS Synthesis and Modeling Project, the CARIACO Ocean Time-Series (formerly known as the CARbon Retention In A Colored Ocean) Program has studied the relationship between surface primary production, physical forcing variables like the wind, and the settling flux of particulate carbon in the Cariaco Basin.

The objective of these time-series effort is to provide well-sampled seasonal resolution of biogeochemical variability at a limited number of ocean observatories, provide support and background measurements for process-oriented research, as well as test and validate observations for biogeochemical models. Since their creation, the BATS, CARIACO and HOT time-series site data have been available for use by a large community of researchers.

Data from those three US funded, ship-based, time-series sites can be accessed at each site directly or by selecting the site name from the Projects section below.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1646709

[[table of contents](#) | [back to top](#)]