

Microbial enzyme activities: peptidase activities of gravity-filtered seawater samples from the RV\Polarstern cruise ARKXXVII/3 in the Central Arctic Ocean and Laptev Sea, Aug-Sept. 2012

Website: <https://www.bco-dmo.org/dataset/742967>

Data Type: Cruise Results, experimental

Version: 1

Version Date: 2018-06-25

Project

» [Latitudinal and depth-related contrasts in enzymatic capabilities of pelagic microbial communities: Predictable patterns in the ocean?](#) (Patterns of activities)

Contributors	Affiliation	Role
Arnosti, Carol	University of North Carolina at Chapel Hill (UNC-Chapel Hill)	Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

This dataset includes microbial peptidase hydrolysis rates measured on particles collected from gravity filtered seawater. Links to archived CTD data are also provided. Five substrates linked to a 7-amido-4-methyl coumarin (MCA) fluorophore, one amino acid – leucine – and four oligopeptides – the chymotrypsin substrates alanine-alanine-phenylalanine (AAF) and alanine-alanine-proline-phenylalanine (AAPF), and the trypsin substrates glutamine-alanine-arginine (QAR) and glutamic acid-glycine-arginine (EGR) – were used to measure exo- and endo-acting peptidase activities, respectively.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:88.809 E:130.5795 S:79.6502 W:31.21

Temporal Extent: 2012-08-09 - 2012-09-22

Dataset Description

This dataset includes microbial peptidase hydrolysis rates measured on particles collected from gravity filtered seawater. Links to archived CTD data are also provided. Five substrates linked to a 7-amido-4-methyl coumarin (MCA) fluorophore, one amino acid – leucine – and four oligopeptides – the chymotrypsin substrates alanine-alanine-phenylalanine (AAF) and alanine-alanine-proline-phenylalanine (AAPF), and the trypsin substrates glutamine-alanine-arginine (QAR) and glutamic acid-glycine-arginine (EGR) – were used to measure exo- and endo-acting peptidase activities, respectively.

Methods & Sampling

Water was collected via Niskin bottles mounted on a rosette, equipped with a CTD.

Experiments on (operationally defined) particles were carried out by gravity-filtering water through 3 µm pore size filters. 1/12th sections of the 3 µm pore-size filters were submerged in 15 mL artificial seawater; enzyme activities were measured as described below.

Five substrates linked to a 7-amido-4-methyl coumarin (MCA) fluorophore, one amino acid – leucine – and four oligopeptides – the chymotrypsin substrates alanine-alanine-phenylalanine (AAF) and alanine-alanine-proline-phenylalanine (AAPF), and the trypsin substrates glutamine-alanine-arginine (QAR) and glutamic acid-glycine-arginine (EGR) – were used to measure exo- and endo-acting peptidase activities, respectively. Hydrolysis rates of the substrates were measured as an increase in fluorescence as the fluorophore was hydrolyzed from the substrate over time [as in Hoppe, 1993; Obayashi and Suzuki, 2005]. Incubations with the 5 low molecular weight substrates were set up in 4 ml cuvettes. For particle-associated incubations, 1/12th of a filter was submerged in 4 ml artificial seawater in a cuvettes; substrates were added at 100 micromolar concentrations. Fluorescence was measured at 0, 4, 8, and 12 h.

Substrate was added 100 micromolar concentrations. Fluorescence was measured over at 0 hrs, 24 h, 48h, and 72 h, using a mini fluorimeter. Hydrolysis rates were calculated from the rate of increase of fluorescence in the incubation over time relative to a set of standards of known concentration of fluorophore.

All incubations were conducted at 0 C in the dark.

Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- reduced decimal precision of rate columns from 9 to 6 places

[[table of contents](#) | [back to top](#)]

Data Files

File
ARK27-3_GF_MCA_joined.csv (Comma Separated Values (.csv), 18.33 KB) MD5:d5556faa46add62a6751414cd30f8a0d
Primary data file for dataset ID 742967

[[table of contents](#) | [back to top](#)]

Related Publications

Balmonte, J. P., Teske, A., & Arnosti, C. (2018). Structure and function of high Arctic pelagic, particle-associated and benthic bacterial communities. *Environmental Microbiology*, 20(8), 2941–2954. Portico. <https://doi.org/10.1111/1462-2920.14304>
Results

Hoppe, HG. (1993). Use of fluorogenic model substrates for extracellular enzyme activity (EEA) measurement of bacteria, p. 423-431. In P. F. Kemp, B. F. Sherr, E. B. Sherr, and J. J. Cole (ed.), *Handbook of methods in aquatic microbial ecology*. Lewis Publishers, Boca Raton, FL [978-0873715645](#)
Methods

Obayashi, Y., & Suzuki, S. (2005). Proteolytic enzymes in coastal surface seawater: Significant activity of endopeptidases and exopeptidases. *Limnology and Oceanography*, 50(2), 722–726.
doi:[10.4319/lo.2005.50.2.0722](https://doi.org/10.4319/lo.2005.50.2.0722)
Methods

Parameters

Parameter	Description	Units
station_no	refers to station number for cruise	unitless
depth_no	sequence of depths sampled (1 is surface; higher numbers at greater depths)	unitless
depth_m	actual depth at which water collected	meters
cast_no	cast number (refers to cast of CTD/Niskin bottles on cruise)	unitless
ISO_DateTime_UTC	date and time in ISO format (yyyy-mm-ddTHH:MM:SS)	unitless
Latitude	latitude; north is positive	decimal degeed
Longitude	longitude; east is postivie	decimal degeed
substrate	substrates for measurement of enzymatic activities. ara:arabinogalactan; chn:chondroitin sulfate; fuc:fucoidan; lam:laminarin ; pul:pullulan; xyl:xylan	unitless
timepoint	sampling point post-incubation	unitless
time_elapsed_hr	incubation time	hours
rep1_rate	replicate 1 hydrolysis rate	nanomoles/liter/hour (nmol L ⁻¹ h ⁻¹)
rep2_rate	replicate 2 hydrolysis rate	nanomoles/liter/hour (nmol L ⁻¹ h ⁻¹)
average	average of hydrolysis rates	nanomoles/liter/hour (nmol L ⁻¹ h ⁻¹)
std_dev	std deviation of hydrolysis rates	nanomoles/liter/hour (nmol L ⁻¹ h ⁻¹)
comments	url of CTD data in Pangaea database	unitless
filter_um	filter pore size for gravity filtration experiments	microns

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	CTD - profiler
Generic Instrument Description	The Conductivity, Temperature, Depth (CTD) unit is an integrated instrument package designed to measure the conductivity, temperature, and pressure (depth) of the water column. The instrument is lowered via cable through the water column. It permits scientists to observe the physical properties in real-time via a conducting cable, which is typically connected to a CTD to a deck unit and computer on a ship. The CTD is often configured with additional optional sensors including fluorometers, transmissometers and/or radiometers. It is often combined with a Rosette of water sampling bottles (e.g. Niskin, GO-FLO) for collecting discrete water samples during the cast. This term applies to profiling CTDs. For fixed CTDs, see https://www.bco-dmo.org/instrument/869934 .

Dataset-specific Instrument Name	
Generic Instrument Name	Fluorometer
Generic Instrument Description	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.

Dataset-specific Instrument Name	
Generic Instrument Name	Gel Permeation Chromatograph
Generic Instrument Description	Instruments that separate components in aqueous or organic solution based on molecular size generally for molecular weight determination. Gel permeation chromatography (GPC) is a type of size exclusion chromatography (SEC), that separates analytes on the basis of size.

Dataset-specific Instrument Name	
Generic Instrument Name	Niskin bottle
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[[table of contents](#) | [back to top](#)]

Deployments

ARK-XXVII-3

Website	https://www.bco-dmo.org/deployment/741293
Platform	R/V Polarstern
Start Date	2012-08-02
End Date	2012-10-08
Description	Project: Latitudinal and depth-related contrasts in enzymatic capabilities of pelagic microbial communities: Predictable patterns in the ocean? For other files related to this cruise, see https://www.pangaea.de/?q=ARK+XXVII%2F3 .

[[table of contents](#) | [back to top](#)]

Project Information

Latitudinal and depth-related contrasts in enzymatic capabilities of pelagic microbial communities: Predictable patterns in the ocean? (Patterns of activities)

Coverage: Atlantic Ocean, Arctic Ocean, Pacific Ocean, Greenland

NSF Award Abstract:

Heterotrophic microbial communities are key players in the marine carbon cycle, transforming and respiring organic carbon, regenerating nutrients, and acting as the final filter in sediments through which organic matter passes before long-term burial. Microbially-driven carbon cycling in the ocean profoundly affects the global carbon cycle, but key factors determining rates and locations of organic matter remineralization are unclear. In this study, researchers from the University of North Carolina at Chapel Hill will investigate the ability of pelagic microbial communities to initiate the remineralization of polysaccharides and proteins, which together constitute a major pool of organic matter in the ocean. Results from this study will be predictive on a large scale regarding the nature of the microbial response to organic matter input, and will provide a mechanistic framework for interpreting organic matter reactivity in the ocean.

Broader Impacts: This study will provide scientific training for undergraduate and graduate students from underrepresented groups. The project will also involve German colleagues, thus strengthening international scientific collaboration.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1332881

[[table of contents](#) | [back to top](#)]