

# [DEPRECATED] Averages and standard deviation of stable isotopes (d13C and d15N) across species for all macrofauna found on each substrate (carbonate rocks, woods, and bones) of the colonization experiment held in Mound 12

**Website:** <https://www.bco-dmo.org/dataset/747645>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2018-10-08

## Project

» [Collaborative research: Quantifying the biological, chemical, and physical linkages between chemosynthetic communities and the surrounding deep sea](#) (Costa Rica Seeps)

Contributors	Affiliation	Role
<a href="#">Levin, Lisa A.</a>	University of California-San Diego Scripps (UCSD-SIO)	Principal Investigator
<a href="#">Rouse, Gregory</a>	University of California-San Diego Scripps (UCSD-SIO)	Co-Principal Investigator
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset has been deprecated (obsoleted). It has been replaced by dataset <https://www.bco-dmo.org/dataset/747575>.

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Dataset Description

**Status (2021-02-09):** This dataset has been deprecated (obsoleted). It has been replaced by dataset <https://www.bco-dmo.org/dataset/747575>.

Averages and standard deviation of stable isotopes (d13C and d15N) across species for all macrofauna found on each substrate (carbonate rocks, woods, and bones) of the colonization experiment held in Mound 12. Substrates were collected using HOV Alvin on R/V Atlantis cruise AT37-13.

## Methods & Sampling

Carbonate rocks, woods, and bones deployed for 7 years at Mound 12 across seepage gradients were collected using HOV Alvin, and washed in milliQ water onboard. Before preserving, the sample was kept cold and animals were picked to sample tissue for isotope analysis. The tissue (aprox. 0.5 mg) was placed in pre weighted tin boats or combusted vials (500C overnight) and frozen at -70C. The remaining washed sample was sieved through 0.3 mm mesh, separating the sample in two fractions (a fine fraction with the meiofauna, and a coarser one with the macrofauna), both preserved in 96% Ethanol. In the laboratory, tissue samples were oven dried (60C) overnight, weighed and acidified with 12.5-25 ul 1N H3PO4 to remove inorganic C. Stable isotopes measurements (d13C and d15N) were carried out using a Costech elemental analyzer coupled to a

## Data Processing Description

BCO-DMO Processing:  
- modified parameter names.

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### Replaced by New Version

Levin, L. A., Rouse, G., Pereira, O. S. (2021) **Averages and standard deviation across species for all macrofauna found on each carbonate rock collected during R/V Atlantis cruise AT37-13 in the Pacific margin of Costa Rica from May to June 2017**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 2) Version Date 2021-02-17 doi:10.26008/1912/bco-dmo.747575.2 [ [view at BCO-DMO](#) ]

*Relationship Description: Dataset 747645 "AT37-13 Colonization Experiment Isotopes" has been replaced by dataset 747575 v2.*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
substrate	Type of substrate	unitless
alvin_dive	Number of the Alvin dive from which the sample was collected	unitless
habitat	Seepage activity (active or inactive)	unitless
d13C_avg	Average of carbon isotope signature of animals on that substrate	per mil
d15N_avg	Average of nitrogen isotope signature of animals on that substrate	per mil
d13C_stdev	Standard deviation of carbon isotope signature of animals on that substrate	per mil
d15N_stdev	Standard deviation of nitrogen isotope signature of animals on that substrate	per mil

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Costech elemental analyzer
<b>Generic Instrument Name</b>	Elemental Analyzer
<b>Dataset-specific Description</b>	Stable isotopes measurements (d13C and d15N) were carried out using a Costech elemental analyzer coupled to a Micromass Iso-prime isotope ratio mass spectrometer (EA/IRMS) at Washington State University.
<b>Generic Instrument Description</b>	Instruments that quantify carbon, nitrogen and sometimes other elements by combusting the sample at very high temperature and assaying the resulting gaseous oxides. Usually used for samples including organic material.

<b>Dataset-specific Instrument Name</b>	Micromass Iso-prime isotope ratio mass spectrometer
<b>Generic Instrument Name</b>	Isotope-ratio Mass Spectrometer
<b>Dataset-specific Description</b>	Stable isotopes measurements (d13C and d15N) were carried out using a Costech elemental analyzer coupled to a Micromass Iso-prime isotope ratio mass spectrometer (EA/IRMS) at Washington State University.
<b>Generic Instrument Description</b>	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

[ [table of contents](#) | [back to top](#) ]

## Deployments

### AT37-13

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/714567">https://www.bco-dmo.org/deployment/714567</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2017-05-20
<b>End Date</b>	2017-06-11
<b>Description</b>	More cruise information is available from Rolling Deck to Repository (R2R): * <a href="https://www.rvdata.us/search/cruise/AT37-13">https://www.rvdata.us/search/cruise/AT37-13</a> * <a href="https://doi.org/10.7284/907684">https://doi.org/10.7284/907684</a>

### AT37-13\_Alvin\_Dives

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/715760">https://www.bco-dmo.org/deployment/715760</a>
<b>Platform</b>	HOV Alvin
<b>Start Date</b>	2017-05-21
<b>End Date</b>	2017-06-08
<b>Description</b>	Collections of seep organisms in sediments and on rocks.

[ [table of contents](#) | [back to top](#) ]

## Project Information

### **Collaborative research: Quantifying the biological, chemical, and physical linkages between chemosynthetic communities and the surrounding deep sea (Costa Rica Seeps)**

**Coverage:** Costa Rica Pacific Margin

#### *NSF abstract:*

If life were to disappear from the deep sea, would we notice? We only have a cursory understanding of this vast region and the connectivity among its communities and the rest of the oceans, and yet the ecosystems of the deep sea have been implicated in the larger function of the global marine ecosystems. We now rely on the deep ocean for food, energy, novel drugs and materials, and for its role in the global cycling of carbon, as well as for supporting services such as habitat creation, nutrient replenishment for shallow waters, and the maintenance of biodiversity. Cold seeps, active areas of the seafloor where methane and other chemicals are released, are key features along the continental margins worldwide. To characterize how methane seep communities interact with the surrounding ecosystems and vice versa, we will study methane seeps off the Pacific coast of Costa Rica in 2017 and 2018. It is the sphere of influence around the seep, both along the seafloor and up into the water column, that we seek to better understand. We will map the structure and the chemistry surrounding these habitats using a novel 3-dimensional framework, combining typical transects with vertical characterizations of the water column just above the seafloor. This will include measurements of methane flux into the water column and changes in the overlying carbonate chemistry and oxygen levels that are critical to our understanding of the effect of warming, oxygen loss and ocean acidification in this region. Within this framework, we will collect seep organisms in sediments and on rocks (including all sizes from microbes to large animals), and transplant some of these from within the area of seep influence to the background deep sea, and vice-versa. Together, these studies will help us to measure the size of the seep sphere of influence, and also demonstrate the role of these seeps within the deep sea and the greater, global, marine ecosystem. We will share this information with a group of teachers during a series of workshops in the San Diego area, at an exhibit at the Birch Aquarium, and through the work of an artist who has worked extensively with marine organisms in extreme environments.

Chemosynthetic ecosystems are inextricably linked to the broader world-ocean biome and global biogeochemical cycles in ways that we are just beginning to understand. This research will identify the form, extent, and nature of the physical, chemical, and biological linkages between methane seeps and the surrounding deep-sea ecosystem. The proposed research builds critical understanding of the structural and functional processes that underpin the ecosystem services provided by chemosynthetic ecosystems. We target a critical continental margin, Costa Rica, where methane fates and dynamics loom large and play out in an setting that reflects many oceanographic stressors. We will use quantitative sampling and manipulative studies within a 3-dimensional oceanographic framework. We will ask what are the shapes of the diversity and density functions for organisms of different size classes and trophic position over the transition from the seep habitat through the ecotone to the background deep sea? Further, we will ask how do depth, dissolved oxygen concentrations, pH and carbonate ion availability, relative rates of fluid flux, and substrate (biogenic, authigenic carbonate, sediments) alter these linkages and interactions with the surrounding deep sea? Evidence for distinct transitional communities and biotic patterns in density and alpha and beta diversity will be quantified and placed in a global biogeographic context. All of these investigations will occur across biological size spectra: for microorganisms (archaea, bacteria, microeukaryotes), the macrofauna, and the megafauna that form biogenic habitats. Our research results will be interpreted in the context of potential effects of global ocean change in the equatorial Pacific to determine how the linkages with the surrounding deep sea will be altered as anthropogenic impacts proceed in the future.

#### **Related publications:**

Levin, L.A., V.J. Orphan, G.W. Rouse, W. Ussler, A. E. Rathburn, G. S. Cook, S. Goffredi, E. Perez, A. Waren, B. Grupe, G. Chadwick, B. Strickrott. (2012). A hydrothermal seep on the Costa Rica margin: Middle ground in a continuum of reducing ecosystems. *Proc. Royal Soc. B.* 279: 2580-88 doi: [10.1098/rspb.2012.0205](https://doi.org/10.1098/rspb.2012.0205)

Sahling, H., Masson, D. G., Ranero, C. R., Hühnerbach, V., Weinrebe, W., Klauke, I., & Suess, E. (2008). Fluid seepage at the continental margin offshore Costa Rica and southern Nicaragua. *Geochemistry, Geophysics, Geosystems* 9: doi: [10.1029/2008GC001978](https://doi.org/10.1029/2008GC001978)

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1634172</a>

[ [table of contents](#) | [back to top](#) ]