

Event log from R/V Sikuliaq SKQ201701S from January to February 2017

Website: <https://www.bco-dmo.org/dataset/755088>

Data Type: Cruise Results

Version: 1

Version Date: 2019-01-10

Project

» [Collaborative Research: A metabolic index to predict the consequences of climate change for midwater ecosystems](#) (Metabolic Index)

| Contributors | Affiliation | Role |
|---|---|---------------------------|
| Wishner, Karen | University of Rhode Island (URI) | Principal Investigator |
| Roman, Christopher Neil | University of Rhode Island (URI) | Co-Principal Investigator |
| Seibel, Brad | University of South Florida (USF) | Co-Principal Investigator |
| Biddle, Mathew | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

Event log from R/V Sikuliaq cruise SKQ201701S, which took place from January to February 2017.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:22.42 E:-116.5 S:21.22 W:-118.05

Temporal Extent: 2017-01-22 - 2017-02-11

Dataset Description

Date, time, and location of sampling operations.

Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- added date_MST and date.UTC to hold dates reformatted into yyyy-mm-dd.

Data Files

| File |
|--|
| eventlog.csv (Comma Separated Values (.csv), 12.40 KB) MD5:097302580ca7cb374c7e31dd2ca04fd8 Primary data file for dataset ID 755088 |

Related Publications

Childress, J. J., Barnes, A. T., Quetin, L. B., & Robison, B. H. (1978). Thermally protecting cod ends for the recovery of living deep-sea animals. *Deep Sea Research*, 25(4), 419–422. doi:[10.1016/0146-6291\(78\)90568-4](https://doi.org/10.1016/0146-6291(78)90568-4)
Methods

J. J. Childress, B. A. Seibel, Life at stable low oxygen levels: adaptations of animals to oceanic oxygen minimum layers. *J. Exp. Biol.* 201, 1223–1232 (1998). <http://jeb.biologists.org/content/jexbio/201/8/1223.full.pdf>
Methods

Roman, C., Ullman, D. S., Hebert, D., & Licht, S. (2019). The Wire Flyer Towed Profiling System. *Journal of Atmospheric and Oceanic Technology*, 36(2), 161–182. doi:[10.1175/jtech-d-17-0180.1](https://doi.org/10.1175/jtech-d-17-0180.1)
Methods

Wiebe, P. H., Morton, A. W., Bradley, A. M., Backus, R. H., Craddock, J. E., Barber, V., ... Flierl, G. R. (1985). New development in the MOCNESS, an apparatus for sampling zooplankton and micronekton. *Marine Biology*, 87(3), 313–323. doi:10.1007/bf00397811 <https://doi.org/10.1007/BF00397811>
Methods

Wishner, K. F., Seibel, B. A., Roman, C., Deutsch, C., Outram, D., Shaw, C. T., ... Riley, S. (2018). Ocean deoxygenation and zooplankton: Very small oxygen differences matter. *Science Advances*, 4(12), eaau5180. doi:[10.1126/sciadv.aau5180](https://doi.org/10.1126/sciadv.aau5180)
Results

Related Datasets

IsRelatedTo

Seibel, B., Roman, C., Wishner, K. (2021) **Respirometry data for pelagic crustaceans, cephalopods, and fish collected on R/V Sikuliaq cruise SKQ201701S from January to February 2017**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-07-22 doi:10.26008/1912/bco-dmo.855732.1 [[view at BCO-DMO](#)]

Wishner, K., Outram, D., Seibel, B., Roman, C. (2021) **Abundances of copepod species in each net from MOCNESS tows in the Eastern Tropical North Pacific collected on four research cruises from 2007-2017**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-07-09 doi:10.26008/1912/bco-dmo.855395.1 [[view at BCO-DMO](#)]

Parameters

| Parameter | Description | Units |
|-----------|-------------|-------|
| | | |

| | | |
|--------------------|--|-----------------|
| Event_no | event number | unitless |
| Gear_Type | gear type | unitless |
| Cast_no | cast number | unitless |
| Station | station | unitless |
| Tow_Type | tow type | unitless |
| Date_Local_MST | Date Local MST (MMDDYYYY) | unitless |
| Date_UTC | Date UTC (MMDDYYYY) | unitless |
| Time_In_Local_MST | Time In Local MST | unitless |
| Time_Out_Local_MST | Time Out Local MST | unitless |
| Time_In_UTC | Time In UTC | unitless |
| Time_Out_UTC | Time Out UTC | unitless |
| Lat_In | Lat in; North is positive, negative denotes South | decimal degrees |
| Lon_In | Lon in; East is positive, negative denotes West | decimal degrees |
| Lat_Out | Lat Out; North is positive, negative denotes South | decimal degrees |
| Lon_Out | Lon Out; East is positive, negative denotes West | decimal degrees |
| Day_Night | Day/Night | unitless |
| Min_Depth | min depth | meters (m) |
| Max_Depth | max depth | meters (m) |
| Comments | comments | unitless |

| | | |
|------------------|---|----------|
| date_time_in.UTC | date and time (UTC) in the water following ISO8601 yyyy-mm-ddTHH:MM | unitless |
| date_time_in.MST | date and time local (MST) in the water following ISO8601 yyyy-mm-ddTHH:MM | unitless |

[[table of contents](#) | [back to top](#)]

Instruments

| | |
|---|---|
| Dataset-specific Instrument Name | CTD |
| Generic Instrument Name | CTD Sea-Bird |
| Dataset-specific Description | CTD (Sikuliaq equipment, SeaBird rosette) |
| Generic Instrument Description | A Conductivity, Temperature, Depth (CTD) sensor package from SeaBird Electronics. This instrument designation is used when specific make and model are not known or when a more specific term is not available in the BCO-DMO vocabulary. Refer to the dataset-specific metadata for more information about the specific CTD used. More information from: http://www.seabird.com/ |

| | |
|---|--|
| Dataset-specific Instrument Name | MOCNESS |
| Generic Instrument Name | MOCNESS |
| Dataset-specific Description | MOCNESS (Wishner: Multiple Opening Closing Net and Environmental Sensing System (Wiebe et al. 1985), 1 m2 system, 8 or 9 nets, 222 µm mesh, updated software and Sea-Bird SBE911plusCTD system (SIO upgrades), day and night vertically- stratified and horizontally-sequenced tows, Wishner et al. 2018) |
| Generic Instrument Description | The Multiple Opening/Closing Net and Environmental Sensing System or MOCNESS is a family of net systems based on the Tucker Trawl principle. There are currently 8 different sizes of MOCNESS in existence which are designed for capture of different size ranges of zooplankton and micro-nekton Each system is designated according to the size of the net mouth opening and in two cases, the number of nets it carries. The original MOCNESS (Wiebe et al, 1976) was a redesigned and improved version of a system described by Frost and McCrone (1974). (from MOCNESS manual) |

| | |
|---|--|
| Dataset-specific Instrument Name | Wire Flyer |
| Generic Instrument Name | Sea-Bird SBE 49 FastCAT CTD Sensor |
| Dataset-specific Description | Wire Flyer (Roman development: oscillating deep towed hydrographic profiler with Sea-Bird 49 FastCAT CTD and Aanderaa 4831F oxygen sensor, Roman et al 2018) |
| Generic Instrument Description | The SBE 49 FastCAT is a CTD sensor for use in autonomous platforms. It contains a SBE 3P temperature sensor, a SBE 4C conductivity sensor and a strain-gauge pressure sensor as standard. It can operate in autonomous (16 Hz per sec) or polled mode (transmits each sample). The sensor is depth-rated to 350 m (plastic housing) or 7000 m (titanium housing). Accuracy: +/- 0.002 deg C (temperature), +/- 0.0003 S/m (conductivity), 0.1% of full scale range (pressure). |

| | |
|---|--|
| Dataset-specific Instrument Name | Tucker trawl net |
| Generic Instrument Name | Tucker Trawl |
| Dataset-specific Description | MTT + MOC (Seibel: large Tucker trawl net (3 m ² mouth, 100 m long) with MOCNESS software and sensors (Childress and Seibel 1998), large insulated closing cod end (Childress et al. 1978), live animal collections) |
| Generic Instrument Description | The original Tucker Trawl, a net with a rectangular mouth opening first built in 1951 by G.H. Tucker, was not an opening/closing system, but shortly thereafter it was modified so that it could be opened and closed. The original had a 183 cm by 183 cm flexible rectangular mouth opening 914 cm long net with 1.8 cm stretched mesh for the first 457 cm and 1.3 cm mesh for last 457 cm. 152 cm of coarse plankton or muslin netting lined the end of the net. Tucker designed the net to collect animals associated with the deep scattering layers, principally euphausiids, siphonophores, and midwater fish. (from Wiebe and Benfield, 2003). Currently used Tucker Trawls usually have 1-m ² openings and can have a single net or multiple nets on the frame. |

[[table of contents](#) | [back to top](#)]

Deployments

SKQ201701S

| | |
|--------------------|---|
| Website | https://www.bco-dmo.org/deployment/755461 |
| Platform | R/V Sikuliaq |
| Start Date | 2017-01-19 |
| End Date | 2017-02-15 |
| Description | See additional cruise information from R2R: https://www.rvdata.us/search/cruise/SKQ201701S |

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: A metabolic index to predict the consequences of climate change for

midwater ecosystems (Metabolic Index)

Coverage: Eastern Tropical North Pacific

Description from NSF award abstract:

With climate change, ocean temperatures are expected to increase which in turn will reduce oxygen availability and increase metabolic oxygen demand in marine organisms. The investigators will conduct shipboard physiological experiments for various marine organisms and determine their distributions in relation to environmental conditions within an oxygen minimum zone (OMZ) in the Eastern Pacific Ocean. The goal will be to model and map a Metabolic Index (MI) to predict how vertical and horizontal distributions for these species might change throughout the world's oceans in the future. The MI is defined as the ratio between environmental oxygen supply and temperature-dependent oxygen demand. Oxygen supply includes both the environmental oxygen concentration across a habitat range and the physiological features of organisms that facilitate oxygen uptake, such as gills and circulatory systems. Thus, the MI will integrate measured tolerance and environmental exposure to low oxygen with environmental data. The investigators will measure tolerance to low oxygen, focusing on under-studied organisms, including the effect of temperature and organism size. They will sample along a natural gradient in oxygen content south of the California Current in the Eastern Pacific. The science team and a videographer will develop a blog about deep-sea biology and climate change using web-based and video technologies. Four graduate students will be funded on this project, and in conjunction with a recently developed course in pelagic ecology, several undergraduates will have the opportunity to participate in seagoing research.

This research fills a critical need for a physiology-based metric that can be used to predict changing marine communities as the oceans warm and hypoxic zones expand. Modern OMZs are extensive and characterized by deep-water (300-800 m) oxygen partial pressures lethal to most marine organisms, yet thriving communities exist there. Climate change is predicted to further deplete oxygen. The investigators will model and map a Metabolic Index (MI) for diverse marine species to help predict how in vertical and horizontal distributions of species may change throughout the world's oceans in the future. The MI will derive oxygen supply and demand data from published and planned measurements of the minimum environmental partial pressure of oxygen to which individual species are exposed (based on their distributions in the water column) and the minimum requirements to support routine aerobic metabolic demand (from shipboard respiration measurements of individuals). During research cruises in the Eastern Pacific along a gradient of OMZ intensity, the investigators will conduct shipboard physiological measurements to determine metabolic demand for understudied mesozooplankton and gelatinous taxa and determine the size- and temperature dependence for diverse species for incorporation into the MI. Vertically-stratified net sampling and in situ photography will identify and characterize unique OMZ community features, such as the lower oxycline biomass peak present in some OMZs and the oxygen-dependence of day and night habitat depths for vertically-migrating species. The MI will be mapped using climatological data to both test and generate hypotheses about the response of oceanic communities to climate change. In preliminary analysis, the MI suggests a metabolic constraint at a MI of ~2 that may act to limit vertical and horizontal habitat ranges.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|-----------------------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1459243 |

[[table of contents](#) | [back to top](#)]