

# Sea urchin biomass at central and western Aleutian Islands, Alaska from visual surveys, July 2014

**Website:** <https://www.bco-dmo.org/dataset/755155>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2019-01-30

## Project

» [Ocean Acidification: Century Scale Impacts to Ecosystem Structure and Function of Aleutian Kelp Forests](#)  
(OA Kelp Forest Function)

## Program

» [Science, Engineering and Education for Sustainability NSF-Wide Investment \(SEES\): Ocean Acidification \(formerly CRI-OA\)](#) (SEES-OA)

| Contributors                       | Affiliation   | Role                      |
|------------------------------------|---|---------------------------|
| <a href="#">Steneck, Robert S.</a> | University of Maine (U Maine DMC)                   | Principal Investigator    |
| <a href="#">Estes, James A.</a>    | University of California-Santa Cruz (UCSC)          | Co-Principal Investigator |
| <a href="#">Rasher, Douglas B.</a> | Bigelow Laboratory for Ocean Sciences               | Co-Principal Investigator |
| <a href="#">Copley, Nancy</a>      | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager      |

## Abstract

Sea urchin biomass at central and western Aleutian Islands, Alaska from visual surveys, July 2014. Estimates were derived from visual surveys, performed via SCUBA.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Program Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:52.91344 E:-176.61505 S:51.88707 W:173.30659

**Temporal Extent:** 2014-07-04 - 2014-07-17

## Methods & Sampling

We characterized the ecological status of each island by quantifying the density, size frequency distribution, and biomass of the sea urchin community (primarily *Strongylocentrotus polyacanthus*) at randomly selected sites, using the same methods that have been employed by us and others over the past 30 years (Estes et al. 2010). Originally, we identified potential study sites by laying a grid over a map of each island, marking every place a grid line intersected the coast; these marks were later assigned GPS waypoints. During the 2014 research cruise, we randomly selected and resampled six sites per island, or in the Semichi Islands (Alaid, Nizki, and Shemya)—island group—as this level of sampling is sufficient to determine the ecological status of an island (Estes et al. 2010). We performed identical community surveys at the sites we studied with respect to algal reef bioerosion (see associated metadata forms and datasets).

At each site, a diver placed a 0.25-m<sup>2</sup> quadrat at 20 feet depth and counted all urchins within the quadrat, then collected the urchins in a bag. The diver then took a random number of kicks along the same depth contour and repeated this process until 20 quadrats were sampled or 200 urchins were collected, whichever occurred first. If 200 urchins were collected quickly, additional density counts were made to yield a better density estimate ( $n = 4$  minimum). Shipside, we measured the size (test diameter; mm) of each collected urchin with calipers. We then calculated its biomass using a known size-weight relationship (Estes et al. 2010). To estimate total urchin biomass for a site (grams per 0.25-m<sup>2</sup>), we summed the biomass of all urchins collected at the site and divided that sum by the number of quadrats deployed.

## Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- re-formatted date from m/d/yyyy to yyyy-mm-dd
- reduced precision from 9 decimal places to 5 for biomass\_convert and sum\_biomass\_class and from 9 to 3 places for urchin\_biomass\_grams

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

| File  |
|---|
| <b>island_biomass.csv</b> (Comma Separated Values (.csv), 116.33 KB)<br>MD5:db74ae3d3a97a47250e32c2892b10ece<br>Primary data file for dataset ID 755155 |

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Estes, J. A., Tinker, M. T., & Bodkin, J. L. (2010). Using Ecological Function to Develop Recovery Criteria for Depleted Species: Sea Otters and Kelp Forests in the Aleutian Archipelago. *Conservation Biology*, 24(3), 852–860. doi:[10.1111/j.1523-1739.2009.01428.x](https://doi.org/10.1111/j.1523-1739.2009.01428.x)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

| Parameter            | Description   | Units           |
|----------------------|---|-----------------|
| island               | name of island  | unitless        |
| site_name            | identity of site  | unitless        |
| latitude             | latitude of study site  | decimal degrees |
| longitude            | longitude of study site   | decimal degrees |
| depth_feet           | depth of benthic survey   | feet            |
| date                 | calendar date of survey formatted as yyyy-mm-dd   | unitless        |
| observer             | last name of observer   | unitless        |
| quadrat_count        | number of 0.25-m <sup>2</sup> quadrats from which estimates were derived  | quadrats        |
| urchin_diameter_mm   | diameter of urchin test (body)  | millimeters     |
| urchin_frequency     | number of urchins collected within a size class   | urchins         |
| size_biomass_convert | urchin biomass; calculated using known size to dry weight relationship:<br>=EXP(-7.857+(2.992*LN(urchin diameter)))     | grams           |
| sum_biomass_class    | total biomass of all urchins collected within a given size class  | grams           |
| urchin_biomass_grams | urchin biomass per 0.25 m <sup>2</sup> per size class; calculated by dividing<br>"sum.biomass.class" by "quadrat.count" | grams           |

[ [table of contents](#) | [back to top](#) ]

## Deployments

### PS1409

|                    |   |
|--------------------|---|
| <b>Website</b>     | <a href="https://www.bco-dmo.org/deployment/755184">https://www.bco-dmo.org/deployment/755184</a>   |
| <b>Platform</b>    | R/V Point Sur   |
| <b>Start Date</b>  | 2014-07-05  |
| <b>End Date</b>    | 2014-07-22  |
| <b>Description</b> | Benthic community studies associated with project "Project: Ocean Acidification: Century Scale Impacts to Ecosystem Structure and Function of Aleutian Kelp Forests". |

## Project Information

### **Ocean Acidification: Century Scale Impacts to Ecosystem Structure and Function of Aleutian Kelp Forests (OA Kelp Forest Function)**

*Extracted from the NSF award abstract:*

Marine calcifying organisms are most at risk to rapid ocean acidification (OA) in cold-water ecosystems. The investigators propose to determine if a globally unique and widespread calcareous alga in Alaska's Aleutian archipelago, *Clathromorphum nereostratum*, is threatened with extinction due to the combined effects of OA and food web alterations. *C. nereostratum* is a slow growing coralline alga that can live to at least 2000 years. It accretes massive 'bioherms' that dominate the regions' rocky substrate both under kelp forests and deforested sea urchin barrens. It develops growth bands (similar to tree rings) in its calcareous skeleton, which effectively record its annual calcification rate over centuries. Pilot data suggest the skeletal density of *C. nereostratum* began to decline precipitously in the 1990's in some parts of the Aleutian archipelago. The investigators now propose to use high-resolution microscopy and microCT imaging to examine how the growth and skeletal density of *C. nereostratum* has changed in the past 300 years (i.e., since the industrial revolution) across the western Aleutians. They will compare their records of algal skeletal densities and their variation through time with reconstructions of past climate to infer causes of change. In addition, the investigators will examine whether the alga's defense against grazing by sea urchins is compromised by ongoing ocean acidification. The investigators will survey the extent of *C. nereostratum* bioerosion occurring at 10 sites spanning the western Aleutians, both inside and outside of kelp forests. At each site they will compare these patterns to observed and monitored ecosystem trophic structure and recent *C. nereostratum* calcification rates. Field observations will be combined with laboratory experiments to determine if it is a decline in the alga's skeletal density (due to recent OA and warming), an increase in grazing intensity (due to recent trophic-level dysfunction), or their interactive effects that are likely responsible for bioerosion patterns inside vs. outside of forests. By sampling *C. nereostratum* inside and outside of forests, they will determine if kelp forests locally increase pH via photosynthesis, and thus buffer the effects of OA on coralline calcification. The combination of field observations with laboratory controlled experiments, manipulating CO<sub>2</sub> and temperature, will help elucidate drivers of calcification and project how these species interactions will likely change in the near future. The project will provide the first in situ example of how ongoing ocean acidification is affecting the physiology of long-lived, carbonate producing organisms in the subarctic North Pacific. It will also be one of the first studies to document whether OA, ocean warming, and food web changes to ecological processes are interacting in complex ways to reshape the outcome of species interactions in nature.

## Program Information

### **Science, Engineering and Education for Sustainability NSF-Wide Investment (SEES): Ocean Acidification (formerly CRI-OA) (SEES-OA)**

**Website:** [https://www.nsf.gov/funding/pgm\\_summ.jsp?pims\\_id=503477](https://www.nsf.gov/funding/pgm_summ.jsp?pims_id=503477)

**Coverage:** global

NSF Climate Research Investment (CRI) activities that were initiated in 2010 are now included under Science, Engineering and Education for Sustainability NSF-Wide Investment (SEES). SEES is a portfolio of activities that highlights NSF's unique role in helping society address the challenge(s) of achieving sustainability. Detailed information about the SEES program is available from NSF ([https://www.nsf.gov/funding/pgm\\_summ.jsp?pims\\_id=504707](https://www.nsf.gov/funding/pgm_summ.jsp?pims_id=504707)).

In recognition of the need for basic research concerning the nature, extent and impact of ocean acidification on oceanic environments in the past, present and future, the goal of the SEES: OA program is to understand

(a) the chemistry and physical chemistry of ocean acidification; (b) how ocean acidification interacts with processes at the organismal level; and (c) how the earth system history informs our understanding of the effects of ocean acidification on the present day and future ocean.

#### **Solicitations issued under this program:**

[NSF 10-530](#), FY 2010-FY2011

[NSF 12-500](#), FY 2012

[NSF 12-600](#), FY 2013

[NSF 13-586](#), FY 2014

NSF 13-586 was the final solicitation that will be released for this program.

#### **PI Meetings:**

[1st U.S. Ocean Acidification PI Meeting](#) (March 22-24, 2011, Woods Hole, MA)

[2nd U.S. Ocean Acidification PI Meeting](#) (Sept. 18-20, 2013, Washington, DC)

3rd U.S. Ocean Acidification PI Meeting (June 9-11, 2015, Woods Hole, MA – Tentative)

#### **NSF media releases for the Ocean Acidification Program:**

[Press Release 10-186 NSF Awards Grants to Study Effects of Ocean Acidification](#)

[Discovery Blue Mussels "Hang On" Along Rocky Shores: For How Long?](#)

[Discovery nsf.gov - National Science Foundation \(NSF\) Discoveries - Trouble in Paradise: Ocean Acidification This Way Comes - US National Science Foundation \(NSF\)](#)

[Press Release 12-179 nsf.gov - National Science Foundation \(NSF\) News - Ocean Acidification: Finding New Answers Through National Science Foundation Research Grants - US National Science Foundation \(NSF\)](#)

[Press Release 13-102 World Oceans Month Brings Mixed News for Oysters](#)

[Press Release 13-108 nsf.gov - National Science Foundation \(NSF\) News - Natural Underwater Springs Show How Coral Reefs Respond to Ocean Acidification - US National Science Foundation \(NSF\)](#)

[Press Release 13-148 Ocean acidification: Making new discoveries through National Science Foundation research grants](#)

[Press Release 13-148 - Video nsf.gov - News - Video - NSF Ocean Sciences Division Director David Conover answers questions about ocean acidification. - US National Science Foundation \(NSF\)](#)

[Press Release 14-010 nsf.gov - National Science Foundation \(NSF\) News - Palau's coral reefs surprisingly resistant to ocean acidification - US National Science Foundation \(NSF\)](#)

[Press Release 14-116 nsf.gov - National Science Foundation \(NSF\) News - Ocean Acidification: NSF awards \\$11.4 million in new grants to study effects on marine ecosystems - US National Science Foundation \(NSF\)](#)

[ [table of contents](#) | [back to top](#) ]

---

## **Funding**

| Funding Source                                | Award                       |
|---|-----------------------------|
| <a href="#">NSF Arctic Sciences (NSF ARC)</a> | <a href="#">PLR-1316141</a> |

[ [table of contents](#) | [back to top](#) ]