

Series 3A: Multiple stressor experiments on *T. pseudonana* (CCMP1014) - Dissolved Inorganic Carbon (DIC) measurements

Website: <https://www.bco-dmo.org/dataset/771333>

Data Type: experimental

Version: 1

Version Date: 2019-06-17

Project

» [Collaborative Research: Effects of multiple stressors on Marine Phytoplankton](#) (Stressors on Marine Phytoplankton)

Contributors	Affiliation	Role
Passow, Uta	University of California-Santa Barbara (UCSB-MSI)	Principal Investigator
Laws, Edward	Louisiana State University College of the Coast and Environment (LSU-CC&E [formerly SC&E])	Co-Principal Investigator
D'Souza, Nigel	University of California-Santa Barbara (UCSB-MSI)	Scientist, Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

The experiments were designed to test the combined effects of three CO₂ concentrations, four temperatures, and three light intensities on growth and photophysiology of the diatom *T. pseudonana* CCMP1014 in a multifactorial design. This dataset contains measurements of Dissolved Inorganic Carbon (DIC) made over the course of the experiments.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Temporal Extent: 2018-07-01 - 2018-10-31

Dataset Description

The experiments in Series 3A were designed to test the combined effects of three CO₂ concentrations, four temperatures, and three light intensities on growth and photophysiology of the diatom *T. pseudonana* CCMP1014 in a multifactorial design. This dataset reports the dissolved inorganic carbon concentrations measured during the experiments.

Methods & Sampling

Three CO₂ concentrations were tested: 410 ppm, 750 ppm, and 1000 ppm respectively. For each CO₂

concentration, four temperatures were tested: 15 degrees-C, 20 degrees-C, 25 degrees-C, and 30 degrees-C. Within each temperature, three light levels were tested: a sub-optimum light (SOL) intensity of 60 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, an optimum light (OL) intensity of 400 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ and an extreme light (EL) intensity of 800 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$. All lights were set at a 12 h day: 12 h dark cycle. For logistical reasons, experiments were partially conducted in series, with all light treatments at two temperatures (either 15 degrees-C and 25 degrees-C or 20 degrees-C and 30 degrees-C) running simultaneously. This was repeated for each CO₂ concentration.

Experiments were conducted in Multicultivator MC-1000 OD units (Photon Systems Instruments, Drasov, Czech Republic). Each unit consists of eight 85 ml test-tubes immersed in a thermostated water bath, each independently illuminated by an array of cool white LEDs set at specific intensity and timing. A 0.2 μm filtered CO₂-air mix (Praxair Distribution Inc.) was bubbled through sterile artificial seawater, and the humidified gas mix was supplied to each tube via gentle sparging through a 2 μm stainless steel diffuser. Flow rates were gradually increased over the course of the incubation to compensate for the DIC uptake of actively growing cells, and ranged from <0.04 Liters per minute (LPM) at the start of the incubations to 0.08 LPM in each tube after 2 days. For each CO₂ and temperature level, replication was achieved by incubating three tubes at sub-optimum light intensities, two tubes at optimum light intensity, and three tubes at extreme light intensities. Each experiment was split into two phases: An acclimation phase spanning 4 days, was used to acclimate cultures to their new environment. Pre-acclimated, exponentially-growing cultures were then inoculated into fresh media and incubated through a 3-day experimental phase during which assessments of growth, photophysiology, and nutrient cycling were carried out daily. All sampling started 5 hours into the daily light cycle to minimize the effects of diurnal cycles.

Experiments were conducted with artificial seawater (ASW) prepared using previously described methods (Kester et. al 1967), and enriched with nitrate (NO₃), phosphate (PO₄), silicic acid (Si[OH]₄), at levels ensuring that the cultures would remain nutrient-replete over the course of the experiment. Trace metals and vitamins were added as in f/2 (Guillard 1975). The expected DIC concentration and pH of the growth media was determined for the different pCO₂ and temperatures using the CO₂SYN calculator (Pierrot et al. 2006), with constants from Mehrbach et al. (1973, refit by Dickson & Millero 1987), and inputs of temperature, salinity, total alkalinity (2376.5 $\mu\text{mol} \cdot \text{kg}^{-1}$), pCO₂, phosphate, and silicic acid. DIC levels in ASW at the start of each phase of the experiments were manipulated by the addition of NaHCO₃, and was then maintained by bubbling a CO₂-Air mix through the cultures over the course of the experiments. The pH of the growth media was measured spectrophotometrically using the m-cresol purple method (Dickson 1993), and adjusted using 0.1N HCl or 0.1M NaOH. The media was distributed into 75 ml aliquots and each aliquot was inoculated with 5 ml of the *T. pseudonana* CCMP 1014 (TP1014) stock culture at the start of the experiments.

Dissolved Inorganic Carbon (DIC) measurements:

DIC was measured in freshly prepared media, and at the end of the experiment phase. 25 ml of the sample was siphoned into clean glass serum vials, fixed with HgCL₂ (0.035% final conc. v/v), and sealed with butyl rubber septa. Samples were stored at 4 degrees-C until analyzed. Prior experiments had confirmed that no gas exchange, and/or change in DIC occurred during sample storage for up to 30 days using this method. Total dissolved inorganic carbon (TCO₂) samples were analyzed using an automated infrared inorganic carbon analyzer (AIRICA). The AIRICA-23 (MARIANDA, Kiel, Germany), is a high precision instrument used to measure total dissolved inorganic carbon in seawater. The system uses a high precision syringe and a mass flow controller to deliver a known volume of sample into a stripper where it is then acidified, converting the inorganic carbon species into CO₂ and delivered under constant flow to nondispersive infrared detector. The CO₂ is then carried using an inert reference gas (N₂) into a LICOR-7000 that measures pCO₂ using the difference in infrared absorbance between a sample and reference cell. The pCO₂ is recorded over time and integrated by the AIRICA software. This integrated value is proportional to the amount of dissolved inorganic carbon evolved from the sample and converted to carbon units using a conversion factor (CT Factor). The CT Factor is determined by calibration of the system against a certified reference material of known value (Dickson et al. 2007. Guide to Best Practices for Ocean CO₂ Measurements). The value is converted to gravimetric units ($\mu\text{mol/kg}$) using the volume, temperature and salinity of the sample. In order to check for analytical stability of the system throughout a run, a certified reference material is used in between every 5 samples. Replicate DIC measurements were averaged.

Problem report: DIC measurements for two treatments (410 ppm CO₂, 20°C), and (410 ppm CO₂, 20°C) were flawed.

Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- reduced precision of pH values to 2 decimal places
- changed "- NA -" to "NA" ("not applicable")

[[table of contents](#) | [back to top](#)]

Data Files

File
3A_DIC.csv (Comma Separated Values (.csv), 3.58 KB) MD5:577e1aba3e15ad9577ee9f4ddc350797
Primary data file for dataset ID 771333

[[table of contents](#) | [back to top](#)]

Related Publications

Dickson, A. G. (1993). The measurement of sea water pH. *Marine Chemistry*, 44(2-4), 131-142.

doi:10.1016/0304-4203(93)90198-w [https://doi.org/10.1016/0304-4203\(93\)90198-W](https://doi.org/10.1016/0304-4203(93)90198-W)

Methods

Dickson, A. G., & Millero, F. J. (1987). A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research Part A. Oceanographic Research Papers*, 34(10), 1733-1743. doi:[10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5)

Methods

Dickson, A.G., Sabine, C.L. and Christian, J.R. (Eds.) 2007. Guide to best practices for ocean CO2 measurements. PICES Special Publication 3, 191 pp. ISBN: 1-897176-07-4. URL:

https://www.nodc.noaa.gov/ocads/oceans/Handbook_2007.html <https://hdl.handle.net/11329/249>

Methods

Guillard, R. R. L. (1975). Culture of Phytoplankton for Feeding Marine Invertebrates. *Culture of Marine Invertebrate Animals*, 29-60. doi:[10.1007/978-1-4615-8714-9_3](https://doi.org/10.1007/978-1-4615-8714-9_3)

Methods

Kester, D. R., Duedall, I. W., Connors, D. N., & Pytkowicz, R. M. (1967). Preparation of Artificial Seawater 1. *Limnology and Oceanography*, 12(1), 176-179. doi:[10.4319/lo.1967.12.1.0176](https://doi.org/10.4319/lo.1967.12.1.0176)

Methods

Mehrbach, C., Culberson, C. H., Hawley, J. E., & Pytkowicz, R. M. (1973). Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*, 18(6), 897-907. doi:[10.4319/lo.1973.18.6.0897](https://doi.org/10.4319/lo.1973.18.6.0897)

Methods

Pierrot, D. E. Lewis, and D. W. R. Wallace. 2006. MS Excel Program Developed for CO2 System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S.

Department of Energy, Oak Ridge, Tennessee. doi: [10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a](https://doi.org/10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a).

Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Phase	Indicates whether the sample was collected during the acclimation phase or the experiment phase of the experiment.	unitless
CO2	Indicates the concentration of CO2 in the CO2-Air mix that was bubbled through the samples over the course of the experiment	parts per million (ppm)
Temp	Indicates the temperature at which the samples were incubated.	degrees Celsius
Day	Indicates the timepoint (day) of sampling. D0 = day 0; D1 = day 1; etc.	unitless
Replicate	Indicates replication within a treatment. "NA" indicates "not applicable"	unitless
SOL_A	Dissolved Inorganic Carbon (DIC) measurements in replicate A incubated at sub optimum light (SOL)	micromol/kilogram
SOL_B	Dissolved Inorganic Carbon (DIC) measurements in replicate B incubated at sub optimum light (SOL)	micromol/kilogram
SOL_C	Dissolved Inorganic Carbon (DIC) measurements in replicate C incubated at sub optimum light (SOL)	micromol/kilogram
OL_A	Dissolved Inorganic Carbon (DIC) measurements in replicate A incubated at optimum light (OL)	micromol/kilogram
OL_B	Dissolved Inorganic Carbon (DIC) measurements in replicate B incubated at optimum light (OL)	micromol/kilogram
EL_A	Dissolved Inorganic Carbon (DIC) measurements in replicate A incubated at extreme light (EL)	micromol/kilogram
EL_B	Dissolved Inorganic Carbon (DIC) measurements in replicate B incubated at extreme light (EL)	micromol/kilogram
EL_C	Dissolved Inorganic Carbon (DIC) measurements in replicate C incubated at extreme light (EL)	micromol/kilogram

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Multicultivator MC-1000 OD (Photon Systems Instruments, Drasov, Czech Republic)
Generic Instrument Name	Cell Cultivator
Dataset-specific Description	Used for incubation of TP1014 cultures.
Generic Instrument Description	An instrument used for the purpose of culturing small cells such as algae or bacteria. May provide temperature and light control and bubbled gas introduction.

Dataset-specific Instrument Name	AIRICA-23 (MARIANDA, Kiel, Germany)
Generic Instrument Name	Inorganic Carbon Analyzer
Dataset-specific Description	An Automated infrared inorganic carbon analyzer (AIRICA) for analysis of dissolved inorganic carbon.
Generic Instrument Description	Instruments measuring carbonate in sediments and inorganic carbon (including DIC) in the water column.

Dataset-specific Instrument Name	LICOR-7000
Generic Instrument Name	Inorganic Carbon Analyzer
Dataset-specific Description	A CO ₂ /H ₂ O Analyzer for analysis of dissolved inorganic carbon.
Generic Instrument Description	Instruments measuring carbonate in sediments and inorganic carbon (including DIC) in the water column.

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Effects of multiple stressors on Marine Phytoplankton (Stressors on Marine Phytoplankton)

The overarching goal of this project is to develop a framework for understanding the response of phytoplankton to multiple environmental stresses. Marine phytoplankton, which are tiny algae, produce as much oxygen as terrestrial plants and provide food, directly or indirectly, to all marine animals. Their productivity is thus important both for global elemental cycles of oxygen and carbon, as well as for the productivity of the ocean. Globally the productivity of marine phytoplankton appears to be changing, but while we have some understanding of the response of phytoplankton to shifts in one environmental parameter at a time, like temperature, there is very little knowledge of their response to simultaneous changes in several parameters. Increased atmospheric carbon dioxide concentrations result in both ocean acidification and increased surface water temperatures. The latter in turn leads to greater ocean stratification and associated changes in light exposure and nutrient availability for the plankton. Recently it has become apparent that the response of phytoplankton to simultaneous changes in these growth parameters is not additive. For example, the effect of ocean acidification may be severe at one temperature-light combination and negligible at another. The researchers of this project will carry out experiments that will provide a theoretical understanding of the relevant interactions so that the impact of climate change on marine phytoplankton can be predicted in an informed way. This project will engage high schools students through training of a teacher and the development of a teaching unit. Undergraduate and graduate students will work directly on the research. A cartoon journalist will create a cartoon story on the research results to translate the findings to a broader general public audience.

Each phytoplankton species has the capability to acclimatize to changes in temperature, light, pCO₂, and nutrient availability - at least within a finite range. However, the response of phytoplankton to multiple simultaneous stressors is frequently complex, because the effects on physiological responses are interactive. To date, no datasets exist for even a single species that could fully test the assumptions and implications of existing models of phytoplankton acclimation to multiple environmental stressors. The investigators will combine modeling analysis with laboratory experiments to investigate the combined influences of changes in pCO₂, temperature, light, and nitrate availability on phytoplankton growth using cultures of open ocean and coastal diatom strains (*Thalassiosira pseudonana*) and an open ocean cyanobacteria species (*Synechococcus* sp.). The planned experiments represent ideal case studies of the complex and interactive effects of environmental conditions on organisms, and results will provide the basis for predictive modeling of the response of phytoplankton taxa to multiple environmental stresses.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1538602

[[table of contents](#) | [back to top](#)]