

# Nitrous oxide concentrations from the R/V Falkor expedition FK160115 in the Central Pacific from January to February 2016

**Website:** <https://www.bco-dmo.org/dataset/775849>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2022-06-08

## Project

» [The ProteOMZ Expedition: Investigating Life Without Oxygen in the Pacific Ocean](#) (ProteOMZ (Proteomics in an Oxygen Minimum Zone))

Contributors	Affiliation	Role
<a href="#">Santoro, Alyson E.</a>	University of Maryland Center for Environmental Science (UMCES/HPL)	Principal Investigator
<a href="#">Saito, Mak A.</a>	Woods Hole Oceanographic Institution (WHOI)	Co-Principal Investigator
<a href="#">Laperriere, Sarah Marie</a>	University of California-Santa Barbara (UCSB)	Scientist
<a href="#">Biddle, Mathew</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager
<a href="#">Soenen, Karen</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Dissolved N<sub>2</sub>O concentrations were measured in discrete samples on a research expedition to the Equatorial Pacific. Water samples were collected using a 24 bottle Niskin rosette equipped with a CTD. N<sub>2</sub>O concentrations were measured using a headspace equilibration method and analyzed on a SRI Greenhouse Gas Monitoring Gas Chromatograph.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:17 E:156 S:-10.563 W:139.8

**Temporal Extent:** 2016-01-17 - 2016-02-04

## Dataset Description

Falkor ProteOMZ nitrous oxide from expedition (FK160115) in the Central Pacific in 2016.

## Methods & Sampling

Samples were collected from the CTD rosette into 160 mL glass serum bottles.

Nitrous oxide concentration: N<sub>2</sub>O concentrations were measured using a headspace equilibration method and analyzed on a SRI Greenhouse Gas Monitoring Gas Chromatograph (GC) equipped with an electron capture detector (ECD), dual HayeSep D packed columns, and a 1-mL sample loop (SRI Instruments, Torrance, California, USA; Elkins 1980; Laperriere et al. 2019). N<sub>2</sub>O concentrations were calculated according to Walter et al. (2006).

## Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- combined date and time columns to form ISO Date format in column ISO\_DateTime\_UTC

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>n2o.csv</b> (Comma Separated Values (.csv), 14.72 KB) MD5:a939e48d4656a988f2ba3790acc425bc Primary data file for dataset ID 775849

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Saunders, J. K., McIlvin, M. R., Dupont, C. L., Kaul, D., Moran, D. M., Horner, T., Laperriere, S. M., Webb, E. A., Bosak, T., Santoro, A. E., & Saito, M. A. (2022). Microbial functional diversity across biogeochemical provinces in the central Pacific Ocean. *Proceedings of the National Academy of Sciences*, 119(37). <https://doi.org/10.1073/pnas.2200014119>  
*Results*

Walter, S., Bange, H. W., Breitenbach, U., & Wallace, D. W. R. (2006). Nitrous oxide in the North Atlantic Ocean. *Biogeosciences*, 3(4), 607–619. doi:[10.5194/bg-3-607-2006](https://doi.org/10.5194/bg-3-607-2006)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
ISO_DateTime_UTC	Date time time of cast following ISO 8601 convention in UTC	unitless
station	Station number	unitless
ctd	CTD cast number	unitless
latitude	latitude of station; North is positive; negative denotes South	decimal degrees
longitude	longitude station East is positive; negative denotes West	decimal degrees
niskin	Niskin bottle number	unitless
depth	depth of sampling	meters (m)
salinity	Salinity from CTD	practical salinity units (PSU)
temperature	Temperature from CTD	Celsius (C)
n2o_1	Nitrous oxide concentration replicate 1	nanomoles per liter (nmol/L)
n2o_2	Nitrous oxide concentration replicate 2	nanomoles per liter (nmol/L)
n2o_3	Nitrous oxide concentration replicate 3	nanomoles per liter (nmol/L)
n2o_4	Nitrous oxide concentration replicate 4	nanomoles per liter (nmol/L)
n2o_5	Nitrous oxide concentration replicate 5	nanomoles per liter (nmol/L)

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	CTD
<b>Generic Instrument Name</b>	CTD - profiler
<b>Dataset-specific Description</b>	Temperature and Salinity are from CTD
<b>Generic Instrument Description</b>	The Conductivity, Temperature, Depth (CTD) unit is an integrated instrument package designed to measure the conductivity, temperature, and pressure (depth) of the water column. The instrument is lowered via cable through the water column. It permits scientists to observe the physical properties in real-time via a conducting cable, which is typically connected to a CTD to a deck unit and computer on a ship. The CTD is often configured with additional optional sensors including fluorometers, transmissometers and/or radiometers. It is often combined with a Rosette of water sampling bottles (e.g. Niskin, GO-FLO) for collecting discrete water samples during the cast. This term applies to profiling CTDs. For fixed CTDs, see <a href="https://www.bco-dmo.org/instrument/869934">https://www.bco-dmo.org/instrument/869934</a> .

<b>Dataset-specific Instrument Name</b>	RI Greenhouse Gas Monitoring Gas Chromatograph
<b>Generic Instrument Name</b>	Gas Chromatograph
<b>Dataset-specific Description</b>	Nitrous oxide concentration: N2O concentrations were measured using a headspace equilibration method and analyzed on a SRI Greenhouse Gas Monitoring Gas Chromatograph (GC) equipped with an electron capture detector (ECD), dual HayeSep D packed columns, and a 1-mL sample loop (SRI Instruments, Torrance, California, USA; Elkins 1980; Laperriere et al. 2019).
<b>Generic Instrument Description</b>	Instrument separating gases, volatile substances, or substances dissolved in a volatile solvent by transporting an inert gas through a column packed with a sorbent to a detector for assay. (from SeaDataNet, BODC)

[ [table of contents](#) | [back to top](#) ]

## Deployments

### FK160115

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/708387">https://www.bco-dmo.org/deployment/708387</a>
<b>Platform</b>	R/V Falkor
<b>Report</b>	<a href="https://service.rvdata.us/data/cruise/FK160115/doc/FK160115_OfficialCruiseReport_Saito_v3.pdf">https://service.rvdata.us/data/cruise/FK160115/doc/FK160115_OfficialCruiseReport_Saito_v3.pdf</a>
<b>Start Date</b>	2016-01-16
<b>End Date</b>	2016-02-11
<b>Description</b>	Project: Using Proteomics to Understand Oxygen Minimum Zones (ProteOMZ) More information is available from the ship operator at <a href="https://schmidtoccean.org/cruise/investigating-life-without-oxygen-in-the...">https://schmidtoccean.org/cruise/investigating-life-without-oxygen-in-the...</a> Additional cruise information is available from the Rolling Deck to Repository (R2R): <a href="https://www.rvdata.us/search/cruise/FK160115">https://www.rvdata.us/search/cruise/FK160115</a>

[ [table of contents](#) | [back to top](#) ]

## Project Information

### The ProteOMZ Expedition: Investigating Life Without Oxygen in the Pacific Ocean (ProteOMZ (Proteomics in an Oxygen Minimum Zone))

**Website:** <https://schmidtocean.org/cruise/investigating-life-without-oxygen-in-the-tropical-pacific/#team>

**Coverage:** Central Pacific Ocean (Hawaii to Tahiti)

From Schmidt Ocean Institute's ProteOMZ Project page:

Rising temperatures, ocean acidification, and overfishing have now gained widespread notoriety as human-caused phenomena that are changing our seas. In recent years, scientists have increasingly recognized that there is yet another ingredient in that deleterious mix: a process called deoxygenation that results in less oxygen available in our seas.

Large-scale ocean circulation naturally results in low-oxygen areas of the ocean called oxygen deficient zones (ODZs). The cycling of carbon and nutrients – the foundation of marine life, called biogeochemistry – is fundamentally different in ODZs than in oxygen-rich areas. Because researchers think deoxygenation will greatly expand the total area of ODZs over the next 100 years, studying how these areas function now is important in predicting and understanding the oceans of the future. This first expedition of 2016 led by Dr. Mak Saito from the Woods Hole Oceanographic Institution (WHOI) along with scientists from University of Maryland Center for Environmental Science, University of California Santa Cruz, and University of Washington aimed to do just that, investigate ODZs.

During the 28 day voyage named “ProteOMZ,” researchers aboard R/V *Falkor* traveled from Honolulu, Hawaii to Tahiti to describe the biogeochemical processes that occur within this particular swath of the ocean’s ODZs. By doing so, they contributed to our greater understanding of ODZs, gathered a database of baseline measurements to which future measurements can be compared, and established a new methodology that could be used in future research on these expanding ODZs.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
Schmidt Ocean Institute (SOI)	<a href="#">R/V Falkor 160115 SOI ProteOMZ Expedition</a>
Alfred P. Sloan Foundation (Sloan)	<a href="#">FG-2016-7129</a>

[ [table of contents](#) | [back to top](#) ]