

Oxygen metrics for retinal function in marine invertebrate larvae determined with electroretinograms

Website: <https://www.bco-dmo.org/dataset/779425>

Data Type: experimental

Version: 1

Version Date: 2019-10-23

Project

» [Vision-mediated influence of low oxygen on the physiology and ecology of marine larvae](#) (Vision under hypoxia)

Contributors	Affiliation	Role
Levin, Lisa A.	University of California-San Diego (UCSD)	Principal Investigator
Oesch, Nicholas	University of California-San Diego (UCSD)	Co-Principal Investigator
Soenen, Karen	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Oxygen metrics for retinal function in marine invertebrate larvae determined with electroretinograms

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Oxygen metrics for retinal function in marine invertebrate larvae determined with electroretinograms.

Please see additional datasets for this paper, including the “RetinalResponse TimeSeries” and the “Experiment Metrics” datasets.

Methods & Sampling

Detailed methods can be found in McCormick, LR; Levin, LA; Oesch, NW. 2019. Vision is highly sensitive to oxygen availability in marine invertebrate larvae. Journal of Experimental Biology 222, jeb200899. doi:10.1242/jeb.200899. Data shown in Figure 1, bottom panel.

Briefly, the time series test recorded electroretinogram (ERG) responses to a 1 s square step of light at a constant irradiance of 3.56 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ repeated every 20 s, providing a nearly continuous measure of ERG response in a tethered, live larva during the experimental manipulation of partial pressure of oxygen ($p\text{O}_2$). There was a constant flow of pH-buffered sterile seawater in the chamber where the larva was held, and after a brief period in “normoxia” (surface-ocean oxygen levels), the $p\text{O}_2$ was decreased, and then held at a low $p\text{O}_2$ before re-oxygenating the solution. This dataset shows the three visual metrics that were calculated to quantify the effects of reduced $p\text{O}_2$ on retinal function. V90, V50, and V10 were calculated for each individual larva as the oxygen where there was 90%, 50%, and 10% retinal function remaining, in respect to retinal responses in normoxia (surface-ocean oxygen levels).

Oxygen was measured using a Microx4 (PreSens) oxygen meter and a Pst-7 oxygen optode probe.

Data Processing Description

All electrophysiology data was recorded and analyzed using Igor Pro 7 Software (Wavemetrics) using custom code. Oxygen data was analyzed using PreSens Measurement Studio 2. Post-processing analysis was completed in R Studio (version 3.3.3).

[[table of contents](#) | [back to top](#)]

Data Files

File
oxygen_metrics.csv (Comma Separated Values (.csv), 3.66 KB) MD5:e772a8884b27ef3b2db4112b29c6b3a8 Primary data file for dataset ID 779425

[[table of contents](#) | [back to top](#)]

Related Publications

McCormick, L. R., Levin, L. A., & Oesch, N. W. (2019). Vision is highly sensitive to oxygen availability in marine invertebrate larvae. The Journal of Experimental Biology, 222(10), jeb200899. doi:[10.1242/jeb.200899](https://doi.org/10.1242/jeb.200899)
Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Species	Species name.	unitless
Experiment_Name	Original experiment name that can be linked back to raw datafile collected in Igor.	unitless
Metric	One of three oxygen metrics for retinal function calculated in McCormick et al.	unitless
O2_umol_l	Oxygen concentration in umol O2/L of the given metric.	micromole per liter (umol/L)
O2_ml_l	Oxygen concentration in mL O2/L of the given metric.	milliliter per liter (ml/l)
O2_umol_kg	Oxygen concentration in umol O2/kg of the given metric.	micromole per kg (umol/kg)
O2_pO2	The partial pressure of oxygen of the given metric.	kiloPascale (kPa)

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	Oxygen Microelectrode Sensor
Dataset-specific Description	Oxygen was measured using a Microx4 (PreSens) oxygen meter and a Pst-7 oxygen optode probe.
Generic Instrument Description	Any microelectrode sensor that measures oxygen.

Project Information

Vision-mediated influence of low oxygen on the physiology and ecology of marine larvae (Vision under hypoxia)

Coverage: Southern California Bight, Northeast Pacific Ocean

NSF abstract:

Oxygen is being lost in the ocean worldwide as a result of ocean warming and the input of nutrients from land. Vision requires a large amount of oxygen, and may be less effective or require more light when oxygen is in short supply. This is especially true for active marine animals with complex eyes and visual capabilities, including active arthropods (crabs), cephalopods (squid), and fish. The California coastal waters exhibit a sharp drop in oxygen and light with increasing water depth. This project examines how visual physiology and ecology in young (larval) highly visual marine animals respond to oxygen loss, with a focus on key fisheries and aquaculture species. Experiments and observations will test the hypothesis that oxygen stress will change the light required for these organisms to see effectively, influencing the water depths where they can live and survive. The project will provide interdisciplinary experiences to students and an early career scientist and inform both the public (through outreach at the Birch Aquarium at Scripps Institution of Oceanography) and policy makers about the effects of oxygen decline in the ocean.

Negative effects of oxygen loss on vision have been described for humans and other terrestrial organisms, but never in the marine environment, despite the large changes in oxygen that can occur with depth and over time in the ocean, and the high metabolic demand of visual systems. This project will test the effects of low oxygen on vision in 3 combinations of eye design and photo-transduction mechanisms: compound eye with rhabdomeric photoreceptors (arthropods), simple eye with rhabdomeric photoreceptors (cephalopods), and simple eye with ciliary photoreceptors (fish). A series of oxygen- and light-controlled laboratory experiments will be conducted on representative taxa of each group including the tuna crab, *Pleuroncodes planipes*; the market squid, *Doryteuthis opalescens*, and the white sea bass, *Atractoscion nobilis*. In vivo electrophysiology and behavioral phototaxis experiments will identify new oxygen metrics for visual physiology and function, and will be compared to metabolic thresholds determined in respiration experiments. Hydrographic data collected over 3 decades by the CalCOFI program in the Southern California Bight will be evaluated with respect to visual and metabolic limits to determine the consequences of oxygen variation on the critical luminoxyscape (range of oxygen and light conditions required for visual physiology and function in target species) boundary in each species. Findings for the three vision-based functional groups may test whether oxygen-limited visual responses offer an additional explanation for the shoaling of species distributions among highly visual pelagic taxa in low oxygen, and will help to focus future research efforts and better understand the stressors contributing to habitat compression with expanding oxygen loss in the ocean.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1829623

[[table of contents](#) | [back to top](#)]