

GPX log of survey dives(?) from coastal reefs of Ormoc Bay, Leyte, Philippines, 2012-2018

Website: <https://www.bco-dmo.org/dataset/781652>

Data Type: Other Field Results

Version: 1

Version Date: 2019-11-06

Project

» [RAPID: Mega-typhoon impacts on the metapopulation resilience of coral reef fishes](#) (Reef Fish Resilience)

Contributors	Affiliation	Role
Pinsky, Malin	Rutgers University	Principal Investigator
Stuart, Michelle	Rutgers University	Co-Principal Investigator, Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

GPX log of survey dives from coastal reefs of Ormoc Bay, Leyte, Philippines, 2012-2018. GPX is a log of 15-second interval GPS locations during dives.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:11.0165 E:124.8083 S:10.6299 W:124.555

Temporal Extent: 2012-05-05 - 2018-04-10

Dataset Description

GPX log of survey dives from the west coast of Ormoc Bay, Leyte, the Philippines in the municipalities of Albueria (10.91667, 124.69667) and Bay Bay City (11.07611, 124.87528), 2012-2018. GPX is a log of 15-second interval GPS locations during dives.

Methods & Sampling

The GPS reader was floating on the surface and tethered to the divers. Location should not be considered exact. With currents and wind, the GPS reader could drift a number of meters away from directly overhead the divers.

Data entry was made with Excel or Google sheets, depending on the internet connection.

Data cleaning was performed in R using the tidyverse package.

Data Processing Description

BCO-DMO Data Processing:

- reformatted 'time' column to ISO_DateTime format (yyyy-mm-ddTHH:MM:SS)

[[table of contents](#) | [back to top](#)]

Data Files

File
gpx_log.csv (Comma Separated Values (.csv), 14.66 MB) MD5:868d8d741e87ea65bba2aeadc0c70a73 Primary data file for dataset ID 781652

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
lat	latitude reading from gps	decimal degrees
lon	longitude reading from gps	decimal degrees
ISO_DateTime_local	Local time of gps reading formatted as yyyy-mm-ddTHH:MM:SS	unitless
unit	number of gps unit	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Biomark 601 PIT tag reader
Generic Instrument Name	tracking tag
Dataset-specific Description	Passive Integrated Transponder (PIT) tags help scientists track individual organisms by providing a reliable lifetime 'barcode' for an individual animal. PIT tags are dormant until activated; they therefore do not require any internal source of power throughout their lifespan. To activate the tag, a low-frequency radio signal is emitted by a scanning device that generates a close-range electromagnetic field. The tag then sends a unique alpha-numeric code back to the reader (Keck 1994). Scanners are available as handheld, portable, battery-powered models and as stationary, automated models that are usually used for automated scanning.
Generic Instrument Description	Devices attached to living organisms with the purpose of determining the location of those organisms as a function of time after tagging and release.

Deployments

SCUBA Pinsky Leyte

Website	https://www.bco-dmo.org/deployment/642952
Platform	SCUBA Pinsky Leyte
Start Date	2012-05-05
End Date	2018-04-10
Description	Field seasons (SCUBA) in Leyte, Philippines to study coral reef fish resilience. West coast of Leyte, Philippines in the municipalities of Albuera (10.91667, 124.69667) and Bay Bay City (10.676940, 124.799170)

Project Information

RAPID: Mega-typhoon impacts on the metapopulation resilience of coral reef fishes (Reef Fish Resilience)

Coverage: West coast of Leyte Island, Visayas, Philippines

Description from NSF award abstract:

When Typhoon Haiyan hit the Philippines it had sustained winds of 305 to 315 kph and was the strongest storm ever to make landfall. Storms are one of the most important disturbances to coral reef ecosystems. Previous research has primarily emphasized that habitat recovery is important for the recovery of reef fish communities after disturbance. We understand little, however, about the role of larval dispersal in mediating species responses to disturbance. Reef fish function as metapopulations connected by larval dispersal among reefs, and larval connectivity is therefore a critical process for their dynamics. A field site directly in Typhoon Haiyan's path provides an ideal opportunity to address the role of larval dispersal during recovery. Over the course of four field seasons (2008 to 2013), nearly two thousand clownfish were surveyed along 20km of coastline. Clownfish possess the same basic life history as most reef fish (sedentary adults and pelagic larvae), but are sufficiently rare and visible that genetic parentage methods can be used to follow larval dispersal. This study site is therefore a unique location in which to understand the metapopulation impacts of a massive storm. This project will focus on three hypotheses: 1) Habitat destruction determines the short-term impacts of storms disturbance, 2) Metapopulation processes shape recolonization after disturbance, and 3) Disturbance allows rare competitors to increase in abundance. The project will address these questions with a combination of fixed and random transects to assess reef habitat and reef fish abundance and diversity, as well as detailed, spatially explicit surveys of anemones and clownfish. Genetic mark-recapture and parentage methods with yellowtail clownfish will pinpoint the origin of new recruits that recolonize the reef post-typhoon.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1430218