

# Elemental stoichiometry for *Emiliana huxleyi* across a range of 12 temperatures from 8.5-28.6C

**Website:** <https://www.bco-dmo.org/dataset/782921>

**Data Type:** experimental

**Version:** 1

**Version Date:** 2019-11-26

## Project

» [How does intensity and frequency of environmental variability affect phytoplankton growth?](#) (Enviro variability and phytoplankton growth)

Contributors	Affiliation	Role
<a href="#">Hutchins, David A.</a>	University of Southern California (USC)	Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset includes elemental stoichiometry for *Emiliana huxleyi* across a range of 12 temperatures from 8.5-28.6C. Global warming will be combined with predicted increases in thermal variability in the future surface ocean, but how temperature dynamics will affect phytoplankton biology and biogeochemistry is largely unknown. Here, we examine the responses of the globally important marine coccolithophore *Emiliana huxleyi* to thermal variations at two frequencies (1 d and 2 d) at low (18.5 °C) and high (25.5 °C) mean temperatures. Elevated temperature and thermal variation decreased growth, calcification and physiological rates, both individually and interactively. The 1 d thermal variation frequencies were less inhibitory than 2 d variations under high temperatures, indicating that high-frequency thermal fluctuations may reduce heat-induced mortality and mitigate some impacts of extreme high-temperature events. Cellular elemental composition and calcification was significantly affected by both thermal variation treatments relative to each other and to the constant temperature controls. The negative effects of thermal variation on *E. huxleyi* growth rate and physiology are especially pronounced at high temperatures. These responses of the key marine calcifier *E. huxleyi* to warmer, more variable temperature regimes have potentially large implications for ocean productivity and marine biogeochemical cycles under a future changing climate.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Temporal Extent:** 2017-01-01 - 2017-10-31

## Dataset Description

This dataset includes elemental stoichiometry for *Emiliana huxleyi* across a range of 12 temperatures from 8.5-28.6C. Global warming will be combined with predicted increases in thermal variability in the future surface ocean, but how temperature dynamics will affect phytoplankton biology and biogeochemistry is largely unknown. Here, we examine the responses of the globally important marine coccolithophore *Emiliana huxleyi* to thermal variations at two frequencies (1 d and 2 d) at low (18.5 °C) and high (25.5 °C) mean temperatures. Elevated temperature and thermal variation decreased growth, calcification and physiological rates, both

individually and interactively. The 1 d thermal variation frequencies were less inhibitory than 2 d variations under high temperatures, indicating that high-frequency thermal fluctuations may reduce heat-induced mortality and mitigate some impacts of extreme high-temperature events. Cellular elemental composition and calcification was significantly affected by both thermal variation treatments relative to each other and to the constant temperature controls. The negative effects of thermal variation on *E. huxleyi* growth rate and physiology are especially pronounced at high temperatures. These responses of the key marine calcifier *E. huxleyi* to warmer, more variable temperature regimes have potentially large implications for ocean productivity and marine biogeochemical cycles under a future changing climate.

These data are published in Wang, X., Fu, F., Qu, P., Kling, J. D., Jiang, H., Gao, Y., & Hutchins, D. A. (2019). How will the key marine calcifier *Emiliana huxleyi* respond to a warmer and more thermally variable ocean?. *Biogeosciences*, 16(22), 4393-4409. doi:10.5194/bg-2019-179.

## Methods & Sampling

Chlorophyll a, total particulate carbon (TPC), particulate organic carbon (POC), particulate organic nitrogen (PON), and particulate organic carbon (POP) were filtered onto GF/F filters and analyzed following the methodology used in Fu et al., 2007. Particulate inorganic carbon was defined as the difference between TPC and POC after POC filters had been subjected to concentrated HCl fumes for 24 hours to remove all inorganic carbon. Calcification, photosynthesis, and carbon fixation rates were all measured following the procedures outlined in Feng et al., 2008.

All data was processed using either R (v 3.4.4) or Microsoft Excel 2016.

## Data Processing Description

### BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- combined two Excel sheet (low and high temp tables)
- changed parameter names by removing spaces and units
- reduced precision from 14 to 4 decimal places for all values

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Ehux_thermal_curve_elemental_ratio.csv</b> (Comma Separated Values (.csv), 1.22 KB) MD5:731eb1894922579e41f2812992cb78b0
Primary data file for dataset ID 782921

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Feng, Y., Warner, M. E., Zhang, Y., Sun, J., Fu, F.-X., Rose, J. M., & Hutchins, D. A. (2008). Interactive effects of increased pCO<sub>2</sub>, temperature and irradiance on the marine coccolithophore *Emiliana huxleyi* (Prymnesiophyceae). *European Journal of Phycology*, 43(1), 87–98. doi:[10.1080/09670260701664674](https://doi.org/10.1080/09670260701664674)  
*Methods*

Fu, F.-X., Warner, M. E., Zhang, Y., Feng, Y., & Hutchins, D. A. (2007). Effects of Increased temperature and CO<sub>2</sub> on photosynthesis, growth, and elemental ratios in marine *Synechococcus* and *Prochlorococcus* (cyanobacteria). *Journal of Phycology*, 43(3), 485–496. doi:[10.1111/j.1529-8817.2007.00355.x](https://doi.org/10.1111/j.1529-8817.2007.00355.x)  
*Methods*

Wang, X., Fu, F., Qu, P., Kling, J. D., Jiang, H., Gao, Y., & Hutchins, D. A. (2019). How will the key marine calcifier

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Temperature	treatment temperature	degrees Celsius
TPC_PON	TPC to PON ratio (mol/mol)	unitless
PIC_POC	PIC to POC ratio (mol/mol)	unitless
Chla_POC	Chla to POC ratio	milligrams/gram
POC_PON	POC to PON ratio (mol/mol)	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	440 elemental analyzer (Costech Inc., CA)
<b>Generic Instrument Name</b>	Elemental Analyzer
<b>Dataset-specific Description</b>	Used to measure organic/inorganic carbon and nitrogen.
<b>Generic Instrument Description</b>	Instruments that quantify carbon, nitrogen and sometimes other elements by combusting the sample at very high temperature and assaying the resulting gaseous oxides. Usually used for samples including organic material.

<b>Dataset-specific Instrument Name</b>	Perkin Elmer (CA) Liquid Scintillation Counter
<b>Generic Instrument Name</b>	Liquid Scintillation Counter
<b>Dataset-specific Description</b>	Used to process radioactive assays.
<b>Generic Instrument Description</b>	Liquid scintillation counting is an analytical technique which is defined by the incorporation of the radiolabeled analyte into uniform distribution with a liquid chemical medium capable of converting the kinetic energy of nuclear emissions into light energy. Although the liquid scintillation counter is a sophisticated laboratory counting system used to quantify the activity of particulate emitting ( $\beta$ and $\alpha$ ) radioactive samples, it can also detect the Auger electrons emitted from $^{51}\text{Cr}$ and $^{125}\text{I}$ samples. Liquid scintillation counters are instruments assaying alpha and beta radiation by quantitative detection of visible light produced by the passage of rays or particles through a suitable scintillant incorporated into the sample.

[ [table of contents](#) | [back to top](#) ]

## Project Information

### How does intensity and frequency of environmental variability affect phytoplankton growth? (Enviro variability and phytoplankton growth)

**Coverage:** laboratory experiment

#### *NSF Award Abstract:*

Microscopic plants called phytoplankton are key members of global oceanic ecosystems, since their photosynthesis supports the majority of the marine food chain and produces about as much oxygen as land plants. Because of this, oceanographers have often carried out experiments examining how factors such as temperature and carbon dioxide levels may affect phytoplankton growth. Most previous experiments have used constant levels of temperature and carbon dioxide, but it is clear from looking at measurements from real ocean ecosystems that these two factors often vary greatly over timescales of days to weeks. Using field and laboratory experiments along with computer modeling, this project will test how the growth of several major groups of phytoplankton differs under constant conditions of temperature and carbon dioxide, compared to conditions in which these factors fluctuate in intensity and frequency. This research will give marine scientists a better picture of how phytoplankton may respond to a varying natural environment today and in the future, and therefore help us to understand how ocean food webs function to support critical living resources such as fisheries. The project will train graduate and undergraduate students and a postdoctoral researcher, and the lead scientists will be involved in an ocean science education program for largely minority high school students from a downtown Los Angeles school district.

The goal of this project is to use laboratory culture and natural community experiments to understand how realistically fluctuating temperature and  $\text{pCO}_2$  conditions may affect globally important phytoplankton groups in ways that differ from the artificial constant exposures used in previous work. Culture experiments will test how the intensity and frequency of short-term thermal and carbonate fluctuations affects the growth responses of diazotrophic and picoplanktonic cyanobacteria, coccolithophores, and diatoms under both current and projected future environmental conditions. These lab results will be supported and extended by parallel experiments using mixed natural assemblages from the California upwelling regime, allowing us to test these same questions using phytoplankton communities that experience large seasonal shifts between highly dynamic thermal and carbonate system conditions during the spring upwelling season, and relatively much more static conditions during fall stratification events. These results will be synthesized using a new generation of numerical models that employ novel approaches to incorporating realistic environmental variations to allow more accurate predictions of phytoplankton responses to a dynamic environment in today's marine ecosystems, and in the future changing ocean.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1538525</a>

[ [table of contents](#) | [back to top](#) ]