Mass spectrometry proteomics data investigating morphological plasticity in a sulfur-oxidizing bacterium from the SUP05 clade enhances dark carbon fixation from cultures grown under under aerobic and anaerobic conditions

Website: https://www.bco-dmo.org/dataset/819147

Data Type: experimental

Version: 1

Version Date: 2019-12-17

Project

» Characterizing the contribution of bacteria from the SUP05 clade to autotrophic and heterotrophic carbon cycling across ocean gradients (SUP05 Activities)

Contributors	Affiliation	Role
Morris, Robert	University of Washington (UW)	Principal Investigator
Ingalls, Anitra E.	University of Washington (UW)	Co-Principal Investigator
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
 - Methods & Sampling
- Related Publications
- Parameters
- <u>Instruments</u>
- Project Information
- Funding

Dataset Description

The mass spectrometry proteomics data files have been deposited to the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository with the data set identifier PXD013243 and title "Morphological plasticity in a sulfur-oxidizing bacterium from the SUP05 clade enhances dark carbon fixation"

(see https://www.ebi.ac.uk/pride/archive/projects/PXD013243).

These data were published in Shah et al. (2019).

Methods & Sampling

Methodology:

Ca. T. autotrophicus strain EF1 protein extracts were analyzed using a mass spectrometer. Protein identifications were determined using COMET and checked using PeptideProphet, ProteinProphet, and iProphet. The final list of protein identifications were then filtered to ensure that at least one peptide was observed consistently across individual injections and that the protein probability within each replicate was greater than 95%. A total of 802 proteins were observed, representing 53% of all predicted proteins in strain EF1.

Sampling and analytical procedures:

Protein samples for comparative analyses were extracted from axenic cultures grown under aerobic and anaerobic conditions.

Instruments:

Waters ACQUITY M-class UPLC coupled to a Thermo QExactive HF Orbitrap high-resolution mass spectrometer (HRMS) equipped with a nano-electrospray (NSI) source.

[table of contents | back to top]

Related Publications

Shah, V., Zhao, X., Lundeen, R. A., Ingalls, A. E., Nicastro, D., & Morris, R. M. (2019). Morphological Plasticity in a Sulfur-Oxidizing Marine Bacterium from the SUP05 Clade Enhances Dark Carbon Fixation. mBio, 10(3). doi:10.1128/mbio.00216-19 https://doi.org/10.1128/mBio.00216-19 Results

[table of contents | back to top]

Parameters

Parameters for this dataset have not yet been identified

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	Waters ACQUITY M-class UPLC coupled to a Thermo QExactive HF Orbitrap high-resolution mass spectrometer (HRMS) equipped with a nano-electrospray (NSI) source	
Generic Instrument Name	Mass Spectrometer	
Generic Instrument Description	General term for instruments used to measure the mass-to-charge ratio of ions; generally used to find the composition of a sample by generating a mass spectrum representing the masses of sample components.	

[table of contents | back to top]

Project Information

Characterizing the contribution of bacteria from the SUP05 clade to autotrophic and heterotrophic carbon cycling across ocean gradients (SUP05 Activities)


NSF Abstract:

Life in the ocean is dominated by diverse communities of microorganisms that are fueled by the remains of algae. The relative activity of different microbial community members can determine the extent of greenhouse gas capture, recycling of nutrients and food production by the ocean. Currently, there is a critical gap in our ability to quantify the balance between these processes. A major roadblock in our ability to understand the microbial players is our inability to grow even the most abundant organisms in the lab and study their physiology. With the recent cultivation of three bacteria common to the ocean, the possibility exists to make major advances in our understanding of their contribution to ocean processes and how they might change in the future. This project will characterize these three bacteria in culture to develop tools to determine their activities in the field. Because these bacteria are common in low oxygen environments, this work is particularly well poised to address how microbial communities respond to oxygen-depleted waters. This project will train graduate and undergraduate students in an important research area through collaboration with the University

of Washington Integrative Graduate Education and Research Traineeship program, the Program for Climate change and the University of Washington's Educational Outreach Program. Results will be presented at national and international meetings and through publication. All data will be freely available through the Biological and Chemical Oceanographic Data Management Office (BCO-DMO).

The balance between autotrophy and heterotrophy determines the size of the ocean's carbon sink and the flux of material and energy available to food chains. Despite efforts to identify the contributions of major bacterial and archaeal lineages to carbon fluxes, little is known about the environmental conditions that determine the underlying metabolisms controlling carbon flux. Members of the SUP05 clade of sulfur oxidizing gammaproteobacteria are abundant in the dark ocean and can dominate expanding suboxic and anoxic zones, where they have the potential to both produce and consume organic matter. The recent cultivation of 3 representatives from the SUP05 clade has revealed striking metabolic diversity within the clade. These isolates represent an unprecedented opportunity to elucidate the effects of carbon limitation, energy limitation and oxygen tension on the production and consumption of organic matter by a critical group of marine bacteria. The work proposed here would 1) determine the balance between autotrophic carbon fixation and heterotrophic organic matter assimilation in diverse cultured representatives under varying environmental conditions 2) determine the kinetics of reactions fueling autotrophy 3) characterize biomarkers along gradients of autotrophic and heterotrophic production induced by varying environmental conditions and 4) detect these molecular markers in the field along with cell abundances to determine the contribution of autotrophy, heterotrophy, sulfur oxidation and denitrification to SUP05 metabolism in a field site that transects an oxygen minimum zone in the North Pacific Ocean. This project will advance understanding of SUP05, a biological driver of elemental cycling in the mesopelagic, one of the largest habitats on earth and one that is experiencing large climate driven changes, including expansion of oxygen minimum zones.

More information about this project can be found on the Morris Lab page: http://morrislab.ocean.washington.edu/



SUP05 Bacteria (image courtesy of Robert Morris)

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1558483

[table of contents | back to top]