

Kelp forest mooring DIC, TA, pCO₂, and aragonite saturation state estimations inside the kelp canopy (36° 37.297' N, 121° 54.102' W.) at Hopkins Marine Station, recorded between June and October 2018.

Website: <https://www.bco-dmo.org/dataset/823008>

Data Type: Other Field Results

Version: 1

Version Date: 2020-09-02

Project

» [Collaborative Research: RUI: Building a mechanistic understanding of water column chemistry alteration by kelp forests: emerging contributions of foundation species](#) (Kelp forest biogeochemistry)

Contributors	Affiliation	Role
Nickols, Kerry J.	California State University Northridge (CSUN)	Principal Investigator, Project Coordinator
Dunbar, Robert B.	Stanford University	Co-Principal Investigator
Hirsh, Heidi	Stanford University	Scientist, Contact
Monismith, Stephen G.	Stanford University	Scientist
Mucciarone, David	Stanford University	Scientist
Takeshita, Yuichiro	Monterey Bay Aquarium Research Institute (MBARI)	Scientist
Traiger, Sarah	United States Geological Survey (USGS)	Scientist
Soenen, Karen	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Kelp forest mooring DIC, TA, pCO₂, and aragonite saturation state estimations inside the kelp canopy (36° 37.297' N, 121° 54.102' W.) at Hopkins Marine Station, recorded between June and October 2018.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: Lat:36.6216 Lon:-121.9017
Temporal Extent: 2018-06-07 - 2018-10-04

Dataset Description

These data are published in Hirsh *et al.*, see related publications section.

Methods & Sampling

In order to estimate other carbonate parameters from sensor pH data, we require an additional carbonate parameter. To accomplish this, we created a local empirical relationship to estimate total alkalinity (TA) based on discrete samples, and combined this with sensor data to calculate DIC, pCO₂, and aragonite saturation state (ΩAr).

TA was estimated using a multiple linear regression (MLR) approach, with temperature (T) and salinity (S) as inputs (Alin *et al.*, 2012; Carter *et al.*, 2018). This approach has been demonstrated to be effective in the open ocean (Carter *et al.*, 2018) and along the California Coast below 50 m depth (Alin *et al.*, 2012). Takeshita *et al.* (2015) demonstrated that this MLR approach can be extended to nearshore environments and showed TA could be estimated to $\pm 6 \mu\text{mol kg}^{-1}$ (RMSE) inside a kelp forest in Southern California by applying an offset to the equations from Alin *et al.* (2012).

Here, we fit an equation of the form used in Alin *et al.* (2012) to the discrete samples ($n = 271$) collected in the kelp forest, and obtained the following equation:

$$\text{TA est} = \alpha_0 + \alpha_1 (T - T_r) + \alpha_2 (S - S_r) + \alpha_3 (T - T_r) \times (S - S_r)$$

with a RMSE of $\pm 7.3 \mu\text{mol kg}^{-1}$. T_r and S_r are the mean temperature and salinity for the deployment. The estimated total alkalinity (TA_{est}) was combined with sensor pH to estimate DIC, pCO₂, and ΩAr using equilibrium constants from Lueker *et al.* (2000) and CO₂SYS (van Heuven *et al.*, 2011). For a range of pH from 7.56 to 8.30 the standard uncertainty for estimated carbonate system parameters was 8.98 to 13.43 $\mu\text{mol kg}^{-1}$ DIC, 54.22 to 9.42 μatm pCO₂, and 0.0501 to 0.223 ΩAr. These uncertainties were calculated using the 'errors' function in seacarb (Gattuso *et al.*, 2020, version 3.2.13 - Accessed 6 Mar 2020) with inputs of TA_{est} and measured pH, temperature and salinity over depth inside the kelp forest.

Data Processing Description

CO₂SYS (Matlab) was used to estimate DIC, pCO₂, and ΩAr from estimated TA and pH. The 'errors' function in seacarb (R) was used to estimate uncertainties.

BCO-DMO processing notes:

- Adjusted column headers to comply with database requirements
- Converted Timestamp to ISO format

[[table of contents](#) | [back to top](#)]

Data Files

File
carbonate.csv (Comma Separated Values (.csv), 22.84 MB) MD5:d80e450a7eff5be93ad4c4e350b6c7c3
Primary data file for dataset ID 823008

[[table of contents](#) | [back to top](#)]

Related Publications

Alin, S. R., Feely, R. A., Dickson, A. G., Hernández-Ayón, J. M., Juranek, L. W., Ohman, M. D., & Goericke, R. (2012). Robust empirical relationships for estimating the carbonate system in the southern California Current

System and application to CalCOFI hydrographic cruise data (2005-2011). Journal of Geophysical Research: Oceans, 117(C5), n/a-n/a. doi:10.1029/2011jc007511 <https://doi.org/10.1029/2011JC007511>
Methods

Carter, B. R., Feely, R. A., Williams, N. L., Dickson, A. G., Fong, M. B., & Takeshita, Y. (2017). Updated methods for global locally interpolated estimation of alkalinity, pH, and nitrate. Limnology and Oceanography: Methods, 16(2), 119-131. doi:[10.1002/lom3.10232](https://doi.org/10.1002/lom3.10232)
Methods

Hirsh, H. K., Nickols, K. J., Takeshita, Y., Traiger, S. B., Mucciarone, D. A., Monismith, S., & Dunbar, R. B. (2020). Drivers of Biogeochemical Variability in a Central California Kelp Forest: Implications for Local Amelioration of Ocean Acidification. Journal of Geophysical Research: Oceans, 125(11). Portico. <https://doi.org/10.1029/2020jc016320> <https://doi.org/10.1029/2020JC016320>
Results

Lavigne H, Gattuso JP (2014) Seacarb: seawater carbonate chemistry with R, R package version 3.0. Available from <http://CRAN.R-project.org/package=seacarb>
Methods

Lueker, T. J., Dickson, A. G., & Keeling, C. D. (2000). Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. Marine Chemistry, 70(1-3), 105-119. doi:10.1016/S0304-4203(00)00022-0 [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0)
Methods

Takeshita, Y., Frieder, C. A., Martz, T. R., Ballard, J. R., Feely, R. A., Kram, S., ... Smith, J. E. (2015). Including high-frequency variability in coastal ocean acidification projections. Biogeosciences, 12(19), 5853-5870. doi:[10.5194/bg-12-5853-2015](https://doi.org/10.5194/bg-12-5853-2015)
Methods

Van Heuven, S., Pierrot, D., Rae, J. W. B., Lewis, E., & Wallace, D. W. R. (2011). MATLAB Program Developed for CO₂ System Calculations. ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center (CDIAC). https://doi.org/10.3334/CDIAC/OTG.CO2SYS_MATLAB_V1.1 https://doi.org/10.3334/CDIAC/otg.CO2SYS_MATLAB_v1.1
Software

[[table of contents](#) | [back to top](#)]

Related Datasets

IsDerivedFrom

Hirsh, H., Nickols, K. J., Takeshita, Y., Traiger, S., Monismith, S. G., Mucciarone, D., Dunbar, R. B. (2020) **Data from moored instruments (pH, dissolved oxygen, temperature, salinity, PAR, pressure) at 9 depths outside and inside the kelp canopy at Hopkins Marine Station, recorded between June and October 2018.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2020-09-02 doi:10.26008/1912/bco-dmo.822549.1 [[view at BCO-DMO](#)]
Relationship Description: The derived carbonate system data comes from the kelp mooring data (pH, temperature, salinity)

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Depth_ID	ID to distinguish depth	unitless
Mooring_ID	Mooring name	unitless
Latitude	Latitude of mooring location, south is negative	decimal degrees
Longitude	Longitude of mooring location, west is negative	decimal degrees
MAB	Meters above bottom	meters (m)
DIC	Disolved Inorganic Carbon	micromoles per kilogram (umol/kg)
TA	Total Alkalinity	micromoles per kilogram (umol/kg)
Omega	Aragonite saturation state	aragonite (Ω_{Ar})
pCO2	Partial pressure of CO2	microatmosphere (uatm)
ISO_DateTime_UTC	Timestamp (date and time) in ISO format, UTC (yyyy-mm-ddThh:mmZ)	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	mFET and SeapHOx
Generic Instrument Name	pH Sensor
Dataset-specific Description	pH was directly measured (via mFET and SeapHOx) on the mooring at 6 depths (and estimated for 2 weeks at the surface based on the near-surface pH-O2 relationship). Please see metadata for kelp mooring data for additional pH details (https://www.bco-dmo.org/dataset/822549).
Generic Instrument Description	An instrument that measures the hydrogen ion activity in solutions. The overall concentration of hydrogen ions is inversely related to its pH. The pH scale ranges from 0 to 14 and indicates whether acidic (more H+) or basic (less H+).

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: RUI: Building a mechanistic understanding of water column chemistry alteration by kelp forests: emerging contributions of foundation species (Kelp forest biogeochemistry)

Coverage: Central California 36.6 N 122 W

NSF Award Abstract:

Kelp forest ecosystems are of ecological and economic importance globally and provide habitat for a diversity of fish, invertebrates, and other algal species. In addition, they may also modify the chemistry of surrounding waters. Uptake of carbon dioxide (CO₂) by giant kelp, *Macrocystis pyrifera*, may play a role in ameliorating the effects of increasing ocean acidity on nearshore marine communities driven by rising atmospheric CO₂. Predicting the capacity for kelp forests to alter seawater chemistry requires understanding of the oceanographic and biological mechanisms that drive variability in seawater chemistry. The project will identify specific conditions that could lead to decreases in seawater CO₂ by studying 4 sites within the southern Monterey Bay in Central California. An interdisciplinary team will examine variations in ocean chemistry in the context of the oceanographic and ecological characteristics of kelp forest habitats. This project will support an early career researcher, as well as train and support a postdoctoral researcher, PhD student, thesis master's student, and up to six undergraduate students. The PIs will actively recruit students from underrepresented groups to participate in this project through Stanford University's Summer Research in Geosciences and Engineering (SURGE) program and the Society for Advancement of Hispanics/Chicanos and Native Americans in Science (SACNAS). In addition, the PIs and students will actively engage with the management community (Monterey Bay National Marine Sanctuary and California Department of Fish and Wildlife) to advance products based on project data that will assist the development of management strategies for kelp forest habitats in a changing ocean.

This project builds upon an extensive preliminary data set and will link kelp forest community attributes and hydrodynamic properties to kelp forest biogeochemistry (including the carbon system and dissolved oxygen) to understand mechanistically how giant kelp modifies surrounding waters and affects water chemistry using unique high-resolution measurement capabilities that have provided important insights in coral reef biogeochemistry. The project sites are characterized by different oceanographic settings and kelp forest characteristics that will allow examination of relationships between kelp forest inhabitants and water column chemistry. Continuous measurements of water column velocity, temperature, dissolved oxygen, pH, and photosynthetically active radiation will be augmented by twice-weekly measurements of dissolved inorganic carbon, total alkalinity, and nutrients as well as periods of high frequency sampling of all carbonate system parameters. Quantifying vertical gradients in carbonate system chemistry within kelp forests will lead to understanding of its dependence on seawater residence time and water column stratification. Additional biological sampling of kelp, benthic communities, and phytoplankton will be used to 1) determine contributions of understory algae and calcifying species to bottom water chemistry, 2) determine contributions of kelp canopy growth and phytoplankton to surface water chemistry, and 3) quantify the spatial extent of surface chemistry alteration by kelp forests. The physical, biological, and chemical data collected across multiple forests will allow development of a statistical model for predictions of kelp forest carbonate system chemistry alteration in different locations and under future climate scenarios. Threshold values of oceanographic conditions and kelp forest characteristics that lead to alteration of water column chemistry will be identified for use by managers in mitigation strategies such as targeted protection or restoration.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1737096
NSF Division of Ocean Sciences (NSF OCE)	OCE-1737176

[[table of contents](#) | [back to top](#)]