

# Ichthyoplankton density and plankton displacement volume data collected using NOAA SEAMAP Bongo Plankton Nets on the NSF RAPID Plankton Cruises in the Northwestern Gulf of Mexico in 2017-2018

**Website:** <https://www.bco-dmo.org/dataset/829115>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2020-11-16

## Project

» [Collaborative Research: RAPID-HARVEY: Response of plankton assemblages and trophodynamics to a historic, hurricane-induced floodwater plume in a subtropical, pelagic environment](#) (RAPID Plankton)

Contributors	Affiliation	Role
<a href="#">Geist, Simon</a>	Texas A&M, Corpus Christi (TAMU-CC)	Principal Investigator
<a href="#">Robinson, Kelly L.</a>	University of Louisiana at Lafayette	Co-Principal Investigator, Contact
<a href="#">Schnetzer, Astrid</a>	North Carolina State University (NCSU)	Co-Principal Investigator
<a href="#">Stauffer, Beth</a>	University of Louisiana at Lafayette	Co-Principal Investigator
<a href="#">Bond, Pamela</a>	NOAA Southeast Fisheries Science Center (NOAA SEFSC)	Scientist
<a href="#">Drass, Denise</a>	NOAA Southeast Fisheries Science Center (NOAA SEFSC)	Scientist
<a href="#">Zapfe, Glenn</a>	NOAA Southeast Fisheries Science Center (NOAA SEFSC)	Scientist
<a href="#">McAskill, Shannan</a>	Texas A&M, Corpus Christi (TAMU-CC)	Student
<a href="#">Paxton, Stormy</a>	Texas A&M, Corpus Christi (TAMU-CC)	Student
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset includes ichthyoplankton density and plankton displacement volume data collected using NOAA SEAMAP Bongo Plankton Nets during three NSF RAPID Plankton cruises and one NOAA Plankton cruise. RAPID Plankton Cruise 1 took place on R/V Point Sur, cruise number PS1813, from October 28-November 3, 2017. RAPID Plankton Cruise 2 took place on R/V Pelican, cruise number PE18-18, from January 6- 12, 2018. RAPID Plankton Cruise 3 took place on R/V Pelican, cruise number PE18-21, from March 19- 23, 2018. The NOAA Plankton cruise took place on R/V Gordon Gunter, cruise number GU17-04, from September 04-30, 2017.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

---

## Coverage

**Spatial Extent:** N:29.5 E:-93.69 S:27.3 W:-95

**Temporal Extent:** 2017-09-22 - 2018-03-23

## Methods & Sampling

The Bongo Plankton Net was deployed according to the "NOAA SEAMAP Field Operations Manual for collection of data, Chapter V., National Marine Fisheries Service and Gulf States Marine Fisheries Commission, October 2001, Revision No.4."

Target Fishing Depth as much of the water column as possible stopping 2-4m above the bottom at bottom depths <200m, and upper 200m at bottom depths >200m; ship speed 1.5-2.0 knots; target wire angle of 45°; winch speeds during pay out and retrieving according to Chapter V., Table A.

Nets were rinsed and processed on board and samples were preserved in 200 proof Ethanol (ETOH) according to NOAA SEAMAP Operations Manual upon retrieval of nets. After 24 hours, ETOH was exchanged with fresh ETOH. Plankton samples were stored in 1000ml plastic jars until processing in the Geist Early Life History Research Laboratory at Texas A&M University Corpus Christi during 2017 and 2020. Total Zooplankton Displacement Volume was determined using a calibrated 1000ml measuring cylinder by subtracting the volume of ethanol without sample from the volume of ethanol with sample. The plankton sample was separated from ethanol using a concentrator sieve with a mesh size less than the net mesh size. Then the plankton sample was resuspended in Ethanol. Larval fish were picked under dissecting microscopes using feather steel tweezers. Every sample was looked through twice for quality check purposes.

## Data Processing Description

### Data Processing:

Filtered Volume was calculated according to the manufacturer manual of the flowmeter (General Oceanics):

Filtered Volume = distance towed \* net opening area,

with distance towed calculated as:

Flowmeter Count Difference \* Flowmeter Constant / 999999.

Total Plankton Displacement Volume and Total Larval Fish Density were standardized per 100m<sup>3</sup> filtered water volume and per area as under 10m<sup>2</sup> water column using the same formula as NOAA SEAMAP Pascagoula Office (POC G. Zapfe).

### BCO-DMO Processing:

- concatenated separate files into one dataset;
- converted date format to YYYY-MM-DD;
- removed commas from the FM\_CO field;
- added ISO date-time fields;
- added Cruise\_ID numbers.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>bongo.csv</b> (Comma Separated Values (.csv), 9.42 KB) MD5:4fb10fb5bcfa5bc89984a1768c1aebd
Primary data file for dataset ID 829115

## Related Publications

National Marine Fisheries Service and Gulf States Marine Fisheries Commission. (2001). SEAMAP Field Operations Manual for Collection of Data. Revision 4. <https://www.fisheries.noaa.gov/inport/item/53243>  
*Methods*

## Parameters

Parameter	Description	Units
Date	Date; format: YYYY-MM-DD	unitless
Cruise_ID	Cruise identifier	unitless
CR_N	Project Internal Cruise Name	unitless
P_SN	Project Internal Station Number	unitless
NOAA_SN	Respective NOAA SEAMAP/GOMECC Station Number	unitless
GEAR_T	Plankton Net Gear Type	unitless
MESH_W	Plankton Net Mesh Width	micrometers (um)
NET_NO	Net number, if gear has more than 1 net	unitless
NET_DIA	Diameter of the net	meters (m)
STA_LAT	Generalized Station Latitude	decimal degrees
STA_LON	Generalized Station Longitude	decimal degrees
BOT_D	Assigned Bottom Depth	meters (m)
MAX_ND	Max Net Depth	meters (m)
MIN_ND	Min Net Depth	meters (m)
D_TW	Depth of Tow $FD\_TW = MAX\_ND - MIN\_ND$	meters (m)

TGI_GMT	Time Gear In (GMT), Net starting fishing; format: hh:mm	unitless
TGO_GMT	Time Gear Out (GMT), Net stopping fishing; format: hh:mm	unitless
TGI_CST	Time Gear In (CST), Net starting fishing; format: hh:mm	unitless
TGO_CST	Time Gear Out (CST), Net stopping fishing; format: hh:mm	unitless
TT	Time towed, Duration of Net fishing; format: hh:mm:ss	unitless
FM_M	Flowmeter Manufacturer	unitless
FM_IC	Flowmeter initial count	unitless
FM_FC	Flowmeter final count	unitless
FM_CD	Flowmeter count difference	unitless
FM_CO	Flowmeter Constant, Manufacturer Supplied	unitless
DIS_T	Distance of Net towed through the water based on flowmeter readings	meters (m)
NET_OA	Net Opening Area	square meters (m <sup>2</sup> )
FIL_V	Filtered Volume	cubic meters (m <sup>3</sup> )
EV_WP	ETOH with Plankton (ml)	milliliters (ml)
EV_WOP	ETOH without Plankton (ml)	milliliters (ml)
PLA_D	Plankton Displacement Volume = ETOH with Plankton - ETOH without Plankton	milliliters (ml)
PLA_DD	Total Plankton Displacement Volume Density per 100 cubic meters filtered water volume	milliliters per 100 cubic meters (ml/100m <sup>3</sup> )
PLA_DA	Total Plankton Displacement Volume under 10m <sup>2</sup> area of the water column, $PLA\_DA = PLA\_D * ((FD\_TW / FIL\_V) * 10)$	milliliters under 10 square meters (ml/10m <sup>2</sup> )

TFILA	Total Larval Fish Number in sample	number of larval fish
TFILA_DV	Total Larval Fish Density per filtered water volume	larval fish per 100 cubic meters
TFILA_DA	Total Larval Fish Density under 10m <sup>2</sup> area, TFILA_DA=TFILA * ((FD_TW / FIL_V) * 10)	larval fish under 10 square meters (Ind./ 10m <sup>2</sup> )
TGI_ISO_DateTime_GMT	Date and Time Gear In (GMT), Net starting fishing; formatted to ISO8601 standard: YYYY-MM-DDThh:mmZ	unitless
TGO_ISO_DateTime_GMT	Date and Time Gear Out (GMT), Net stopping fishing; formatted to ISO8601 standard: YYYY-MM-DDThh:mmZ	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	NOAA SEAMAP Bongo Plankton Net
<b>Generic Instrument Name</b>	Bongo Net
<b>Dataset-specific Description</b>	NOAA SEAMAP Bongo Plankton Net (60cm opening diameter, 335µm mesh width), equipped with General Oceanics Mechanical Flowmeter. Online depth sensor (Seabird) was supplied by research vessel to allow safe deployment. One of the two nets was used to determine larval fish densities. The net was deployed over the side.
<b>Generic Instrument Description</b>	A Bongo Net consists of paired plankton nets, typically with a 60 cm diameter mouth opening and varying mesh sizes, 10 to 1000 micron. The Bongo Frame was designed by the National Marine Fisheries Service for use in the MARMAP program. It consists of two cylindrical collars connected with a yoke so that replicate samples are collected at the same time. Variations in models are designed for either vertical hauls (OI-2500 = NMFS Paironet-Style, MARMAP Bongo, CalVET) or both oblique and vertical hauls (Aquatic Research). The OI-1200 has an opening and closing mechanism that allows discrete "known-depth" sampling. This model is large enough to filter water at the rate of 47.5 m <sup>3</sup> /minute when towing at a speed of two knots. More information: Ocean Instruments, Aquatic Research, Sea-Gear

<b>Dataset-specific Instrument Name</b>	General Oceanics Mechanical Flowmeter
<b>Generic Instrument Name</b>	Mechanical Flowmeter
<b>Generic Instrument Description</b>	Manufactured by General Oceanics, a mechanical flow meter is used with plankton tows to determine the volume of water which flows through the net. Flow meters are also used in rivers, estuaries, canals, sewer outfalls, pipes, and harbor entrances to determine water velocity and flow distance information.

<b>Dataset-specific Instrument Name</b>	Dissecting microscope
<b>Generic Instrument Name</b>	Microscope - Optical
<b>Dataset-specific Description</b>	Larval fish were picked under dissecting microscopes using feather steel tweezers.
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### GU17-04

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/829174">https://www.bco-dmo.org/deployment/829174</a>
<b>Platform</b>	R/V Gordon Gunter
<b>Report</b>	<a href="https://datadocs.bco-dmo.org/docs/302/RAPID_Plankton/data_docs/GU-17-04_0097_SEAMAP_Fall_Plankton_Cruise_Report.pdf">https://datadocs.bco-dmo.org/docs/302/RAPID_Plankton/data_docs/GU-17-04_0097_SEAMAP_Fall_Plankton_Cruise_Report.pdf</a>
<b>Start Date</b>	2017-09-04
<b>End Date</b>	2017-09-30

### PS1813

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/827940">https://www.bco-dmo.org/deployment/827940</a>
<b>Platform</b>	R/V Point Sur
<b>Start Date</b>	2017-10-28
<b>End Date</b>	2017-11-03

### PE18-18

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/827941">https://www.bco-dmo.org/deployment/827941</a>
<b>Platform</b>	R/V Pelican
<b>Start Date</b>	2018-01-06
<b>End Date</b>	2018-01-13
<b>Description</b>	More cruise information is available from the Rolling Deck to Repository (R2R): <a href="https://www.rvdata.us/search/cruise/PE18-18">https://www.rvdata.us/search/cruise/PE18-18</a>

### PE18-21

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/827944">https://www.bco-dmo.org/deployment/827944</a>
<b>Platform</b>	R/V Pelican
<b>Start Date</b>	2018-03-18
<b>End Date</b>	2018-03-23
<b>Description</b>	More information is available from the Rolling Deck to Repository (R2R): <a href="https://www.rvdata.us/search/cruise/PE18-21">https://www.rvdata.us/search/cruise/PE18-21</a>

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

**Collaborative Research: RAPID-HARVEY: Response of plankton assemblages and trophodynamics to a historic, hurricane-induced floodwater plume in a subtropical, pelagic environment (RAPID Plankton)**

**Website:** <https://cajunplankton.com/nsf-rapid-plankton/>

**Coverage:** Northwestern Gulf of Mexico 95-96°W, 30-27.5°N

### *NSF Award Abstract:*

This project will examine how plankton in the northwestern Gulf of Mexico respond to large floodwater plumes generated by extreme weather events like Hurricane Harvey at time scales relevant to its development and evolution (days to months). The goal is to understand how the timing, magnitude, and constituent loads of a massive pulse of freshwater to the Louisiana-Texas shelf are: (1) driving changes in phytoplankton, zooplankton, and larval fish communities and distributions over monthly, seasonal, and annual time scales and, (2) what the consequences of those changes are to food web interactions within the plankton. The timing of Hurricane Harvey flood water disturbance coincides with the summer-fall spawning seasons for economically important Gulf of Mexico fisheries (e.g. red drum, sea trouts, snappers), raising additional questions of longer term effects of food web disruptions on recruitment. This project will train two undergraduate students and four PhD-level graduate students across three institutions, as well as support three early-career investigators. Pre- and post-floodwater plume data and samples will be shared with the broader scientific community within one year of collection to facilitate their immediate use by scientists beyond the research team. The team will give coordinated public talks at established regional science communication series and through other existing regional outreach partnerships to extend the educational scope of the project. Finally, results from this research will be incorporated in course curriculum and shared through scientific presentations and publications in peer-reviewed journals.

Record-breaking rain delivered by Hurricane Harvey to Southeast Texas in late August 2017 has resulted in a massive floodwater plume being delivered to coastal waters in the northern Gulf of Mexico (GOM). This project will investigate the short- and mid-term effects of that plume on planktonic (from pico- to ichthyoplankton) community composition and trophic interactions in that system. Building on data collected in July 2017 during a GOM Ecosystems and Carbon Cycle (GOMECC-3) cruise, and from historical datasets in the region, plankton assemblages, abundance, and food web interactions will be assessed during three research cruises 2, 4, and 8 months after the event. Oceanographic data and samples will be collected and processed using standard and state of the art gear, including ZooScan, FlowCAM, flow cytometry, and next generation sequencing. Onboard micro- and mesozooplankton grazing experiments will be conducted to understand the trophodynamic interactions and relationships between different plankton groups under changing environmental conditions. Diet and growth rate analyses of larval fish will be undertaken and related to phyto- and zooplankton (i.e. prey) abundance and community composition data. Application of the same gear types and methods during the three project cruises will ensure comparability of these new data to existing samples and datasets. These post-Harvey data will be compared to immediately-preceding and long-term data collected in the area by NOAA's Southeast Area Monitoring and Assessment Program (SEAMAP), allowing for investigation of the temporal evolution of planktonic assemblages and interpretation of plankton regime shifts in seasonal, multiyear, and decadal contexts.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1760674</a>

[ [table of contents](#) | [back to top](#) ]