

O2 saturation from microsensor profiles of ex situ sediment cores collected in the Chesapeake Bay and measured during 2017-2018

Website: <https://www.bco-dmo.org/dataset/847846>

Data Type: Other Field Results

Version: 1

Version Date: 2021-04-06

Project

» [Collaborative Research: Probing the Metabolic and Electrical Interactions of Cable Bacteria in Anoxic Sediments](#) (Anoxic Sediment Bacteria Interactions)

| Contributors | Affiliation | Role |
|--------------------------------|---|------------------------|
| Malkin, Sairah | University of Maryland Center for Environmental Science (UMCES/HPL) | Principal Investigator |
| Rauch, Shannon | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

This dataset includes O2 saturation from microsensor profiles of ex situ sediment cores collected in the Chesapeake Bay and measured during 2017-2018.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:38.55728 E:-76.42794 S:38.55505 W:-76.49402

Temporal Extent: 2017-03-28 - 2018-08-24

Methods & Sampling

Methodology:

High-resolution microsensor profiling of O₂, pH, and H₂S was performed on replicate sediment cores retrieved from the sampling sites, with 1 or 2 replicate profiles made per sediment core per analyte, using commercial microsensors operated with a motorized micromanipulator (Unisense A.S., Denmark). Oxygen sensor data were calibrated with a 2-point calibration using air saturated water and anoxic zone of sediments. Detailed methodology is given in Malkin et al., 2014.

Sampling and Analytical Procedures:

Replicate sediment cores were collected using a gravity corer (Uwitec; clear PVC liners, Ø = 8.6 cm), capped, kept in the dark at bottom water temperature in a water bath, and transported back to the laboratory, where they were held in a climate-controlled room. Microsensor profiling was conducted within 1 day of core retrieval.

Instruments:

Unisense A.S. microsenors (Denmark) and motorized micromanipulator system were used for data collection. The 4 channel microsensor multimeter was used, with the following microsenors: Ox-50, pH-200, and H2S-100. Each sensor was used consecutively and so resultant data is provided separately as individual datasets. SensorTrace Suite software was used for data collection.

Data Processing Description

BCO-DMO Processing:

- changed date format to YYYY-MM-DD;
- converted longitude from positive degrees West to negative degrees East.

[[table of contents](#) | [back to top](#)]

Data Files

| File |
|--|
| O2Sat.csv (Comma Separated Values (.csv), 195.23 KB) MD5:e6a0f32c2cd58c25436c93ae51110a0a Primary data file for dataset ID 847846 |

[[table of contents](#) | [back to top](#)]

Related Publications

Malkin, S. Y., Rao, A. M., Seitaj, D., Vasquez-Cardenas, D., Zetsche, E.-M., Hidalgo-Martinez, S., ... Meysman, F. J. (2014). Natural occurrence of microbial sulphur oxidation by long-range electron transport in the seafloor. The ISME Journal, 8(9), 1843–1854. doi:[10.1038/ismej.2014.41](https://doi.org/10.1038/ismej.2014.41)
Methods

[[table of contents](#) | [back to top](#)]

Related Datasets

IsRelatedTo

Malkin, S. (2021) **H2S from microsensor profiles of ex situ sediment cores collected in the Chesapeake Bay and measured during 2017-2018**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-04-07 doi:10.26008/1912/bco-dmo.847949.1 [[view at BCO-DMO](#)]

Malkin, S. (2021) **pH from microsensor profiles of ex situ sediment cores collected in the Chesapeake Bay and measured during 2017-2018**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-04-07 doi:10.26008/1912/bco-dmo.847923.1 [[view at BCO-DMO](#)]

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|-----------------|---|---------------------------|
| date_local | date of field sampling (local time zone; EST); format: YYYY-MM-DD | unitless |
| Year | year of field sampling | unitless |
| Month | month of field sampling | unitless |
| Day | day of month of field sampling | unitless |
| site | site of sediment collection | site |
| lat | latitude | degrees North |
| lon | longitude | degrees East |
| site_depth | depth of sampling site | meters (m) |
| CoreRep | core replicate (A or B) | unitless |
| ProfileRep | replicate profiles within a core (A or B) | unitless |
| sensor_depth_mm | depth of microsensor | millimeters (mm) |
| value_O2Sat | calibrated O2 saturation value of microsensor | percentage air saturation |
| Flag | indication of measurement problem or caution | unitless |

[[table of contents](#) | [back to top](#)]

Instruments

| | |
|---|---|
| Dataset-specific Instrument Name | gravity corer (Uwitec) |
| Generic Instrument Name | Gravity Corer |
| Generic Instrument Description | The gravity corer allows researchers to sample sediment layers at the bottom of lakes or oceans. The coring device is deployed from the ship and gravity carries it to the seafloor. (http://www.whoi.edu/instruments/viewInstrument.do?id=1079). |

| | |
|---|---|
| Dataset-specific Instrument Name | Unisense A.S. microsensors (Denmark) |
| Generic Instrument Name | Unisense oxygen microsensor |
| Dataset-specific Description | Unisense A.S. microsensors (Denmark) and motorized micromanipulator system were used for data collection. The 4 channel microsensor multimeter was used, with the following microsensors: Ox-50, pH-200, and H2S-100. Each sensor was used consecutively. |
| Generic Instrument Description | The Unisense oxygen microsensor is a miniaturized Clark-type dissolved oxygen instrument, including glass micro-sensors with minute tips (diameters ranging from 1 to 800 um). A gold sensing cathode is polarized against an internal reference and, driven by external partial pressure, oxygen from the environment penetrates through the sensor tip membrane and is reduced at the sensing cathode surface. A picoammeter converts the resulting reduction current to a signal. The sensor also includes a polarized guard cathode, which scavenges oxygen in the electrolyte, thus minimizing zero-current and pre-polarization time. See more on the manufacturer's website: https://www.unisense.com/ |

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Probing the Metabolic and Electrical Interactions of Cable Bacteria in Anoxic Sediments (Anoxic Sediment Bacteria Interactions)

Coverage: Chesapeake Bay sediments; Mid-Atlantic Coastal sediments

NSF Award Abstract:

Marine sediments represent the world's largest repository of stored organic carbon, and understanding how microorganisms break down this carbon is an imperative for understanding global carbon cycling. Yet long-standing questions remain regarding how networks of microorganisms work together to accomplish the complete breakdown of organic carbon in marine sediments. Sediment microbes interact in a myriad of ways that couple their metabolism to the break down of organic carbon, including by sharing products of metabolism. Accumulating evidence further suggests that some microorganisms can interact by transferring electrons directly to other unrelated microorganisms. This ability occurs across diverse microorganisms and appears to be widespread in the biosphere, particularly in anaerobic environments such as marine sediments. This project addresses emerging questions about the identity and metabolic linkages between microorganisms that work together in natural anaerobic marine and estuarine sediments to break down organic carbon. The investigators approach these questions by focusing on the influence of a keystone bacterium on its surrounding microbial community. "Cable bacteria" are a recently discovered group of long filamentous bacteria that act as electrical conductors in aquatic sediments providing a conduit for electrons to commute from deeper sulfidic sediments up to the surface oxygen layer by the process of centimeter-scale electron transport. Since their discovery about 6 years ago, these bacteria have been observed in a wide range of depositional sedimentary environments, often at extremely high cell densities. Where these bacteria are abundant, such as in coastal marine muds, they drive intense localized changes in pH and strongly influence the mineral cycling. This research explores the direct and indirect influence of cable bacteria on the metabolic activity of associated microorganisms. This project also advance the education and training of two early-career investigators, two PhD students, and undergraduate students. The skills and expertise gained from these PhD research projects will enable the students to be competitive in academic pursuits and in bioinformatics and technology applications relevant to private industry. The scientific discoveries emerging from this work is being incorporated into undergraduate and graduate level courses in marine microbial ecology. The research team will reach out to the broader community by hosting public lectures promoting a better understanding of environmental microbial ecology.

The proposed work is to investigate the role of cable bacteria in structuring sediment microbial communities.

Due to their growth strategy and morphology, cable bacteria are particularly amenable to experimental manipulation, providing an outstanding opportunity to better understand community interactions among microorganisms in a natural and complex anaerobic environment. The investigators will explore the interactions and relationships between cable bacteria and their associated microbial community by manipulating the growth and activity of cable bacteria and quantifying the resultant microbial community response. Specifically, this project aims to (1) identify microorganisms whose growth is enhanced by cable bacteria, (2) identify metabolic processes linked with cable bacteria activity using metatranscriptomics, (3) test specific metabolic links between sediment microorganisms and cable bacteria activity using a DNA-stable isotope probing (SIP) approach, and (4) visually confirm the identity and quantify key microorganisms associated with cable bacteria using microscopy. As more is learned about the identity and the mechanisms by which microorganisms are metabolically linked in anoxic sediments, we will be better able to understand and make predictions about how microorganisms function in their environment and how they can be utilized in bioengineered systems.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|-----------------------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1756877 |
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1756851 |

[[table of contents](#) | [back to top](#)]