

# T. rotula microbiome global sample

**Website:** <https://www.bco-dmo.org/dataset/860347>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2021-09-13

## Project

» [Connecting local, regional and global scales of gene flow in planktonic marine diatoms](#) (Diatom Gene Flow)

Contributors	Affiliation	Role
<a href="#">Rynearson, Tatiana A.</a>	University of Rhode Island (URI-GSO)	Principal Investigator
<a href="#">Ahern, Olivia</a>	University of Rhode Island (URI-GSO)	Contact
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset includes NCBI identifiers and descriptions of *T. rotula* microbiome global samples.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:51.18 E:174.81 S:-41.3 W:-125.09

**Temporal Extent:** 2010-01-26 - 2010-11-19

## Methods & Sampling

Single cells and chains of *T. rotula* were subjected to three serial washes in separate aliquots of 200  $\mu$ l sterile seawater, with <2  $\mu$ l seawater carried between washes. Individual cells or chains were then cultured with associated bacteria in 1ml f/20 in sterile polystyrene plates at *in situ* temperature and 90-120  $\mu$ mol photons  $m^{-2} s^{-1}$  on a 12:12-h light:dark cycle. Isolates were grown for two weeks, or until the phytoplankton concentration reached  $\sim 1,000$  cells  $mL^{-1}$ , whichever occurred sooner. Single cell isolates were filtered onto 0.2- $\mu$ m filters (Millipore) and stored at -80  $^{\circ}C$  until DNA extraction. DNA was extracted from *T. rotula* isolates and bacterial co-cultures following Whittaker and Ryneerson (2017).

## Data Processing Description

The V3-V4 rDNA region of the 16S rRNA gene was amplified from all samples using a dual index approach in 20  $\mu$ l reactions containing 1 $\mu$ l genomic DNA, 0.4 U of Taq (Lucigen) polymerase, and a final concentration of 200  $\mu$ M dNTPs, 2 mM MgCl<sub>2</sub>, and 0.5  $\mu$ M of each primer (16S V3 Forward - CCTACGGGNGGCWSCAG and 16S V4 Reverse - GGACTACNVGGGTWTCTAAT). The PCR was run for 30 seconds at 98  $^{\circ}C$ , followed by 30-35 cycles at 98  $^{\circ}C$  for 10 seconds, 55  $^{\circ}C$  for 30 sec and 72  $^{\circ}C$  for 30 sec, with a final extension at 72  $^{\circ}C$  for 2 min. Triplicate 20  $\mu$ l reactions were pooled and gel-purified for each sample.

Raw paired-end libraries were quality controlled using the ea-utils V.1.04.807 script fastq-mcf to a mate-pair quality mean of 25 with a 10 basepair (bp) sliding window and a minimum length of 150 bp. Forward and reverse reads were merged using the USEARCH V.8.1.1861 script fastq\_mergepairs with a minimum overlap length of 10 bp and zero differences in the overlap region followed by clustering using Minimum Entropy Decomposition (MED) V.2.0 with standard settings. Chloroplasts were filtered using the QIIME V.1.9.1 script filter\_fasta.py followed by an additional clustering step in MED with a minimum substantive abundance (-M) of 53 sequences required to form a single node. Chimeras were identified and removed using the USEARCH V.11.0.667 script uchime2\_ref in sensitive mode against the Silva v128 database. Silva database v128 was used to assign taxonomy to OTUs and libraries were then rarefied using the script multiple\_rarefactions\_even\_depth.py ten times to an even depth of 501 sequences per sample and OTU tables were merged using merge\_otu\_tables.py in QIIME V.1.9.1.

#### BCO-DMO Processing:

- flagged N/A as missing data identifier (replaced ;with nd);
- converted date format to YYYY-MM-DD
- renamed fields to conform with BCO-DMO naming conventions;
- removed commas from the Sample\_Site column.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>t_rotula_microbiome_global.csv</b> (Comma Separated Values (.csv), 12.58 KB) MD5:f65f7f7ac6b9bd465a21395e913c2f44
Primary data file for dataset ID 860347

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Ahern, O. M., Whittaker, K. A., Williams, T. C., Hunt, D. E., & Ryneerson, T. A. (2021). Host genotype structures the microbiome of a globally dispersed marine phytoplankton. *Proceedings of the National Academy of Sciences*, 118(48). <https://doi.org/10.1073/pnas.2105207118>

*Results*

Whittaker, K. A., & Ryneerson, T. A. (2017). Evidence for environmental and ecological selection in a microbe with no geographic limits to gene flow. *Proceedings of the National Academy of Sciences*, 114(10), 2651–2656. doi:[10.1073/pnas.1612346114](https://doi.org/10.1073/pnas.1612346114)

*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
SRA_Library_Name	Sample ID	unitless
Desc	Library description	unitless
Sample_Site	Sample site	unitless
Sample_Name	Timepoint identifier	unitless
Sample_date	Date of collection; format: YYYY-MM-DD	unitless
Lat	Latitude	degrees North
Long	Longitude	degrees East
Chl_a	Chlorophyll a concentration	milligrams per liter (mg/L)
Salinity	Sea surface salinity	parts per thousand (ppt)
SST	Sea surface temperature	degrees Celsius
T_rotula_Cell_Abundance	T. rotula cell abundance at time of collection	cells per milliliter (cells/mL)
Study_Accession	NCBI SRA accession for the project	unitless
Bioproject	NCBI BioProject ID number	unitless
SRA_Accession	NCBI SRA accession numbers	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Illumina MiSeq 2x250 bp PE reads
<b>Generic Instrument Name</b>	Automated DNA Sequencer
<b>Generic Instrument Description</b>	A DNA sequencer is an instrument that determines the order of deoxynucleotides in deoxyribonucleic acid sequences.

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Microscope - Optical
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Spectrophotometer
<b>Generic Instrument Description</b>	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Thermal Cycler
<b>Generic Instrument Description</b>	A thermal cycler or "thermocycler" is a general term for a type of laboratory apparatus, commonly used for performing polymerase chain reaction (PCR), that is capable of repeatedly altering and maintaining specific temperatures for defined periods of time. The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. They can also be used to facilitate other temperature-sensitive reactions, including restriction enzyme digestion or rapid diagnostics. (adapted from <a href="http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html">http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html</a> )

<b>Dataset-specific Instrument Name</b>	YSI
<b>Generic Instrument Name</b>	Water Quality Multiprobe
<b>Dataset-specific Description</b>	A YSI instrument measured SST and salinity.
<b>Generic Instrument Description</b>	An instrument which measures multiple water quality parameters based on the sensor configuration.

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Connecting local, regional and global scales of gene flow in planktonic marine diatoms (Diatom Gene Flow)

**Coverage:** global diatom samples; laboratory-based analyses

#### *Description from NSF award abstract:*

Diatoms are ubiquitous, unicellular, eukaryotes that generate about 40% of the organic carbon fixed annually in the sea. Interpretation of diatom species distributions and abundances in relation to environmental conditions has relied on two assumptions: (1) cells with identical morphologies represent the same species and (2) high potentials for dispersal and gene flow in passively drifting diatoms prevent local adaptation. Recent studies have challenged both assumptions, suggesting diatoms possess rich patterns of genetic and physiological variation both within and between species. Although there is emerging evidence of intra-specific population differentiation on local scales (~100km), it is commonly assumed that planktonic microbes are homogeneously distributed on global scales (e.g. Fenchel and Finlay 2004). There is currently no data on diatoms to support this assumption. Aside from intriguing data on local scales, nothing is known about regional and global-scale population genetics and biogeography of diatoms.

The research proposed here will focus on the essential questions of if and how populations of planktonic diatoms are connected at local, regional and global scales. Connectivity among populations can influence a species' ecology, adaptive potential, evolutionary longevity and ultimately speciation potential. The proposed research will examine how local populations are connected to each other on regional scales and how regional dynamics connect to global-scale biogeographies using two model diatom species. rDNA sequence variation will be used to test whether broad species distributions observed in diatoms result from cryptic speciation. Within species, microsatellite markers will be used to identify genetically distinct populations, determine their relatedness to each other and examine spatial patterns of differentiation. The degree of physiological variation that accompanies genetic differentiation between populations will also be examined. Samples will be collected in a framework of existing oceanography and biodiversity programs, permitting genetic data to be interpreted in the context of larger, often long-term, studies. Because little is known about diatom biogeography, this work will begin to shed light on the connections between local and global population dynamics. Because the proposed research will represent the first large-scale sampling of diatom population genetics, it will also serve to generate many new hypotheses about the mechanisms that regulate ecological processes such as bloom formation over space and time and evolutionary processes such as the development of reproductive isolation and eventual speciation in planktonic organisms.

#### *Related publications:*

Ryner, T.A., E.O. Lin and E.V. Armbrust. 2009. Metapopulation structure in the planktonic diatom *Ditylum brightwellii* (Bacillariophyceae). *Protist*, 160(1):111-121. doi:[10.1016/j.protis.2008.10.003](https://doi.org/10.1016/j.protis.2008.10.003)

Whittaker, K., Rignanes, D., Olson, R., Ryner, T., 2012. Molecular subdivision of the marine diatom *Thalassiosira rotula* in relation to geographic distribution, genome size, and physiology. *BMC Evolutionary Biology*, 12:209. doi:[10.1186/1471-2148-12-209](https://doi.org/10.1186/1471-2148-12-209)

Boyd, P.W., Ryner, T.A., Armstrong, E.A., Fu, F., Hayashi, K., Hu, Z., Hutchins, D.A., Kudela, R.M., Litchman, E., Mulholland, M.R., Passow, U., Strzepek, R.F., Whittaker, K.A., Yu, E., Thomas, M.K., 2013. Marine Phytoplankton Temperature versus Growth Responses from Polar to Tropical Waters - Outcome of a Scientific Community-Wide Study. *PLoS One*, 8(5), e63091. doi:[10.1371/journal.pone.0063091](https://doi.org/10.1371/journal.pone.0063091)

[ [table of contents](#) | [back to top](#) ]

---

## **Funding**

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0727227</a>

[ [table of contents](#) | [back to top](#) ]