

Phytoplankton and heterotrophic bacterial cell abundance in seawater collected during R/V Savannah cruise SAV-19-02 in the NW Atlantic Ocean in the Spring of 2019

Website: <https://www.bco-dmo.org/dataset/864302>

Data Type: Cruise Results

Version: 1

Version Date: 2021-11-02

Project

» [Collaborative Research: Assessing the role of compound-specific phosphorus hydrolase transformations in the marine phosphorus cycle](#) (P-hydrolase)

Contributors	Affiliation	Role
Duhamel, Solange	Columbia University	Principal Investigator
Diaz, Julia	University of Georgia (UGA)	Co-Principal Investigator
Djaoudi, Kahina	University of Arizona (UA)	Contact
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Phytoplankton and heterotrophic bacterial cell abundance in seawater collected during R/V Savannah cruise SAV-19-02 from March to April of 2019 in the Northwestern Atlantic from the surface to 50 m depth.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:31.7635 E:-79.8421 S:31.0175 W:-80.7965

Temporal Extent: 2019-03-30 - 2019-04-10

Methods & Sampling

Sampling and analytical procedures:

Seawater samples were collected in triplicate from Niskin bottles (12 L), at three stations (St. 1, St. 2 and St. 3) along a transect from coastal Georgia to offshore waters. Latitude (lat) and longitude (lon) of sampling sites are provided in .mat files. Stations 1 and 2 were on the continental shelf (15–30 m depth), while Station 3 was on the shelf break (220 m depth) adjacent to the Gulf Stream. At each station, samples were collected at 2–3 depths between the surface (4–5 m) and 50 m for Picoeukaryotes, Nano-eukaryotes, *Synechococcus* (PhytoplanktonCellAbundance.mat) and heterotrophic bacteria (HeterotrophicBacteriaCellAbundance.mat).

Samples for cell abundance (2 mL) were fixed with 0.5% glutaraldehyde (final concentration), flash frozen in liquid nitrogen and then transferred into a -80 °C freezer until analysis. Frozen samples were thawed at room

temperature and were analyzed using the Guava EasyCyte HT flow cytometer (Millipore). Instrument specific beads were used to calibrate the cytometer.

Samples were analyzed at a low flow rate ($0.24 \mu\text{L s}^{-1}$) during 3 min. For heterotrophic bacteria cell counts, samples were incubated with SYBR Green II solution 1:10 (Molecular Probes) for 15 min in the dark, in order to stain the nucleic acids. Bacterial cells were detected and enumerated based on diagnostic plots of forward scatter vs. green fluorescence. Group-specific phytoplankton were distinguished based on plots of forward scatter vs. orange fluorescence (phycoerythrin containing *Synechococcus* sp.), and SSC vs. red fluorescence (eukaryotes).

Instruments: Sampling was performed using Niskin bottles (12 L) mounted on a rosette. Cell abundance of both phytoplankton and heterotrophic bacteria were determined using the Guava EasyCyte HT flow cytometer (Millipore).

Location: Northwestern Atlantic surface waters. Depth: surface-50 m.

Data Processing Description

Data were organized using MATLAB and output as .mat files. Gaps in data were filled with NaN in the .mat files.

[[table of contents](#) | [back to top](#)]

Parameters

Parameters for this dataset have not yet been identified

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Guava EasyCyte HT flow cytometer (Millipore)
Generic Instrument Name	Flow Cytometer
Dataset-specific Description	Cell abundance of both phytoplankton and heterotrophic bacteria were determined using the Guava EasyCyte HT flow cytometer (Millipore).
Generic Instrument Description	Flow cytometers (FC or FCM) are automated instruments that quantitate properties of single cells, one cell at a time. They can measure cell size, cell granularity, the amounts of cell components such as total DNA, newly synthesized DNA, gene expression as the amount messenger RNA for a particular gene, amounts of specific surface receptors, amounts of intracellular proteins, or transient signalling events in living cells. (from: http://www.bio.umass.edu/micro/immunology/facs542/facswhat.htm)

Dataset-specific Instrument Name	
Generic Instrument Name	Niskin bottle
Dataset-specific Description	Sampling was performed using Niskin bottles (12 L) mounted on a rosette.
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[[table of contents](#) | [back to top](#)]

Deployments

SAV-19-02

Website	https://www.bco-dmo.org/deployment/864191
Platform	R/V Savannah
Start Date	2019-03-30
End Date	2019-04-11
Description	Cruise synonym: Zephyr (Zooming in on Enzymatic PhosphoHYdrolysis Reactions)

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Assessing the role of compound-specific phosphorus hydrolase transformations in the marine phosphorus cycle (P-hydrolase)

NSF Award Abstract:

Phosphorus (P) is an essential building block for life. Because P is in short supply over vast areas of the ocean, P availability may control biological productivity, such as photosynthesis and carbon fixation, which has implications for uptake of the greenhouse gas carbon dioxide and thus climate regulation. Marine microorganisms must satisfy their nutritional requirement for P by obtaining it from seawater, where P is present in a variety of chemical forms, from simple phosphate ions (Pi) to complex dissolved organic phosphorus (DOP) molecules. The concentration of DOP vastly exceeds Pi over most ocean areas, therefore DOP is a critically important source of P for marine microbial nutrition and productivity. However, much remains unknown about the contribution of specific DOP compounds to the P nutrition, productivity, and structure of marine microbial communities. In this project, the investigators will conduct field experiments in the Atlantic Ocean and perform a series of controlled laboratory studies with pure enzymes and microbial cultures to determine how and to what extent different DOP compounds are degraded to Pi in the marine environment. Furthermore, the contribution of these compound-specific DOP molecules to microbial P nutrition, carbon fixation, and community structure will be determined, thus advancing the current state of knowledge regarding the factors that control the activity and distribution of microbial species in the ocean, and the ocean's role in the climate system. This project will support two female junior investigators, a postdoctoral researcher, and graduate and undergraduate students. The undergraduate students will be recruited from the Marine Sciences program at Savannah State University, an Historically Black Colleges and Universities. In

addition, results will be incorporated into new hands-on K-12 educational tools to teach students about microbial P biogeochemistry, including a digital game and formal lesson plans with hands-on demos. These tools will be validated with K-12 educators and will be widely accessible to the public through various well-known online platforms. These activities will thus reach a broad audience including a significant fraction of underrepresented groups.

P is a vital nutrient for life. Marine microorganisms utilize P-hydrolases, such as alkaline phosphatase (AP), to release and acquire phosphate (Pi) from a wide diversity of dissolved organic P (DOP) compounds, including P-esters (P-O-C bonds), phosphonates (P-C), and polyphosphates (P-O-P). Compound-specific DOP transformations have the potential to exert critical and wide-ranging impacts on marine microbial ecology (e.g. variable DOP bioavailability among species), biogeochemistry (e.g. P geologic sequestration via formation of calcium Pi), and global climate (e.g. aerobic production of the greenhouse gas methane by dephosphorylation of methylphosphonate). However, the mechanisms and comparative magnitude of specific DOP transformations, in addition to their relative contributions to microbial community-level P demand, productivity, and structure, are not completely understood. This study will fill these knowledge gaps by tracking the fate of specific DOP pools in the marine environment. Specifically, this project will test four hypotheses in the laboratory using recombinant enzymes and axenic cultures representative of marine eukaryotic and prokaryotic plankton from high and low nutrient environments, and in the field using observational and experimental approaches along natural Pi gradients in the Atlantic Ocean. In particular, the investigators will reveal potential differences in the hydrolysis and utilization of specific DOP compounds at the community- (bulk enzymatic assays), taxon- (cell sorting of radiolabeled cells in natural samples), species- (axenic cultures) and molecular-levels (pure enzyme kinetic studies and cell-associated proteomes and exoproteomes). Results from our proposed work will provide a robust understanding of the enzymatic basis involved in the transformation of specific forms of DOP and create new knowledge on the relative contribution of these specific P sources to Pi production, marine microbial nutrition, community structure, primary productivity, and thus global carbon cycling and climate. In particular, our refined measurements of the concentration of bioavailable DOP and our unique estimates of DOP remineralization fluxes will provide critical new information to improve models of marine primary production and P cycling.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1736967
NSF Division of Ocean Sciences (NSF OCE)	OCE-1737083
NSF Division of Ocean Sciences (NSF OCE)	OCE-2001212
NSF Division of Ocean Sciences (NSF OCE)	OCE-1948042

[[table of contents](#) | [back to top](#)]