

# Phytoplankton nitrate uptake rates collected during the 2012-2013 Palmer Field Season (WAP Carbon export project)

**Website:** <https://www.bco-dmo.org/dataset/881069>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2022-08-02

## Project

» [Quantifying Processes Driving Interannual Variability in the Biological Carbon Pump in the Western Antarctic Peninsula](#) (WAP Carbon export)

Contributors	Affiliation	Role
<a href="#">Stukel, Michael R.</a>	Florida State University EOAS (FSU - EOAS)	Principal Investigator, Contact
<a href="#">Newman, Sawyer</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Uptake of nitrate by phytoplankton during the 2012-2013 field season of the Palmer LTER program near Anvers Island in the Western Antarctic Peninsula. Nitrate uptake is a proxy for production based on “new” nitrogen entering the ecosystem and hence should balance export flux when integrated over sufficiently large spatial scales and long temporal scales. Nitrate uptake was measured through the uptake of isotopically labeled nitrate, typically at 5 depths spanning the surface to 65 m depth. Measurements were typically made twice per week through the ice-free summer phytoplankton growing season.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Location:** Western Antarctic Peninsula near Anvers Island

**Spatial Extent:** N:-64.7795 E:-64.0405 S:-64.815 W:-64.0725

**Temporal Extent:** 2012-10-31 - 2013-03-21

## Methods & Sampling

We conducted 12-h nitrate uptake experiments in on-land incubators. Each incubator was cooled with flow-through seawater. Nitrate uptake rates were typically conducted on 1-L samples collected from 0, 5, 10, 20, and 65 m. Samples were placed in polycarbonate bottles and spiked with 15-N labeled nitrate, then placed in mesh shading to approximate 100% surface irradiance (0 m), 50% surface irradiance (5 m), 25% surface irradiance (10 m), 10% surface irradiance (20 m), and dark (65 m). These light levels were chosen to match Palmer LTER net primary production depths and light levels and are close to the seasonal average light levels at these depths, although actual light varied substantially throughout the season (see Stukel et al. 2015). At the end of the 24-hour incubation, bottles were immediately vacuum filtered onto pre-combusted 25-mm GF/F filters. Filters were rinsed with filtered seawater, wrapped in foil and stored at 80°C. On land, samples were fumigated with HCl vapor to remove inorganic carbon, dried, and placed inside a tin cup for C/N and isotopic analysis at the UC Davis stable isotope facility. NO<sub>3</sub><sup>-</sup> uptake rates in each incubation bottle (and associated uncertainties) were determined using equations in Dugdale and Wilkerson (1986). For additional details see Stukel et al. (2015).

## BCO-DMO Processing Description

- Spaces removed from column names and replaced with underscores ("\_")
- Row containing units removed from the data file
- Standardized format of dates in the data file
- Converted dates in the data file from %m-%d-%y format to %Y-%m-%d format
- Added a column ISO\_DateTime\_UTC containing date times represented in UTC
- Removed AM and PM from time field in data file
- Added concatenated datetime field to data file

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>881069_v1_palmer_station_nitrate_uptake.csv</b> (Comma Separated Values (.csv), 17.90 KB) MD5:156075fdff14739eba803d4eef40776f
Primary data file for dataset ID 881069, version 1

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Dugdale, R. C., & Wilkerson, F. P. (1986). The use of 15 N to measure nitrogen uptake in eutrophic oceans; experimental considerations<sup>1,2</sup>. *Limnology and Oceanography*, 31(4), 673–689. Portico.

<https://doi.org/10.4319/lo.1986.31.4.0673>

*Methods*

Stukel, M. R., Asher, E., Couto, N., Schofield, O., Strebel, S., Tortell, P., & Ducklow, H. W. (2015). The imbalance of new and export production in the western Antarctic Peninsula, a potentially “leaky” ecosystem. *Global Biogeochemical Cycles*, 29(9), 1400–1420. Portico. <https://doi.org/10.1002/2015gb005211>

<https://doi.org/10.1002/2015GB005211>

*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Station	Palmer LTER station name from which sample was collected.	unitless
Latitude	Station Latitude in decimal degrees; a positive value indicates a Northern coordinate.	decimal degrees
Longitude	Station Longitude in decimal degrees; a positive value indicates an Eastern coordinate.	decimal degrees
Date	Date and time of sample collection in local time at Palmer Station (UTC-3).	unitless
Incubation_Start_Time	Time of incubation start in local time at Palmer Station (UTC-3).	unitless
ISO_DateTime_UTC	Date and time of incubation start in UTC.	unitless
Depth	Depth at which sample was collected.	meters (m)
Incubation_Light_Level	Fraction of surface irradiance.	unitless
Nitrate	Nitrate + Nitrite concentration.	micromoles per Liter (umol/L)
NO3_Uptake	Nitrate uptake.	micromoles per Liter per day (umol/L/day)

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Quantifying Processes Driving Interannual Variability in the Biological Carbon Pump in the Western Antarctic Peninsula (WAP Carbon export)

**Coverage:** Western Antarctic Peninsula (Palmer LTER Study Region)

#### *NSF Award Abstract:*

Algae in the surface ocean convert carbon dioxide into organic carbon through photosynthesis. The biological carbon pump transports this organic carbon from the atmosphere to the deep ocean where it can be stored for tens to hundreds of years. Annually, the amount transported is similar to that humans are currently emitting by burning fossil fuels. However, at present we cannot predict how this important process will change with a warming ocean. These investigators plan to develop a 15+ year time-series of vertical carbon transfer for the Western Antarctic Peninsula; a highly productive Antarctic ecosystem. This region is also rapid transition to warmer temperatures leading to reduced sea ice coverage. This work will help researchers better understand how the carbon cycle in the Western Antarctic Peninsula will respond to climate change. The researchers will develop the first large-scale time-series of carbon flux anywhere in the ocean. This research will also support the education and training of a graduate student and support the integration of concepts in

Antarctic research into two undergraduate courses designed for non-science majors and advanced earth science students. The researchers will also develop educational modules for introducing elementary and middle-school age students to important concepts such as gross and net primary productivity, feedbacks in the marine and atmospheric systems, and the differences between correlation and causation. Results from this proposal will also be incorporated into a children's book, "Plankton do the Strangest Things", that is targeted at 5-7 year olds and is designed to introduce them to the incredible diversity and fascinating adaptations of microscopic marine organisms.

This research seeks to leverage 6 years (2015-2020) of  $^{234}\text{Th}$  samples collected on Palmer LTER program, 5 years of prior measurements (2009-2010, 2012-2014), and upcoming cruises (2021-2023) to develop a time-series of summertime particle flux in the WAP that stretches for 15 years. The  $^{238}\text{U}$ - $^{234}\text{Th}$  disequilibrium approach utilizes changes in the activity of the particle-active radio-isotope  $^{234}\text{Th}$  relative to its parent nuclide  $^{238}\text{U}$  to quantify the flux of sinking carbon out of the surface ocean (over a time-scale of  $\sim$ one month). This proposal will fund  $^{234}\text{Th}$  analyses from nine years' worth of cruises (2015-2023) and extensive analyses designed to investigate the processes driving inter-annual variability in the BCP. These include: 1) physical modeling to quantify the importance of advection and diffusion in the  $^{234}\text{Th}$  budget, 2) time-series analyses of particle flux, and 3) statistical modeling of the relationships between particle flux and multiple presumed drivers (biological, chemical, physical, and climate indices) measured by collaborators in the Palmer LTER program. This multi-faceted approach is critical for linking the measurements to models and for predicting responses to climate change. It will also test the hypothesis that export flux is decreasing in the northern WAP, increasing in the southern WAP, and increasing when integrated over the entire region as a result of earlier sea ice retreat and a larger ice-free zone. The project will also investigate relationships between carbon export and multiple potentially controlling factors including: primary productivity, algal biomass and taxonomic composition, biological oxygen saturation, zooplankton biomass and taxonomic composition, bacterial production, temperature, wintertime sea ice extent, date of sea ice retreat, and climate modes.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1756610</a>
<a href="#">NSF Office of Polar Programs (formerly NSF PLR) (NSF OPP)</a>	<a href="#">OPP-1951090</a>
<a href="#">NSF Office of Polar Programs (formerly NSF PLR) (NSF OPP)</a>	<a href="#">OPP-1340886</a>
<a href="#">NSF Office of Polar Programs (formerly NSF PLR) (NSF OPP)</a>	<a href="#">OPP-1440435</a>

[ [table of contents](#) | [back to top](#) ]