

# Predator size and abundance data from oyster reefs in a northeast Florida estuary collected between April and August 2019 as part of an oyster reciprocal transplant experiment

**Website:** <https://www.bco-dmo.org/dataset/882641>

**Data Type:** Other Field Results, experimental

**Version:** 1

**Version Date:** 2022-12-13

## Project

» [Collaborative research: Quantifying the influence of nonconsumptive predator effects on prey population dynamics](#) (Predatory NCEs and Scale)

Contributors	Affiliation	Role
<a href="#">Kimbrow, David L.</a>	Northeastern University	Principal Investigator
<a href="#">White, J. Wilson</a>	Oregon State University (OSU)	Co-Principal Investigator
<a href="#">Gerlach, Dana Stuart</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Results in this data set come from an experiment conducted at two sites in an estuary in NE Florida, USA. At two sites that encompassed different environmental (salinity, aerial exposure) and biotic (predators) stressors, juvenile oysters were reciprocally transplanted within and between the two locations. At each location, the home and away oyster 'demes' were also randomly assigned between a predator enclosure and control treatment. Just prior to the experiment, the abundance and size of two predators (crab and snail) were quantified at each site.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:29.77002 E:-81.2144 S:29.62923 W:-81.2641

**Temporal Extent:** 2019-04-18 - 2019-08-09

## Methods & Sampling

A reciprocal transplant experiment was conducted at two sites in an estuary in NE Florida, USA that encompassed different environmental (salinity, aerial exposure) and biotic (predators) stressors. Juvenile oysters were reciprocally transplanted within and between the two locations. At each location, the home and away oyster 'demes' were randomly assigned between a predator enclosure and a control treatment.

Predation risk was estimated by surveying three reefs within each site (Butler or Pellicer) for residential predators, which consist of mud crabs (*Panopeus herbstii*) and crown conchs (*Melongena Corona*). In May 2019, the length of reef length was estimated by deploying a transect along the crest of the reef. For each reef, that transect was then partitioned into six intervals. In the center of each interval, a second transect was deployed from the reef crest to the seaward edge of the reef to estimate reef width. At the midpoint of the

reef width within each interval, we deployed a 1 × 1 m quadrat and searched for all mud crabs and crown conchs within each quadrat (n = 6 quadrats per reef). Carapace widths of all mud crabs were measured to estimate crab size, while distance from siphon to pointed end of shell was used to estimate conch size. Because crown conchs are concentrated on the seaward edge of the reef at low tide, we also placed the quadrat at the seaward edge of each of the reef intervals to count and measure size of crown conchs, which resulted in 12 quadrat samples per reef with half from the interior (as described above) and half from the reef edge (designated as conch.low).

## Data Processing Description

### BCO-DMO Processing description:

- Converted dates to format (YYYY-MM-DD)
- Adjusted field/parameter names to comply with BCO-DMO naming conventions
- Added columns for "Latitude" and "Longitude" based on site locations
- Added a conventional header with dataset name, PI names, version date

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Kimbro, D. L., White, J. W., Breef-Pilz, A., Peckham, N., Noble, A., & Chaney, C. (2022). Evidence for local adaptation of oysters to a within-estuary gradient in predation pressure weakens with ontogeny. *Journal of Experimental Marine Biology and Ecology*, 555, 151784. <https://doi.org/10.1016/j.jembe.2022.151784>  
*Results*

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### IsRelatedTo

---

Kimbro, D. L., White, J. (2022) **(DRAFT) Tidal inundation results from oyster reciprocal transplant experiment.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-16 <http://lod.bco-dmo.org/id/dataset/882626> [[view at BCO-DMO](#)]

Kimbro, D. L., White, J. (2022) **(DRAFT) Water salinity and temperature data from oyster reciprocal transplant experiment.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-16 <http://lod.bco-dmo.org/id/dataset/882657> [[view at BCO-DMO](#)]

Kimbro, D. L., White, J. (2022) **Individual oyster results from an oyster reciprocal transplant experiment conducted at two sites in an estuary in NE Florida between August 2019 and May 2020.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-14 <http://lod.bco-dmo.org/id/dataset/880691> [[view at BCO-DMO](#)]

Kimbro, D. L., White, J. (2022) **Survival and growth data from an oyster reciprocal transplant experiment conducted at two sites in an estuary in northeast Florida between August 2019 and May 2020.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-16 <http://lod.bco-dmo.org/id/dataset/882606> [[view at BCO-DMO](#)]

Kimbro, D. L., White, J. (2022) **Water flow data from oyster reciprocal transplant experiment conducted at two sites in an estuary in NE Florida between July 2019 and April 2020.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-16 <http://lod.bco-dmo.org/id/dataset/882674> [[view at BCO-DMO](#)]

Kimbro, D. L., White, J., Breef-Pilz, A. (2022) **Seawater properties at two locations in a northeast Florida estuary measured using HydroCAT CTD between July 2019 and April 2020 as part of an oyster reciprocal transplant experiment.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-12-16 <http://lod.bco-dmo.org/id/dataset/885452> [[view at BCO-DMO](#)]

## Parameters

Parameter	Description	Units
Reef_ID	Unique identifier for three different reefs within each site (NERR zones Butler or Pellicer)	unitless
Date	Date of data collection	unitless
Latitude	Latitude of sampling site	decimal degrees
Longitude	Longitude of sampling site	decimal degrees
NERR_zone	National Estuarine Research Reserve zone or Site (Butler or Pellicer)	unitless
Quad_num	Meter position on length of reef where sample was taken	meter (m)
Organism_type	Type of predator organism: crown conch or mudcrab; Designation of conch.low is for data collected at the reef edge where crown conchs are concentrated at low tide	unitless
Size	Length of organism; carapace width for crabs or maximum length of conch	millimeter (mm)
Sex	gender of organism (determined only for crabs) where m=male, f=female, gf=gravid female	unitless

## Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	calipers
Generic Instrument Description	A caliper (or "pair of calipers") is a device used to measure the distance between two opposite sides of an object. Many types of calipers permit reading out a measurement on a ruled scale, a dial, or a digital display.

## Project Information

### **Collaborative research: Quantifying the influence of nonconsumptive predator effects on prey population dynamics (Predatory NCEs and Scale)**

**Coverage:** Sub-tropical estuarine waters (29.67,-81.21)

#### *NSF Award Abstract:*

Predators can affect populations of their prey in two ways: by consuming them ("consumptive effects" or "CE"s), or by causing the prey to change behavior to avoid contact with the predator. For example, prey often spend less time feeding and more time watching out for predators, which comes with the cost of lower food intake and thus slower growth. Such "non-consumptive effects" (NCEs) have been described for a wide range of terrestrial and marine prey species, from elk to clams, but mostly in short-term (< 1 month) experiments. These prior results suggest that in some cases, the behavioral changes (NCEs) have a bigger effect on prey populations than consumption by predators (CEs). However, those short-term, controlled experiments may artificially inflate the perceived importance of NCEs. Over longer time periods, prey may adapt or become acclimated to predation risk, and NCEs may become less important. Additionally, environmental variability (e.g., differences in the availability of the prey's food between study sites) may have a bigger effect on prey populations than NCEs do. This project will use a combination of short- (months) and long-term (years) field experiments and mathematical models to evaluate the role of NCEs on Florida oyster reefs. The prey species in this study is the eastern oyster, an important marine resource in the southeast US for harvesting and habitat creation; the main oyster predator is a mud crab. In this study, results from mathematical models of oyster populations will be compared to experimental data from the field to see whether including NCEs in the model leads to better model predictions. Better understanding of NCEs in oysters should improve management of that important marine resource. Furthermore, the mathematical model will be used to develop broader, generalizable conclusions about the importance of NCEs that could be applied to other important prey species. This project will provide data useful for oyster resource management, will support public education regarding the ecological importance of NCEs, and will enhance the scientific engagement of underrepresented groups in the study region. The project will support a partnership with the Guana Tolomato Matanzas National Estuarine Research Reserve in Florida, including data sharing, sponsoring an oyster management symposium, and funding the development of multimedia scientific outreach materials at the reserve that will be used by a large and diverse population of K-12 students in the surrounding community. The project will train a postdoctoral researcher, two graduate students, two undergraduate students, and research results will be disseminated by those students and the principal investigators at scientific conferences, in journal publications, and in online content through an ongoing partnership with a Florida public television station.

Predators can alter prey population dynamics by causing fear-based shifts in prey traits (nonconsumptive effect, NCE). The importance of NCEs for prey populations - relative to direct consumption by predators (consumptive effects, CE) - remains uncertain, particularly because short-term studies of NCEs cannot estimate their effect over multiple prey generations. This project addresses that knowledge gap by combining short- and long-term field experiments with population models to investigate the importance of NCEs on oyster population dynamics in a Florida estuary. The central question is whether accounting for NCEs improves the ability to predict long-term trends in oyster population abundance. Several types of NCEs are present in this system: exposure to water-containing predator odors reduces oyster larval recruitment and causes juvenile oysters to increase shell thickness, reducing their somatic growth. In addition to CE and NCEs, environmental gradients in stress, food, and propagule delivery are also present in this system. Those environmental factors can have strong effects on post-settlement survivorship, growth, and recruitment of oysters, so the relative importance of predator CE and NCEs may vary along those spatial gradients as well. This project will consist of four components. (1) A series of short-term field experiments to test how NCEs vary with predator density and environmental variables, and whether one of the NCEs (increased shell thickness) actually reduces vulnerability to predators. (2) A population model, parameterized using experimental results; model simulations will quantify how the relative importance of NCEs should vary over time, space, and environmental gradients. (3) A longer-term (3.5 year) field experiment; the results from this experiment will be compared to model predictions to test whether accounting for NCEs improves predictions of long-term variation in oyster population dynamics. (4) A general form of the model will be developed to broadly investigate the effect of NCEs on non-equilibrium, transient population dynamics. By combining models and field experiments, this project will bridge the gap between the theoretical understanding of how NCEs affect population dynamics and empirical tests of that theory, advancing the field towards the goal of predicting how multiple interacting factors structure communities.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1736943</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1820540</a>

[ [table of contents](#) | [back to top](#) ]