

Tail wave amplitude from 5 high-speed video frames at 3 temperatures from *Oikopleura dioica* particle tracking experiments conducted in December 2015

Website: <https://www.bco-dmo.org/dataset/897825>

Data Type: experimental

Version: 1

Version Date: 2023-06-22

Project

» [Collaborative Research: Comparative feeding by gelatinous grazers on microbial prey](#) (Gelatinous Grazer Feeding)

Contributors	Affiliation	Role
Gemmell, Brad J.	University of South Florida (USF)	Principal Investigator
Sutherland, Kelly Rakow	University of Oregon	Principal Investigator, Contact
Conley, Keats R.	University of Oregon	Scientist
Hiebert, Terra C.	University of Oregon	Scientist
von Dassow, George	University of Oregon	Scientist
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

These data include tail beat kinematics measurements and particle tracking from the appendicularian *Oikopleura dioica* during experiments conducted in December 2015 at the Sars Centre for Marine Molecular Biology in Bergen, Norway. The data were collected from high-speed video frames. The experiments comprised 3 temperature treatments. This dataset includes measurements of tail wave amplitude. Tail beat kinematic measurements were used to describe how *O. dioica* drives flow across food concentrating filters and how it is affected by temperature.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Temporal Extent: 2015-12-05 - 2015-12-10

Methods & Sampling

All experimental animals were obtained from the appendicularian culture facility at the Sars Centre for Marine Molecular Biology in Bergen, Norway in December 2015. *Oikopleura dioica* were filmed individually following

Gemmell et al. (2014). Images were recorded using an Edgertronic high-speed camera (1280 × 1024-pixel resolution, 500 frames per second) with brightfield illumination from a fiber optic light source, or a Photron FastCam Mini Ux100 (1280x1024, 125-1000 frames per second) with darkfield illumination from a tilting mirror base. The filming vessel was positioned on a manually adjustable stage between the light source and the camera. A long working-distance microscope objective (4x or 40x) was mounted to an adjustable-height optics clamp positioned between the filming vessel and the camera. Videos were converted to image stacks in QuickTime Pro. Day 1 animals were filmed in a 50-milliliter glass cuvette in treatments comprising 3 temperatures: 5° Celsius, 15° Celsius, and 25° Celsius.

The videos are attached as Supplemental Files (there is one .zip folder for each temperature treatment).

Data Processing Description

Data Processing:

Tail beat kinematics were measured in ImageJ. Wave amplitude was determined from 5 frames per animal, with each frame taken at the maximum angle of the tail-trunk junction. Amplitude was measured as the maximum height from a line originating at the tail-trunk junction and bisecting the tail at the inflection point. Two amplitude measurements were taken for each frame.

BCO-DMO Processing:

- renamed fields to comply with BCO-DMO naming conventions;
- bundled the videos into 3 .zip files (see Supplemental Files).

[[table of contents](#) | [back to top](#)]

Data Files

File
wave_amplitude.csv (Comma Separated Values (.csv), 13.61 KB) MD5:25539f9efd59106d8c4ea214b732ddd1
Primary data file for dataset ID 897825

[[table of contents](#) | [back to top](#)]

Supplemental Files

File
15C.zip (ZIP Archive (ZIP), 1.94 GB) MD5:5ee628c77dd3b5079923bb2d4004a5dc This .zip contains 17 .mov video files from the 15 degrees Celsius treatments. See the file inventory (Appendicularian_Video_Inventory.csv) for a description of each file.
25C.zip (ZIP Archive (ZIP), 2.01 GB) MD5:02928e5f9fd97b1f5f73cd4dc8f5e6c4 This .zip contains 16 .mov video files from the 25 degrees Celsius treatments. See the file inventory (Appendicularian_Video_Inventory.csv) for a description of each file.
5C.zip (ZIP Archive (ZIP), 2.49 GB) MD5:6888cc317e68626484f97c5fd5a53df6 This .zip contains 20 .mov video files from the 5 degrees Celsius treatments. See the file inventory (Appendicularian_Video_Inventory.csv) for a description of each file.
Appendicularian_Video_Inventory.csv (Comma Separated Values (.csv), 2.77 KB) MD5:3984ad11c2e6a3e4373883742f6988ad Inventory/description of video files contained in the three .zip files - one for each temperature treatment. This inventory was originally provided as an Excel file and was converted to .csv by BCO-DMO.

Related Publications

Gemmell, B. J., Jiang, H., & Buskey, E. J. (2014). A new approach to micro-scale particle image velocimetry (μ PIV) for quantifying flows around free-swimming zooplankton. *Journal of Plankton Research*, 36(5), 1396–1401. <https://doi.org/10.1093/plankt/fbu067>
Methods

Hiebert, TC, Gemmell, BJ, von Dassow, G, Conley, KR, & Sutherland, KR. (Under Review). The hydrodynamics and kinematics of the appendicularian tail underpin peristaltic pumping.
Results

Related Datasets

IsRelatedTo

Gemmell, B. J., Sutherland, K. R., Conley, K. R., Hiebert, T. C., von Dassow, G. (2023) **Filter area measurements from *Oikopleura dioica* tail beat kinematics and particle tracking experiments conducted in December 2015**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2023-06-15 doi:10.26008/1912/bco-dmo.897665.1 [[view at BCO-DMO](#)]

Gemmell, B. J., Sutherland, K. R., Conley, K. R., Hiebert, T. C., von Dassow, G. (2023) **Inlet particle speeds from 5 high-speed video frames at 3 temperature treatments from *Oikopleura dioica* particle tracking experiments conducted in December 2015**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2023-06-15 doi:10.26008/1912/bco-dmo.897682.1 [[view at BCO-DMO](#)]

Gemmell, B. J., Sutherland, K. R., Conley, K. R., Hiebert, T. C., von Dassow, G. (2023) **Tail beat frequency in the appendicularian *Oikopleura dioica* during particle tracking experiments comprised of three temperature treatments conducted in December 2015**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2023-06-15 doi:10.26008/1912/bco-dmo.897617.1 [[view at BCO-DMO](#)]

Parameters

Parameter	Description	Units
Temperature	Temperature treatment	degrees Celsius
Video	Video identifier. The 10-digit video file names correspond to the .mov file names. The .mov files are provided in the attached .zip folders as Supplemental Files.	unitless
Frame	Video frame number	unitless
Time	Time at frame measured	seconds
Wavelength	Sine wavelength	millimeters (mm)
Tail_Length	Total tail length	millimeters (mm)
Amplitude_A	Maximum wave amplitude; two amplitude measurements were made per frame, and were denoted A and B.	millimeters (mm)
Amplitude_A_Norm	Maximum wave amplitude normalized to tail length	millimeters (mm)
Amplitude_B	Maximum wave amplitude; two amplitude measurements were made per frame, and were denoted A and B.	millimeters (mm)
Amplitude_B_norm	Maximum wave amplitude normalized to tail length	millimeters (mm)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Edgertronic high-speed camera
Generic Instrument Name	high-speed camera
Generic Instrument Description	A high-speed imaging camera is capable of recording rapid phenomena with high-frame rates. After recording, the images stored on the medium can be played back in slow motion. The functionality in a high-speed imaging device results from the frame rate, or the number of individual stills recorded in the period of one second (fps). Common video cameras will typically record about 24 to 40 fps, yet even low-end high-speed cameras will record 1,000 fps.

Dataset-specific Instrument Name	Photron FastCam Mini Ux100
Generic Instrument Name	high-speed camera
Generic Instrument Description	A high-speed imaging camera is capable of recording rapid phenomena with high-frame rates. After recording, the images stored on the medium can be played back in slow motion. The functionality in a high-speed imaging device results from the frame rate, or the number of individual stills recorded in the period of one second (fps). Common video cameras will typically record about 24 to 40 fps, yet even low-end high-speed cameras will record 1,000 fps.

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Comparative feeding by gelatinous grazers on microbial prey (Gelatinous Grazer Feeding)

Coverage: North Pacific Subtropical Gyre, at a field site 3 nautical miles offshore of Kona, Hawai'i (19.710746 N, 22.75 W) & Sars Centre for Marine Molecular Biology in Bergen, Norway

NSF Award Abstract:

The oceans are dominated by microscopic plants and animals (microorganisms) that are at the base of the food web and drive energy and carbon cycles on global scales. Soft jellylike animals called gelatinous grazers specialize in feeding on microorganisms using nets made out of mucus. Gelatinous grazers are abundant in the ocean and have high feeding rates on microorganisms so could have a very strong influence on the abundance and diversity of microorganisms and could change how oceanic food webs are currently understood. However, gelatinous grazers are very fragile and patchy in their distributions so it has been difficult to determine the magnitude and dynamics of these important predator-prey relationships on a meaningful scale using traditional approaches, thus they have typically been disregarded in food web studies. Learning more about the predator-prey relationship between gelatinous grazers and microorganisms will improve understanding of the structure, mechanics, and dynamics of the ocean's food web, which is a critical economic and ecosystem resource on Earth. This project is determining grazing rates by gelatinous animals on microbes to inform food web models. The project also trains students to communicate, disseminate, and interpret scientific findings. These broader impacts goals will be attained through partnerships at the University of Oregon (Applied Scientific Communication) and Portland State University (Advanced Technical Writing), training of 1 PhD student, 2 undergraduates, and 4 science communication interns, and development of a week-long workshop and establish student mentorship relationships towards production of communication products.

The project integrates laboratory and oceanographic approaches to address several specific aspects of the predator-prey relationship between gelatinous grazers and ocean microorganisms. Five distinct types of gelatinous grazers, each with different feeding morphologies and life history, will be studied in an oceanographic setting with an abundant and diverse natural microbial population. These target organisms include pelagic tunicates (salps, appendicularians, doliolods and pyrosomes) and thecosome pteropods. The approach quantifies: 1) grazing rates in the natural ocean environment, 2) particle selectivity with a focus on size and shape and, 3) the morphological and hydrodynamic properties of feeding that underlie the measured grazing rates and particle selection. The project uses a variety of techniques including sampling via SCUBA diving, laboratory experiments, high speed/high resolution videography, flow cytometry, and DNA sequencing techniques.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1851537

[[table of contents](#) | [back to top](#)]