

Total dissolvable iron concentrations in rainwater from weekly sampling at Tudor Hill Bermuda between November 2018 and March 2020.

Website: <https://www.bco-dmo.org/dataset/906757>

Data Type: Other Field Results

Version: 1

Version Date: 2023-08-25

Project

- » [Operation of a Community Marine-Atmospheric Sampling Facility at Tudor Hill, Bermuda](#) (THMAO)
- » [NSFGEO-NERC: Collaborative Research: Using Time-series Field Observations to Constrain an Ocean Iron Model](#) (BAIT)

Programs

- » [United States Surface Ocean Lower Atmosphere Study](#) (U.S. SOLAS)
- » [U.S. GEOTRACES](#) (U.S. GEOTRACES)

Contributors	Affiliation	Role
Sedwick, Peter N.	Old Dominion University (ODU)	Principal Investigator
Sohst, Bettina	Old Dominion University (ODU)	Technician
Gerlach, Dana Stuart	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

These data include total-dissolvable iron concentrations in rainwater and corresponding rainfall amounts for composite samples collected during approximately weekly intervals on the sampling tower at Tudor Hill, Bermuda, between November 2018 and March 2020. The data allow estimates of the wet deposition of iron to the Bermuda region over the period of the BAIT project, which included cruises in the Bermuda Atlantic Time-series Study (BATS) region in March, May, August and November 2019. The rain samples were collected by staff of the Bermuda Institute of Ocean Sciences (the Tudor Hill tower facility is managed by Dr. Andrew Peters, with funding from NSF), and sample processing and analyses were completed in the laboratory of Dr. Peter Sedwick at Old Dominion University.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Coverage

Spatial Extent: Lat:32.265 Lon:-64.879

Temporal Extent: 2018-11-26 - 2020-03-16

Dataset Description

Weekly sampling was conducted at Tudor Hill, Bermuda from November 2018 to March 2020. The sampling

tower platform is approximately 30 meters above sea level with coordinates of 32.265°N, 64.879°W.

Methods & Sampling

Composite samples of bulk rainwater were collected atop the 23 meter height sampling tower at Tudor Hill, Bermuda, on an approximately weekly basis from November 2018 through March 2020, bracketing the four BAIT project cruises. Rainwater samples were collected in acid-cleaned 2-L wide-mouth fluorinated high-density polyethylene bottles (Nalgene) using an automatic rain sampler (N-Con Systems ADS 00-120); Rain samples were subsequently acidified to 0.4% (v/v) with 6 M ultrapure hydrochloric acid (Fisher Optima) in the collection bottles, and then after two months the acidified, unfiltered samples were transferred into acid-cleaned 125 mL low-density polyethylene bottles (Nalgene) for analysis of "total-dissolvable Fe" (TDFe; Sedwick et al., 2007). Field blanks for the rainwater (125 mL ultrapure deionized water) were deployed on the Tudor Hill tower and processed in the same manner as samples, but without opening the rain sampler.

Iron concentrations in acidified rainwater were determined by inductively-coupled plasma mass spectrometry (ICP-MS, Thermo Fisher Scientific ElementXR) without preconcentration, using calibration standards prepared in 0.4% ultrapure hydrochloric acid (Fisher Optima) and yttrium as an internal standard. The field blank value for rainwater was 0.17 nmol TDFe, which equates to a concentration of 1.7 nmol L^{-1} for a typical rain sample volume of 100 mL. The limit of detection for iron in blank-corrected rain samples was estimated as 0.002 nmol per sample, from three times the standard deviation on the mean of replicate analyses of the single field blank (in the absence of a replicate field blank for rain). Repeat ICP-MS analyses of iron in the rainwater samples indicate an analytical uncertainty of less than $\pm 5\%$ (one-sigma). In the absence of duplicate samples for rainwater, we assume an overall analytical uncertainty of less than $\pm 25\%$ (one-sigma) for TDFe in rain, similar to that estimated from our analyses of total iron in replicate aerosol samples.

Problems/Issues:

For some sampling periods there were problems with the rain sampler closing, or with spillage or loss of the sample bottle, or issues with the rain gauge; in these cases data are labelled with a QC flag of 2. No rain samples were collected during the period of 16 September-October 14, 2019, owing to the passage of Hurricane Humberto and subsequent loss of electrical power at the Tudor Hill sampling site.

Data Processing Description

Inductively-coupled plasma mass spectrometer: Instrumental data were collected using ElementXR processing software (Thermo Fisher Scientific), and post-analysis calculations were performed using Microsoft Excel.

BCO-DMO Processing Description

- Imported data from source file "BAIT_Rain_Fe_Data.xlsx" into the BCO-DMO data system
- Added columns for latitude and longitude of sampling tower
- Modified parameter (column) names to conform with BCO-DMO naming conventions
- Replaced spaces with underscores in parameter (column) names

[[table of contents](#) | [back to top](#)]

Data Files

File
906757_v1_bait_rain_fe.csv (Comma Separated Values (.csv), 3.16 KB) MD5:178580ca3b57285066710e21490368b7
Primary data file for dataset ID 906757 version 1

[[table of contents](#) | [back to top](#)]

Related Publications

Sedwick, P. N., Sholkovitz, E. R., & Church, T. M. (2007). Impact of anthropogenic combustion emissions on the fractional solubility of aerosol iron: Evidence from the Sargasso Sea. *Geochemistry, Geophysics, Geosystems*, 8(10), n/a-n/a. <https://doi.org/10.1029/2007gc001586> <https://doi.org/10.1029/2007GC001586>
Methods

Sedwick, P. N., Sohst, B. M., Buck, K. N., Caprara, S., Johnson, R. J., Ohnemus, D. C., Sofen, L. E., Tagliabue, A., Twining, B. S., & Williams, T. E. (2023). Atmospheric Input and Seasonal Inventory of Dissolved Iron in the Sargasso Sea: Implications for Iron Dynamics in Surface Waters of the Subtropical Ocean. *Geophysical Research Letters*, 50(6). Portico. <https://doi.org/10.1029/2022GL102594>
Results

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Latitude	Latitude of sampling tower at Tudor Hill, Bermuda	decimal degrees
Longitude	Longitude of sampling tower at Tudor Hill, Bermuda	decimal degrees
Sample_ID	Unique identifier for each rainwater sample	unitless
Start_date	Start date for rainwater sampling (local)	unitless
End_date	End date for rainwater sampling (local)	unitless
Sampling_period	Number of days over which rainwater was collected	day
TDFe	Concentration of total dissolvable iron	nanomoles per liter (nmol L ⁻¹)
Rainfall	Rainfall during sampling period (from rain gauge)	millimeters (mm)
Sample_flag	Data quality flag that applies to TDFe and/or Rainfall (1=good, 2=rain sampler or rain gauge malfunction, 3=sample volume insufficient for analysis)	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Thermo Fisher Scientific ElementXR ICP-MS
Generic Instrument Name	Inductively Coupled Plasma Mass Spectrometer
Dataset-specific Description	Iron concentrations in acidified rainwater were determined by inductively-coupled plasma mass spectrometry (ICP-MS, Thermo Fisher Scientific ElementXR)
Generic Instrument Description	An ICP Mass Spec is an instrument that passes nebulized samples into an inductively-coupled gas plasma (8-10000 K) where they are atomized and ionized. Ions of specific mass-to-charge ratios are quantified in a quadrupole mass spectrometer.

Dataset-specific Instrument Name	Rain gauge
Generic Instrument Name	Precipitation Gauge
Dataset-specific Description	Rainfall during sampling period from rain gauge measured in millimeters
Generic Instrument Description	measures rain or snow precipitation

Dataset-specific Instrument Name	N-Con Systems ADS 00-120 automatic rain sampler
Generic Instrument Name	Precipitation Sampler
Dataset-specific Description	Rainwater samples were collected in acid-cleaned bottles using an automatic rain sampler (N-Con Systems ADS 00-120).
Generic Instrument Description	A device that collects a sample of precipitation (rain, hail or snow) as it falls.

[[table of contents](#) | [back to top](#)]

Project Information

Operation of a Community Marine-Atmospheric Sampling Facility at Tudor Hill, Bermuda (THMAO)

Website: <https://tudorhill.bios.asu.edu/>

Coverage: West Bermuda, 32.2646N 64.8791W

NSF Award OCE-2123053 Abstract:

This project supports the Tudor Hill Marine Atmospheric Observatory in Bermuda. Because of its location in the western North Atlantic Ocean, Bermuda has been an important location for many studies of the marine atmosphere. The observatory is well equipped for carrying out research, with field laboratories and a 23 meter tower at Tudor Hill. This allows scientists to study the chemistry and physics of the atmosphere over the oceans. This part of the atmosphere plays an important role in the transfer of moisture, chemicals, and energy between the ocean and the atmosphere. Routine work at the site includes the observation and recording of weather conditions, and sampling of rain and air. In addition, samples and data are collected for a number of research partners, including NOAA, NASA and many US and international universities. As well as supporting research, this project will provide a range of educational opportunities, including through the NSF-funded Research Experience for Undergraduates (REU) program, and through other internships and courses at the Bermuda Institute of Ocean Sciences.

The continued operation and maintenance of the Tudor Hill Marine Atmospheric Observatory facilitates research concerning the chemistry and physics of the marine troposphere and ocean-atmosphere exchange processes. The specific objectives of the project are to: 1) Operate and maintain the atmospheric sampling facility at Tudor Hill, Bermuda; 2) Continue collection of continuous meteorological data and weekly bulk-aerosol and rain samples, which will be archived at BIOS and made freely available to other researchers; and 3) Collect additional atmospheric samples and data for external investigators, and provide for the use of the facility by external investigators. Ongoing partner activities at the facility include monitoring and data acquisition for NASA (AERONET) and NOAA programs (Ozone and greenhouse gas monitoring), and a new sampling location will be established for the US National Atmospheric Deposition Program (NADP) in 2021. The THMAO site will enable research to be undertaken that is central to international initiatives such as IGAC, SOLAS and GEOTRACES. In a regional context, the Tudor Hill facility will complement ongoing oceanographic time-series research in the Sargasso Sea, including Hydrostation S and BATS.

NSF Award OCE-1829686 Abstract:

This project provides continuing support for the Tudor Hill Marine Atmospheric Observatory in Bermuda. The marine boundary layer, the lower level of the atmosphere over the oceans, plays important roles in the global cycles of many chemical elements and this observatory provides a unique facility for the ocean and atmospheric chemistry research communities. At present, many aspects of atmospheric composition, atmospheric transport, and atmosphere-ocean exchange remain poorly understood. This limits our ability to predict and mitigate the effects of anticipated future environmental change. Central to improving our understanding of these topics is the ability to observe and sample the marine boundary layer. In this regard, oceanic island observatories have played a crucial role. By virtue of its location in the western North Atlantic Ocean, Bermuda has been a key location for numerous field studies of the marine troposphere and is well equipped for such research, with the 23 meter high atmospheric sampling tower and site laboratories at Tudor Hill. Routine facility operations include the collection of continuous meteorological data and weekly bulk aerosol and rainwater samples, which are archived at the Bermuda Institute of Ocean Sciences and made freely available to the scientific community. Additionally, samples and data are collected for a variety of external research programs funded by NSF, NOAA, NASA and other agencies. This project is expected to make significant educational contributions at a number of levels, including undergraduate education opportunities through the NSF-funded Research Experience for Undergraduates (REU) program, and through other courses taught at the Bermuda Institute of Ocean Sciences.

The continued operation and maintenance of THMAO site will facilitate research topics that are central to international research initiatives such as IGAC, SOLAS and GEOTRACES. The specific objectives of the project are to: 1) Operate and maintain the atmospheric sampling facility at Tudor Hill, Bermuda; 2) Continue collection of continuous meteorological data and weekly bulk-aerosol and rain samples, which will be archived at BIOS and made freely available to other researchers; and 3) Collect additional atmospheric samples and data for external investigators, and provide for the use of the facility by external investigators. The continued operation and maintenance of the Tudor Hill site facilitates research concerning the chemistry and physics of the marine troposphere and ocean-atmosphere exchange processes. In a regional context, the Tudor Hill facility will complement ongoing time-series research in the Sargasso Sea, including Hydrostation S and BATS.

NSF Award OCE-1735504 Abstract:

This award would provide a year of continuation funding for the Tudor Hill Marine Atmospheric Observatory in Bermuda. The lower level of the atmosphere over the oceans plays important roles in the global cycles of many chemical elements and this observatory provides a unique facility for the ocean and atmospheric chemistry research communities. At present, many aspects of atmospheric composition, atmospheric transport, and atmosphere-ocean exchange remain poorly understood. This limits our ability to predict and mitigate the effects of anticipated future environmental change. Central to improving our understanding of these topics is the ability to observe and sample the marine boundary layer. In this regard, oceanic island observatories have played a crucial role. By virtue of its location in the western North Atlantic Ocean, Bermuda has been a key location for numerous field studies of the marine troposphere and is well equipped for such research, with the 23-m high atmospheric sampling tower and site laboratories at Tudor Hill. Routine facility operations include the collection of continuous meteorological data and weekly bulk aerosol and rainwater samples, which are archived at the Bermuda Institute of Ocean Sciences and made freely available to the scientific community. Additionally, samples and data are collected for a variety of external research programs funded by NSF, NOAA, NASA and other agencies.

The specific objectives of the project are to: 1) Operate and maintain the atmospheric sampling facility at Tudor Hill, Bermuda; 2) Continue collection of continuous meteorological data and weekly bulk-aerosol and rain samples, which will be archived at BIOS and made freely available to other researchers; and 3) Collect additional atmospheric samples and data for external investigators, and provide for the use of the facility by external

investigators. The continued operation and maintenance of the Tudor Hill site facilitates research concerning the chemistry and physics of the marine troposphere and ocean-atmosphere exchange processes. In a regional context, the Tudor Hill facility will complement ongoing time-series research in the Sargasso Sea, including Hydrostation S and BATS. This project is expected to make significant educational contributions at a number of levels, including undergraduate education opportunities through the NSF-funded Research Experience for Undergraduates (REU) program, and through other courses taught at BIOS.

NSFGEO-NERC: Collaborative Research: Using Time-series Field Observations to Constrain an Ocean Iron Model (BAIT)

Coverage: Bermuda Atlantic Time-Series Study region, waters of the western Subtropical North Atlantic Gyre (ca. 30°N-33°N, 62°W-65°W)

NSF and NERC Award Abstract:

Iron is an essential nutrient for the growth of phytoplankton in the oceans. As such, iron plays key roles in regulating marine primary production and the cycling of carbon. It is thus important that models of ocean biology and chemistry consider iron, in order to explore past, present and future variations in marine productivity and the role of the ocean in the global carbon cycle. In this joint project involving researchers in the U.S. and the U.K., supported by both NSF and the Natural Environment Research Council (U.K.), field data from the Bermuda Atlantic Time-series Study (BATS) region will be combined with an established, state-of-the-art ocean biogeochemical model. By leveraging the known seasonal-scale physical, chemical and biological changes in the BATS region, the oceanographic context provided by the BATS core data, and an existing model of the regional physical circulation, the proposed study will yield process-related information that is of general applicability to the open ocean. In particular, the proposed research will focus on understanding the atmospheric input, biological uptake, regeneration and scavenging removal of dissolved iron in the oceanic water column, which have emerged as major uncertainties in the ocean iron cycle. The project will include significant educational and training contributions at the K-12, undergraduate, graduate and postdoctoral levels, as well as public outreach efforts that aim to explain the research and its importance.

The ability of ocean models to simulate iron remains crude, owing to an insufficient understanding of the mechanisms that drive variability in dissolved iron, particularly the involvement of iron-binding ligands, colloids and particles in the surface input, biological uptake, regeneration and scavenging of dissolved iron in the upper ocean. Basin-scale data produced by the GEOTRACES program provide an important resource for testing and improving models and, by extension, our mechanistic understanding of the ocean iron cycle. However such data provide only quasi-synoptic 'snapshots', which limits their utility in isolating and identifying the processes that control dissolved iron in the upper ocean. The proposed research aims to provide mechanistic insight into these governing processes by combining time-series data from the BATS region with numerical modeling experiments. Specifically, seasonally resolved data on the vertical (upper 2,000 meters) and lateral (tens of kilometers) distributions of particulate, dissolved, colloidal, soluble and ligand-bound iron species will be obtained from the chemical analysis of water column samples collected during five cruises, spanning a full annual cycle, shared with the monthly BATS program cruises. These data, along with ancillary data from the BATS program, will be used to test and inform numerical modeling experiments, and thus derive an improved understanding of the mechanisms that control the distribution and dynamics of dissolved iron in the oceanic water column.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

This is a project jointly funded by the National Science Foundation's Directorate for Geosciences (NSF/GEO) and the National Environment Research Council (NERC) of the United Kingdom (UK).

[[table of contents](#) | [back to top](#)]

Program Information

United States Surface Ocean Lower Atmosphere Study (U.S. SOLAS)

Website: <http://www.us-solas.org/>

Coverage: Global

The Surface Ocean Lower Atmosphere Study (SOLAS) program is designed to enable researchers from different disciplines to interact and investigate the multitude of processes and interactions between the coupled ocean and atmosphere.

Oceanographers and atmospheric scientists are working together to improve understanding of the fate, transport, and feedbacks of climate relevant compounds, and also weather and hazards that are affected by processes at the surface ocean.

Oceanographers and atmospheric scientists are working together to improve understanding of the fate, transport, and feedbacks of climate relevant compounds.

Physical, chemical, and biological research near the ocean-atmosphere interface must be performed in synergy to extend our current knowledge to adequately understand and forecast changes on short and long time frames and over local and global spatial scales.

The findings obtained from SOLAS are used to improve knowledge at process scale that will lead to better quantification of fluxes of climate relevant compounds such as CO₂, sulfur and nitrogen compounds, hydrocarbons and halocarbons, as well as dust, energy and momentum. This activity facilitates a fundamental understanding to assist the societal needs for climate change, environmental health, weather prediction, and national security.

The US SOLAS program is a component of the International SOLAS program where collaborations are forged with investigators around the world to examine SOLAS issues ubiquitous to the world's oceans and atmosphere.

[Â» International SOLAS Web site](#)

Science Implementation Strategy Reports

[US-SOLAS](#) (4 MB PDF file)

[Other SOLAS reports](#) are available for download from the US SOLAS Web site

U.S. GEOTRACES (U.S. GEOTRACES)

Website: <http://www.geotraces.org/>

Coverage: Global

GEOTRACES is a [SCOR](#) sponsored program; and funding for program infrastructure development is provided by the [U.S. National Science Foundation](#).

GEOTRACES gained momentum following a special symposium, S02: Biogeochemical cycling of trace elements and isotopes in the ocean and applications to constrain contemporary marine processes (GEOSECS II), at a 2003 Goldschmidt meeting convened in Japan. The GEOSECS II acronym referred to the Geochemical Ocean Section Studies To determine full water column distributions of selected trace elements and isotopes, including their concentration, chemical speciation, and physical form, along a sufficient number of sections in each ocean basin to establish the principal relationships between these distributions and with more traditional hydrographic parameters;

- * To evaluate the sources, sinks, and internal cycling of these species and thereby characterize more completely the physical, chemical and biological processes regulating their distributions, and the sensitivity of these processes to global change; and
- * To understand the processes that control the concentrations of geochemical species used for proxies of the past environment, both in the water column and in the substrates that reflect the water column.

GEOTRACES will be global in scope, consisting of ocean sections complemented by regional process studies. Sections and process studies will combine fieldwork, laboratory experiments and modelling. Beyond realizing the scientific objectives identified above, a natural outcome of this work will be to build a community of marine scientists who understand the processes regulating trace element cycles sufficiently well to exploit this knowledge reliably in future interdisciplinary studies.

Expand "Projects" below for information about and data resulting from individual US GEOTRACES research projects.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1829833
NSF Division of Ocean Sciences (NSF OCE)	OCE-1829686

[[table of contents](#) | [back to top](#)]