

# Porewater geochemistry from sediment pushcores collected by ROV Doc Ricketts from Southern California Seeps in May 2021

**Website:** <https://www.bco-dmo.org/dataset/908217>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2025-08-27

## Project

» [Collaborative Research: Redefining the footprint of deep ocean methane seepage for benthic ecosystems](#)  
(Methanosphere)

| Contributors                        | Affiliation   | Role                      |
|-------------------------------------|---|---------------------------|
| <a href="#">Goffredi, Shana</a>     | Occidental College                                  | Co-Principal Investigator |
| <a href="#">Orphan, Victoria J.</a> | California Institute of Technology (Caltech)        | Co-Principal Investigator |
| <a href="#">Magyar, John S.</a>     | California Institute of Technology (Caltech)        | Scientist                 |
| <a href="#">Rauch, Shannon</a>      | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager      |

## Abstract

This dataset provides geochemical data (major ions) for porewaters from sediment pushcores collected by a remotely-operated vehicle (ROV) deployed from the R/V Western Flyer in May 2021 (WF05-21) at the Del Mar and Santa Monica Seeps and in Newport Canyon, California, USA. Cores were sectioned into 1-3 centimeter (cm) horizons, and porewater was extracted shipboard using an argon-pressurized squeezer. Porewater samples were frozen at sea and thawed just before analysis. This dataset reports concentrations of fluoride, acetate, formate, chloride, bromide, nitrate, sulfate, thiosulfate, and phosphate anions and lithium, sodium, potassium, magnesium, and calcium cations. Measurements were made using the Thermo Dual Dionex Integration HPIC ion chromatography system with either a 250-millimeter (mm) Dionex IonPac AS19-4um column (anions) or a 250 mm Dionex IonPac CS16-4um column (cations), in the Resnick Water and Environment Laboratory at the California Institute of Technology (Caltech).

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
  - [BCO-DMO Processing Description](#)
  - [Problem Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:36.776723 E:-117.78227 S:32.904203 W:-122.085014

**Temporal Extent:** 2021-05-19 - 2021-05-24

## Methods & Sampling

Sediment pushcores were collected by a remotely-operated vehicle (ROV Doc Ricketts) from the R/V Western Flyer in May 2021 (cruise ID WF05-21) at the Del Mar Seep (32.9043, -117.7824), Santa Monica Seep Mound 800 (33.7993, -118.6465), Santa Monica Seep Mound 863 (33.7891, -118.6683), and Newport Canyon (33.5450, -117.9251). Pushcores were manually sectioned into 1- or 3-centimeter (cm) horizons using plastic

rings, and porewaters were extracted using a KC Denmark squeezer. Samples for ion chromatography were frozen shipboard and thawed immediately prior to analysis. Measurements were made using a Thermo Dual Dionex Integrion HPIC ion chromatography system with either a 250-millimeter (mm) Dionex IonPac AS19-4um column (anions) with 50 mm guard column or a 250 mm Dionex IonPac CS16-4um column (cations) with a 50 mm guard column, in the Resnick Water and Environment Laboratory at the California Institute of Technology (Caltech).

## Data Processing Description

100 microliter (uL) samples were diluted 50x in milliQ nanopure water before running. Standard curves were generated using calibration standards diluted similarly to samples, with 100 uL of 500 millimolar (mM) NaCl added to mimic typical seawater sample peak behavior. Analyte peaks were integrated and values were calculated using the standards automatically in the Chromeleon software, with manual QC to ensure regular peak shapes.

### Known issues or problems:

Na and Cl are not quantitative, as the standards have an excess of each spiked in, above the upper limit of quantification. Formate values reported as 'ND' indicate they are below the LoQ but not LoD. Ca elutes near the end of the run, and occasionally is not captured in its entirety.

## BCO-DMO Processing Description

- Converted original file "20230630\_WF05-21\_ions\_for\_NSF\_corrected080825.xlsx" to CSV format.
- Imported file "20230630\_WF05-21\_ions\_for\_NSF\_corrected080825.csv" into the BCO-DMO processing system.
- Flagged 'ND', 'NA', and 'NOT\_MEASURED' as missing data identifiers but retained in the dataset.
- Converted date to YYYY-MM-DD format.
- Saved the final file as "908217\_v1\_porewater\_biogeochem.csv".

## Problem Description

Na and Cl are not quantitative, as the standards have an excess of each spiked in, above the upper limit of quantification. Formate values reported as ND indicate they are below the LoQ but not LoD. Ca elutes near the end of the run, and occasionally is not captured in its entirety. Other values noted as ND were below the quantifiable detection limit, or not detected. The limits of detection for each ion are lithium 22 uM, ammonium 194 uM, potassium 566 uM, magnesium 1.6 mM, calcium 198 uM, formate 15 uM, acetate 21 uM, fluoride 23 uM, bromide 40 uM, nitrate 114 uM, sulfate 561 uM, thiosulfate 25 uM, phosphate 47 uM.

[ [table of contents](#) | [back to top](#) ]

## Data Files

| File  |
|---|
| <b>908217_v1_porewater_biogeochem.csv</b> (Comma Separated Values (.csv), 45.63 KB)<br>MD5:c145cf9fe6947f8746704cc79072d636 |
| Primary data file for dataset ID 908217, version 1.   |

[ [table of contents](#) | [back to top](#) ]

## Parameters

| Parameter | Description | Units |
|-----------|-------------|-------|
|           |             |       |

|             |   |                 |
|-------------|---|-----------------|
| serial      | Internal Orphan Lab identifier (sample serial number)   | unitless        |
| Fluoride    | concentration of fluoride ion   | millimolar (mM) |
| Acetate     | concentration of acetate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                  | millimolar (mM) |
| Formate     | concentration of formate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                  | millimolar (mM) |
| Chloride    | concentration of chloride ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                 | millimolar (mM) |
| Bromide     | concentration of bromide ion  | millimolar (mM) |
| Nitrate     | concentration of nitrate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                  | millimolar (mM) |
| Sulfate     | concentration of sulfate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                  | millimolar (mM) |
| Thiosulfate | concentration of thiosulfate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                              | millimolar (mM) |
| Phosphate   | concentration of phosphate ion. ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.                                | millimolar (mM) |
| Lithium     | concentration of lithium ion (Li <sup>+</sup> ). ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.               | millimolar (mM) |
| Sodium      | concentration of sodium ion (Na <sup>+</sup> )  | millimolar (mM) |
| Ammonium    | concentration of ammonium ion (NH <sub>4</sub> <sup>+</sup> ). ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample. | millimolar (mM) |
| Potassium   | concentration of potassium ion (K <sup>+</sup> )  | millimolar (mM) |
| Magnesium   | concentration of magnesium ion (Mg <sup>2+</sup> )  | millimolar (mM) |
| Calcium     | concentration of calcium ion (Ca <sup>2+</sup> ). ND = below level of quantification;<br>NOT_MEASURED = this ion was not measured for this sample.              | millimolar (mM) |

|                  |  |                  |
|------------------|--|------------------|
| pH               | pH value, measured shipboard. ND = below level of quantification; NOT_MEASURED = was not measured for this sample. | unitless         |
| Dive             | Dive number of the ROV Doc Ricketts  | unitless         |
| Lat              | Latitude   | decimal degrees  |
| Long             | Longitude, with west longitude negative  | decimal degrees  |
| Horizon          | depth (cm) in sediment, measured from top of core  | centimeters (cm) |
| Date.UTC         | core collection date (UTC)   | unitless         |
| Time.UTC         | core collection time (UTC)   | unitless         |
| Depth_m          | ocean depth in meters at the sampling location   | meters (m)       |
| Station          | name of the sampling site  | unitless         |
| Sample_Name      | the sample name, which is composed of the dive number and the ROV pushcore number                                  | unitless         |
| ISO_DateTime.UTC | core collection date and time (UTC) in ISO 8601 format   | unitless         |

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

|   |  |
|---|--|
| <b>Dataset-specific Instrument Name</b> | Thermo Dual Dionex Integrion HPIC ion chromatography system  |
| <b>Generic Instrument Name</b>          | Ion Chromatograph  |
| <b>Dataset-specific Description</b>     | Thermo Dual Dionex Integrion HPIC ion chromatography system with either a 250 mm Dionex IonPac AS19-4um column (anions) with 50 mm guard column or a 250 mm Dionex IonPac CS16-4um column (cations) with a 50 mm guard column, in the Resnick Water and Environment Laboratory at the California Institute of Technology (Caltech).  |
| <b>Generic Instrument Description</b>   | Ion chromatography is a form of liquid chromatography that measures concentrations of ionic species by separating them based on their interaction with a resin. Ionic species separate differently depending on species type and size. Ion chromatographs are able to measure concentrations of major anions, such as fluoride, chloride, nitrate, nitrite, and sulfate, as well as major cations such as lithium, sodium, ammonium, potassium, calcium, and magnesium in the parts-per-billion (ppb) range. (from <a href="http://serc.carleton.edu/microbelife/research_methods/biogeochemical/ic....">http://serc.carleton.edu/microbelife/research_methods/biogeochemical/ic....</a> ) |

|   |   |
|---|---|
| <b>Dataset-specific Instrument Name</b> | pushcores   |
| <b>Generic Instrument Name</b>          | Push Corer  |
| <b>Dataset-specific Description</b>     | Sediment pushcores were collected by ROV.   |
| <b>Generic Instrument Description</b>   | Capable of being performed in numerous environments, push coring is just as it sounds. Push coring is simply pushing the core barrel (often an aluminum or polycarbonate tube) into the sediment by hand. A push core is useful in that it causes very little disturbance to the more delicate upper layers of a sub-aqueous sediment. Description obtained from: <a href="http://web.whoi.edu/coastal-group/about/how-we-work/field-methods/coring/">http://web.whoi.edu/coastal-group/about/how-we-work/field-methods/coring/</a> |

|   |   |
|---|---|
| <b>Dataset-specific Instrument Name</b> |   |
| <b>Generic Instrument Name</b>          | ROV Doc Ricketts  |
| <b>Dataset-specific Description</b>     | ROV Doc Ricketts is operated by the Monterey Bay Aquarium Research Institute (MBARI). ROV Doc Ricketts is capable of diving to 4000 meters (about 2.5 miles). The R/V Western Flyer was the support vessel for Doc Ricketts and was designed with a center well whose floor can be opened to allow Doc Ricketts to be launched from within the ship into the water below. For a complete description see: <a href="https://www.mbari.org/technology/rov-doc-ricketts/">https://www.mbari.org/technology/rov-doc-ricketts/</a>   |
| <b>Generic Instrument Description</b>   | The remotely operated vehicle (ROV) Doc Ricketts is operated by the Monterey Bay Aquarium Research Institute (MBARI). ROV Doc Ricketts is capable of diving to 4000 meters (about 2.5 miles). The R/V Western Flyer is the support vessel for Doc Ricketts and was designed with a center well whose floor can be opened to allow Doc Ricketts to be launched from within the ship into the water below. For a complete description, see: <a href="https://www.mbari.org/at-sea/vehicles/remotely-operated-vehicles/rov-doc...">https://www.mbari.org/at-sea/vehicles/remotely-operated-vehicles/rov-doc...</a> |

## Deployments

### WF05-21

|                   |   |
|-------------------|---|
| <b>Website</b>    | <a href="https://www.bco-dmo.org/deployment/877837">https://www.bco-dmo.org/deployment/877837</a> |
| <b>Platform</b>   | R/V Western Flyer   |
| <b>Start Date</b> | 2021-05-19  |
| <b>End Date</b>   | 2021-05-22  |

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### **Collaborative Research: Redefining the footprint of deep ocean methane seepage for benthic ecosystems (Methanosphere)**

**Coverage:** Gulf of Alaska and Southern California Bight

#### *NSF Award Abstract:*

This research examines the role of deep-sea organisms in determining the fate and footprint of methane, a potent greenhouse gas, on Pacific continental margins. The investigators are evaluating the deep ocean methanosphere defined by the microbial communities that consume methane and the animals that directly feed on or form symbioses with methane-consuming microbes. They are also investigating animal communities that gain energy indirectly from methane, as well as those that take advantage of carbonate rocks, the physical manifestation of methane consumption in seafloor sediments. The study of methane seeps in the deep waters of both Alaska (4400-5500 meters) and Southern California (450-1040 meters) is enabling comparisons of the methanosphere under different food-limitation and oxygen regimes. By applying diverse chemical, isotopic, microscopy, and genetic-based analyses to seep microbes and fauna, this study is advancing understanding of the contribution of methane to deep-sea biodiversity and ecosystem function, information that can inform management and conservation actions in US waters. In addition to training for graduate and undergraduate students at their home institutions, the investigators are collaborating with the Alaska Native Science and Engineering Program (ANSEP). They are recruiting Alaskan undergraduates to participate in the research, contributing to ANSEP's online resources that promote interaction between scientists and middle and high school students, and participating in ANSEP's annual residential Career Exploration in Marine Science programs to engage middle school students in learning about deep-sea ecosystems and the variety of career pathways available in marine related fields.

Microbial production and consumption of methane is dynamic and widespread along continental margins, and some animals within deep-sea methane seeps rely on the oxidation and sequestration of methane for nutrition. At the same time, understanding of methane-dependent processes and symbioses in the deep-sea environment is still rudimentary. The goals of this study are to 1) examine the diversity of animals involved in methane-based symbioses and heterotrophic consumption of methane-oxidizing microbes and how these symbioses extend the periphery of seeps, contributing to non-seep, continental slope food webs; and 2) determine whether carbonates on the seep periphery sustain active methanotrophic microbial assemblages, providing a localized food source or chemical fuel for thiotrophic symbioses, via anaerobic oxidation of methane, or free-living, sulfide-oxidizing bacteria consumed by animals. The investigators are addressing these goals by surveying, sampling, and characterizing microbes, water, sediments, carbonates and animals at a deep seep site on the Aleutian Margin and a shallow site off Southern California. Shipboard experiments and laboratory analyses are using molecular, isotopic, geochemical, and radiotracer tools to understand transfer of methane-sourced carbon from aerobic methanotrophs under multiple oxygen levels, pressures, and photosynthetic food inputs. This approach offers a wide lens by which to examine the methane seep footprint, allow reinterpretation of past observations, and identify new scientific areas for future study. Improved characterization of the deep continental margin methanosphere informs climate science, biodiversity conservation, and resource management.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

| Funding Source   | Award                       |
|--|-----------------------------|
| <a href="#">NSF Division of Ocean Sciences (NSF OCE)</a> | <a href="#">OCE-2048666</a> |

[ [table of contents](#) | [back to top](#) ]