

Fluorometric measurements of chlorophyll a from R/V Roger Revelle cruise RR2201 in the Eastern Indian Ocean (Argo Basin) during February 2022

Website: <https://www.bco-dmo.org/dataset/943455>

Data Type: Cruise Results

Version: 1

Version Date: 2024-11-13

Project

» [Collaborative Research: Mesoscale variability in nitrogen sources and food-web dynamics supporting larval southern bluefin tuna in the eastern Indian Ocean](#) (BLOOFINZ-IO)

Program

» [Second International Indian Ocean Expedition](#) (IIOE-2)

Contributors	Affiliation	Role
Landry, Michael R.	University of California-San Diego Scripps (UCSD-SIO)	Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

This dataset contains fluorometric measurements of chlorophyll a from cruise RR2201 on R/V Roger Revelle (BLOOFINZ-IO, January-March 2022) in the Argo Basin region off NW Australia.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Coverage

Location: Northwest Australia, Argo Basin

Spatial Extent: N:-15.349 E:118.142 S:-17.126 W:114.135

Temporal Extent: 2022-02-03 - 2022-02-22

Methods & Sampling

Seawater was collected with Niskin bottles on a CTD rosette, filtered onto GF/F filters and extracted for 24 hours at -20 degrees Celsius (°C) with 90% acetone. Samples were analyzed for chlorophyll a and phaeopigment concentrations by the acidification method with a Turner 10AU fluorometer.

The RR2201 cruise report can be found at <http://hdl.handle.net/1834/43464>.

Data Processing Description

The data were calculated by standard equations from unacidified (Fo) and acidified (Fa) readings, water sample and solvent volumes, and factors derived from the fluorometer calibration against pure Chla standards.

BCO-DMO Processing Description

- Imported original file "RR2201_FluorometricChla_BCO-DMO Submission.xlsx" into the BCO-DMO system.
- Marked 'nd' as a missing data value (missing data are empty/blank in the final csv file).
- Renamed fields to comply with BCO-DMO naming conventions.
- Converted the "DateTimeUTC" column to ISO 8601 format and renamed "ISO_DateTime_UTC".
- Saved final file as "943455_v1_rr2201_fluorometric_chla.csv".

[[table of contents](#) | [back to top](#)]

Data Files

File
943455_v1_rr2201_fluorometric_chla.csv (Comma Separated Values (.csv), 11.82 KB) MD5:c86d0cab66feade2da84b29132135bd
Primary data file for dataset ID 943455, version 1

[[table of contents](#) | [back to top](#)]

Related Publications

Landry, M. R. (2025). R/V Roger Revelle Cruise RR2201 BLOOFINZ-IO Cruise Report. Eastern Indian Ocean larval tuna habitat study 20 January – 14 March 2022. <http://hdl.handle.net/1834/43464>
General

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Cruise	Cruise ID (RR2201)	unitless
Event	unique event number in UTC time as YYYYMMDD.mmss.###, where ### distinguishes events entered within the same minute	unitless
ISO_DateTime_UTC	Date and time (UTC) in ISO 8601 format	unitless
Latitude	latitude (North is positive; South is negative)	decimal degrees
Longitude	longitude (East is positive; West is negative)	decimal degrees
Cycle_Day	lagrangian experiments following a drogued drifter are noted as Cycle,Day, e.g. C1,D1; C1,D2, etc. EOC = end of cycle; e.g., EOC1	unitless
CTD_Cast	CTD cast number in Event Log	unitless
Depth	sampling depth	meters (m)
Chla	concentration of chla	milligrams per cubic meter (mg m-3)
Phaeo	concentration of phaeopigment	milligrams per cubic meter (mg m-3)
Chl_Integr	integrated Chla from surface to sample depth in CTD cast profile	milligrams per square meter (mg m-2)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Niskin bottles on a CTD rosette
Generic Instrument Name	Niskin bottle
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset-specific Instrument Name	Turner 10AU fluorometer
Generic Instrument Name	Turner Designs Fluorometer 10-AU
Generic Instrument Description	The Turner Designs 10-AU Field Fluorometer is used to measure Chlorophyll fluorescence. The 10AU Fluorometer can be set up for continuous-flow monitoring or discrete sample analyses. A variety of compounds can be measured using application-specific optical filters available from the manufacturer (read more from Turner Designs, turnerdesigns.com , Sunnyvale, CA, USA).

[[table of contents](#) | [back to top](#)]

Deployments

RR2201

Website	https://www.bco-dmo.org/deployment/916293
Platform	R/V Roger Revelle
Report	http://hdl.handle.net/1834/43464
Start Date	2022-01-20
End Date	2022-03-14
Description	See more information at R2R: https://www.rvdata.us/search/cruise/RR2201

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Mesoscale variability in nitrogen sources and food-web dynamics supporting larval southern bluefin tuna in the eastern Indian Ocean (BLOOFINZ-IO)

Coverage: Eastern Indian Ocean, Indonesian Throughflow area, and the Gulf of Mexico

NSF Award Abstract:

The small area between NW Australia and Indonesia in the eastern Indian Ocean (IO) is the only known spawning ground of Southern Bluefin Tuna (SBT), a critically endangered top marine predator. Adult SBT migrate thousands of miles each year from high latitude feeding areas to lay their eggs in these tropical waters, where food concentrations on average are below levels that can support optimal feeding and growth of their larvae. Many critical aspects of this habitat are poorly known, such as the main source of nitrogen nutrient that sustains system productivity, how the planktonic food web operates to produce the unusual types of zooplankton prey that tuna larvae prefer, and how environmental differences in habitat quality associated with ocean fronts and eddies might be utilized by adult spawning tuna to give their larvae a greater chance for rapid growth and survival success. This project investigates these questions on a 38-day expedition in early 2021, during the peak time of SBT spawning. This project is a US contribution to the 2nd International Indian Ocean Expedition (IIOE-2) that advances understanding of biogeochemical and ecological dynamics in the poorly studied eastern IO. This is the first detailed study of nitrogen and carbon cycling in the region linking Pacific and IO waters. The shared dietary preferences of SBT larvae with those of other large tuna and billfish species may also make the insights gained broadly applicable to understanding larval recruitment issues for top consumers in other marine ecosystems. New information from the study will enhance international management efforts for SBT. The shared larval dietary preferences of large tuna and billfish species may also extend the insights gained broadly to many other marine top consumers, including Atlantic bluefin tuna that spawn in US waters of the Gulf of Mexico. The end-to-end study approach, highlights connections among

physical environmental variability, biogeochemistry, and plankton food webs leading to charismatic and economically valuable fish production, is the theme for developing educational tools and modules through the "scientists-in-the-schools" program of the Center for Ocean-Atmospheric Prediction Studies at Florida State University, through a program for enhancing STEM learning pathways for underrepresented students in Hawaii, and through public outreach products for display at the Birch Aquarium in San Diego. The study also aims to support an immersive field experience to introduce talented high school students to marine research, with the goal of developing a sustainable marine-related educational program for underrepresented students in rural northwestern Florida.

Southern Bluefin Tuna (SBT) migrate long distances from high-latitude feeding grounds to spawn exclusively in a small oligotrophic area of the tropical eastern Indian Ocean (IO) that is rich in mesoscale structures, driven by complex currents and seasonally reversing monsoonal winds. To survive, SBT larvae must feed and grow rapidly under environmental conditions that challenge conventional understanding of food-web structure and functional relationships in poor open-ocean systems. The preferred prey of SBT larvae, cladocerans and Corycaeidae copepods, are poorly studied and have widely different implications for trophic transfer efficiencies to larvae. Differences in nitrogen sources - N fixation vs deep nitrate of Pacific origin - to sustain new production in the region also has implications for conditions that may select for prey types (notably cladocerans) that enhance transfer efficiency and growth rates of SBT larvae. The relative importance of these N sources for the IO ecosystem may affect SBT resiliency to projected increased ocean stratification. This research expedition investigates how mesoscale variability in new production, food-web structure and trophic fluxes affects feeding and growth conditions for SBT larvae. Sampling across mesoscale features tests hypothesized relationships linking variability in SBT larval feeding and prey preferences (gut contents), growth rates (otolith analyses) and trophic positions (TP) to the environmental conditions of waters selected by adult spawners. Trophic Positions of larvae and their prey are determined using Compound-Specific Isotope Analyses of Amino Acids (CSIA-AA). Lagrangian experiments investigate underlying process rates and relationships through measurements of water-column ^{14}C productivity, N_2 fixation, $^{15}\text{NO}_3^-$ uptake and nitrification; community biomass and composition (flow cytometry, pigments, microscopy, in situ imaging, genetic analyses); and trophic fluxes through micro- and mesozooplankton grazing, remineralization and export. Biogeochemical and food web elements of the study are linked by CSIA-AA (N source, TP), ^{15}N -constrained budgets and modeling. The project elements comprise an end-to-end coupled biogeochemistry-trophic study as has not been done previously for any pelagic ecosystem.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[[table of contents](#) | [back to top](#)]

Program Information

Second International Indian Ocean Expedition (IIOE-2)

Website: <https://web.whoi.edu/iioe2/>

Coverage: Indian Ocean

Description from the [program website](#):

The Second International Indian Ocean Expedition (IIOE-2) is a major global scientific program which will engage the international scientific community in collaborative oceanographic and atmospheric research from coastal environments to the deep sea over the period 2015-2020, revealing new information on the Indian Ocean (i.e. its currents, its influence upon the climate, its marine ecosystems) which is fundamental for future sustainable development and expansion of the Indian Ocean's blue economy. A large number of scientists from research institutions from around the Indian Ocean and beyond are planning their involvement in IIOE-2 in accordance with the overarching six scientific themes of the program. Already some large collaborative research projects are under development, and it is anticipated that by the time these projects are underway, many more will be in planning or about to commence as the scope and global engagement in IIOE-2 grows.

Focused research on the Indian Ocean has a number of benefits for all nations. The Indian Ocean is complex and drives the region's climate including extreme events (e.g. cyclones, droughts, severe rains, waves and storm surges). It is the source of important socio-economic resources (e.g. fisheries, oil and gas

exploration/extraction, eco-tourism, and food and energy security) and is the background and focus of many of the region's human populations around its margins. Research and observations supported through IIOE-2 will result in an improved understanding of the ocean's physical and biological oceanography, and related air-ocean climate interactions (both in the short-term and long-term). The IIOE-2's program will complement and harmonise with other regional programs underway and collectively the outcomes of IIOE-2 will be of huge benefit to individual and regional sustainable development as the information is a critical component of improved decision making in areas such as maritime services and safety, environmental management, climate monitoring and prediction, food and energy security.

IIOE-2 activities will also include a significant focus on building the capacity of all nations around the Indian Ocean to understand and apply observational data or research outputs for their own socio-economic requirements and decisions. IIOE-2 capacity building programs will therefore be focused on the translation of the science and information outputs for societal benefit and training of relevant individuals from surrounding nations in these areas.

A Steering Committee was established to support U.S. participation in IIOE-2. More information is available on their website at <https://web.whoi.edu/iioe2/>.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1851558

[[table of contents](#) | [back to top](#)]