

Nutrient profile data from discrete water samples collected during R/V Revelle cruise RR2104 and R/V Atlantis cruise AT50-10 in the Eastern Tropical and Subtropical Pacific Ocean from 2021 to 2023

Website: <https://www.bco-dmo.org/dataset/948503>

Data Type: Cruise Results

Version: 1

Version Date: 2025-01-14

Project

» [Collaborative Research: Underexplored Connections between Nitrogen and Trace Metal Cycling in Oxygen Minimum Zones Mediated by Metalloenzyme Inventories](#) (CliOMZ)

Contributors	Affiliation	Role
Santoro, Alyson E.	University of California-Santa Barbara (UCSB)	Principal Investigator
Bayer, Barbara	University of Vienna	Scientist, Contact
Newman, Sawyer	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	Data Manager

Abstract

These data include nutrient measurements from two oceanographic expeditions on R/V Roger Revelle from San Diego to San Diego, USA in June 2021 and on R/V Atlantic (CliOMZ AT50-10 expedition) from Golfo, Costa Rica to San Diego, USA in May-June 2023. Instruments used were a CTD profiler, a fluorometer, and a flow injection analysis system.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Location: Eastern Tropical and Subtropical Pacific Ocean

Spatial Extent: N:33.549 E:-88.9709 S:-10.0003 W:-124.7016

Temporal Extent: 2021-06-06 - 2023-06-16

Methods & Sampling

Water samples were collected during cruises RR2104 and AT50-10 in the eastern tropical and subtropical Pacific Ocean. Discrete water samples were obtained using a CTD rosette sampler equipped with 24 x 10 L Niskin bottles. Different depths were sampled ranging from 10 meters to 2200 meters.

Ammonium concentrations were measured on board from unfiltered 40 mL seawater samples using the o-phthalaldehyde derivatization method with modifications as suggested in (Taylor et al. 2007) on an Aquafluor 8000 handheld fluorometer. Samples for nitrite and nitrate concentration measurements were syringe-filtered

(0.22 μm , Sterivex) and stored at -20°C before concentrations were determined by Cd reduction coupled to colorimetric detection via the Griess assay on a Lachat autoanalyzer.

Data Processing Description

Within numeric fields, "0" values indicate measurement readings that are below detection limits.

BCO-DMO Processing Description

- The special character has been removed from "Nitrite+Nitrate" column in the primary data file. The column/parameter name has been changed to "Nitrite_plus_Nitrate."

[[table of contents](#) | [back to top](#)]

Data Files

File
948503_v1_cliomz_nutrient_profile_data.csv (Comma Separated Values (.csv), 10.98 KB) MD5:3a931d10c8c51cd44d2e980b02d0ca51
Primary data file for dataset ID 948503, version 1

[[table of contents](#) | [back to top](#)]

Related Publications

Taylor, B. D., Keep, C. F., Hall, R. O., Koch, B. J., Tronstad, L. M., Flecker, A. S. & Ulseth, A. J. (2007). Improving the fluorometric ammonium method: matrix effects, background fluorescence, and standard additions. *Journal of the North American Benthological Society* 26(2), 167-177. doi:[10.1899/0887-3593\(2007\)26\[167:ITFAMM\]2.0.CO;2](https://doi.org/10.1899/0887-3593(2007)26[167:ITFAMM]2.0.CO;2)
Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Cruise	Cruise name.	unitless
Date	Date of sampling.	unitless
Latitude	Latitude of sampling location in decimal degrees; a positive value indicates a northern coordinate.	decimal degrees
Longitude	Longitude of sampling location in decimal degrees; a negative value indicates a western coordinate.	decimal degrees
Station	Station number of cruises RR2104 and AT50-10.	unitless
Cast	Cast number of cruises RR2104 and AT50-10.	unitless
Depth	Sampling depth.	meters (m)
Nitrite	Concentration of nitrite from sample.	micromoles per liter (umol/L)
Nitrite_plus_Nitrate	Concentration of nitrite+nitrate from sample.	micromoles per liter (umol/L)
Ammonium	Concentration of ammonium from sample.	micromoles per liter (umol/L)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	QuikChem 8500 Series 2 Flow Injection System (Lachat Instruments)
Generic Instrument Name	Flow Injection Analyzer
Dataset-specific Description	Samples for nitrite and nitrate concentration measurements were syringe-filtered (0.22 µm, Sterivex) and stored at –20°C before concentrations were determined by Cd reduction coupled to colorimetric detection via the Griess assay on a Lachat autoanalyzer.
Generic Instrument Description	An instrument that performs flow injection analysis. Flow injection analysis (FIA) is an approach to chemical analysis that is accomplished by injecting a plug of sample into a flowing carrier stream. FIA is an automated method in which a sample is injected into a continuous flow of a carrier solution that mixes with other continuously flowing solutions before reaching a detector. Precision is dramatically increased when FIA is used instead of manual injections and as a result very specific FIA systems have been developed for a wide array of analytical techniques.

Dataset-specific Instrument Name	Aquafluor 8000 (Turner Designs)
Generic Instrument Name	Fluorometer
Dataset-specific Description	Ammonium concentrations were measured on board from unfiltered 40 mL seawater samples using the o-phthaldialdehyde derivatization method with modifications as suggested in (Taylor et al. 2007) on an Aquafluor 8000 handheld fluorometer.
Generic Instrument Description	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.

Dataset-specific Instrument Name	24x10 L Niskin Bottles
Generic Instrument Name	Niskin bottle
Dataset-specific Description	Discrete water samples were collected using a rosette sampler equipped with 24×10 L Niskin bottles.
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[[table of contents](#) | [back to top](#)]

Deployments

AT50-10

Website	https://www.bco-dmo.org/deployment/916122
Platform	R/V Atlantis
Report	https://www.rvdata.us/search/cruise/AT50-10
Start Date	2023-05-02
End Date	2023-06-09

RR2104

Website	https://www.bco-dmo.org/deployment/948513
Platform	R/V Roger Revelle
Start Date	2021-06-12
End Date	2021-07-01
Description	More information is available from R2R: https://www.rvdata.us/search/cruise/RR2104

Project Information

Collaborative Research: Underexplored Connections between Nitrogen and Trace Metal Cycling in Oxygen Minimum Zones Mediated by Metalloenzyme Inventories (CliOMZ)

Coverage: Eastern Tropical Pacific

NSF abstract:

Though scarce and largely insoluble, trace metals are key components of sophisticated enzymes (protein molecules that speed up biochemical reactions) involved in biogeochemical cycles in the dark ocean (below 1000m). For example, metalloenzymes are involved in nearly every reaction in the nitrogen cycle. Yet, despite direct connections between trace metal and nitrogen cycles, the relationship between trace metal distributions and biological nitrogen cycling processes in the dark ocean have rarely been explored, likely due to the technical challenges associated with their study. Availability of the autonomous underwater vehicle (AUV) Clio, a sampling platform capable of collecting high-resolution vertical profile samples for biochemical and microbial measurements by large volume filtration of microbial particulate material, has overcome this challenge. Thus, this research project plans an interdisciplinary chemistry, biology, and engineering effort to test the hypothesis that certain chemical reactions, such as nitrite oxidation, could become limited by metal availability within the upper mesopelagic and that trace metal demands for nitrite-oxidizing bacteria may be increased under low oxygen conditions. Broader impacts of this study include the continued development and application of the Clio Biogeochemical AUV as a community resource by developing and testing its high-resolution and adaptive sampling capabilities. In addition, metaproteomic data will be deposited into the recently launched Ocean Protein Portal to allow oceanographers and the metals in biology community to examine the distribution of proteins and metalloenzymes in the ocean. Undergraduate students will be supported by this project at all three institutions, with an effort to recruit minority students. The proposed research will also be synergistic with the goals of early community-building efforts for a potential global scale microbial biogeochemistry program modeled after the success of the GEOTRACES program, provisionally called "Biogeoscapes: Ocean metabolism and nutrient cycles on a changing planet".

The proposed research project will test the following three hypotheses: (1) the microbial metalloenzyme distribution of the mesopelagic is spatially dynamic in response to environmental gradients in oxygen and trace metals, (2) nitrite oxidation in the Eastern Tropical Pacific Ocean can be limited by iron availability in the upper mesopelagic through an inability to complete biosynthesis of the microbial protein nitrite oxidoreductase, and (3) nitrite-oxidizing bacteria increase their metalloenzyme requirements at low oxygen, impacting the distribution of both dissolved and particulate metals within oxygen minimum zones. One of the challenges to characterizing the biogeochemistry of the mesopelagic ocean is an inability to effectively sample it. As a sampling platform, we will use the novel biogeochemical AUV Clio that enables high-resolution vertical profile samples for biochemical and microbial measurements by large volume filtration of microbial particulate material on a research expedition in the Eastern Tropical Pacific Ocean. Specific research activities will be orchestrated to test the hypotheses. Hypothesis 1 will be explored by comparison of hydrographic, microbial distributions, dissolved and particulate metal data, and metaproteomic results with profile samples collected by Clio. Hypothesis 2 will be tested by incubation experiments using $^{15}\text{NO}_2^-$ oxidation rates on Clio-collected incubation samples. Hypothesis 3 will be tested by dividing targeted nitrite oxidoreductase protein copies by qPCR (quantitative polymerase chain reaction)-based nitrite oxidizing bacteria abundance (NOB) to determine if cellular copy number varies with oxygen distributions, and by metalloproteomic analyses of NOB cultures. The demonstration of trace metal limitation of remineralization processes, not just primary production, would transform our understanding of the role of metals in biogeochemical cycling and provide new ways with which to interpret sectional data of dissolved and particulate trace metal distributions in the ocean. The idea that oxygen may play a previously underappreciated role in controlling trace metals due not just to metals' physical chemistry, but also from changing biological demand, will improve our ability to predict trace metal distributions in the face of decreasing ocean oxygen content.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1924512

[[table of contents](#) | [back to top](#)]