

Sinking Organic Particle fluxes and stable C isotopes (collected with sediment traps) from the Eastern Tropical North Pacific on the R/V Sikuliaq cruise SKQ201617S in January 2017

Website: <https://www.bco-dmo.org/dataset/948735>

Data Type: Cruise Results

Version: 1

Version Date: 2025-01-17

Project

» [Dimensions: Diversity, assembly and function of microbial communities on suspended and sinking particles in a marine Oxygen Deficient Zone](#) (ETNP_ParticleOmics)

Program

» [Dimensions of Biodiversity](#) (Dimensions of Biodiversity)

Contributors	Affiliation	Role
Keil, Richard	University of Washington (UW)	Principal Investigator
Devol, Allan	University of Washington (UW)	Co-Principal Investigator
Fuchsman, Clara	University of Washington (UW)	Scientist
Duffy, Megan E.	University of Washington (UW)	Student
Neibauer, Jacquelyn A.	University of Washington (UW)	Technician
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Fluxes of sinking organic carbon and nitrogen and the isotopic composition of organic carbon were obtained from free floating, unpoisoned surface tethered sediment traps at St P2 (16.5°N 107°W) in the Eastern Tropical North Pacific Oxygen Deficient Zone in January 2017. These traps were deployed from the R/V Sikuliaq on cruise SKQ201617S. Trap depths ranged between 69 m and 965 m, and trap deployments ranged between 21 and 91 hours with deeper traps deployed for longer. The Oxygen Deficient Zone extended from 105 m to 820 m at this station. Two types of traps were deployed: 1) in shallow waters (150 m), net traps (1.24 m² opening area) were used. For both types of trap, the cod end had bottoms that were open during deployment and during an 8 hour equilibration period at the target depth performed to remove oxygen contamination. Cod ends were closed with a gate valve, using a pre-programmed electronic dissolving link (burn wire) system controlled by an onboard Arduino microcontroller to start collection at the correct depth, and a second gate valve that closed the top of the cod end before retrieval. Some trap deployments functioned as simple sediment traps, and some deployments were combined trap and in situ incubators. The combined trap incubators consisted of upper and lower chambers. The material used to calculate fluxes reported here was collected from the upper chamber and was not incubated. After every deployment, sediment trap material was filtered onto pre-combusted GF-75 filters (0.3 μm nominal pore size). To conform to community standards, zooplankton carcasses were not included in the measurements of carbon and nitrogen flux. Filter samples (particles only) were wafted with HCl overnight to remove carbonate and sent to the University of Washington Isolab facility in the Department of Earth and Space Sciences (Seattle, WA) for C and N analysis. These data were collected to improve our understanding of sinking fluxes of organic matter in the offshore Oxygen Deficient Zone, and to see whether Oxygen Deficient Zones reduce organic matter attenuation Megan Duffy, Jacquelyn Neibauer, and Allan Devol and Rick Keil from the University of Washington deployed these sediment trap systems. Clara Fuchsman and Megan Duffy analyzed the data.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [BCO-DMO Processing Description](#)
 - [Problem Description](#)

- [Data Files](#)
 - [Supplemental Files](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Location: Eastern Tropical North Pacific Station P2 (16.5°N 107°W) Depth profile

Spatial Extent: N:16.56 E:-106.97 S:16.52 W:-107.14

Temporal Extent: 2017-01-07 - 2017-01-14

Methods & Sampling

Free floating, unpoisoned surface tethered sediment traps were used to quantify fluxes of sinking particles at St P2 (16.5°N 107°W) in January 2017, deployed from the R/V Sikuliaq on cruise SKQ201617S. Trap depths ranged between 69 m and 965 m, and trap deployments ranged between 21 and 91 hours with deeper traps deployed for longer. Traps were deployed in arrays of two traps per line. Two types of traps were deployed: 1) in shallow waters (<150 m), traps with a solid plastic cone top (0.46 m² opening area) were used, 2) in deep waters (>150 m), net traps (1.24 m² opening area) modeled from (Peterson et al., 2005) were used. For both types of trap, the cod end had bottoms that were open during deployment and during an 8 hour equilibration period at the target depth performed to remove oxygen contamination. Cod ends were closed with a gate valve, using a pre-programmed electronic dissolving link (burn wire) system controlled by an onboard Arduino microcontroller to start collection at the correct depth, and a second gate valve that closed the top of the cod end before retrieval. The aspect ratio, or height/trap mouth diameter, of the net traps was 2.5 while the aspect ratio of the cone traps was 0.6. These two types of traps have been shown to collect material with similar efficiency when normalized by opening area (Cram et al., 2022). No salt solution was used with these traps and no poisons were used in any chamber.

Some trap deployments functioned as simple sediment traps, and some deployments were combined trap and in situ incubators. The combined trap incubators consisted of upper and lower chambers. The material used to calculate fluxes reported here was collected from the upper chamber and was not incubated. Sinking material was collected in the lower incubation chamber, then isolated by closed a gate valve and injected with ¹⁵N-NO₂⁻. While the incubation occurred (in situ), sinking material was collected in an upper chamber that never encountered the spiked material. However, we do not report d¹⁵N natural stable isotopes for the sediment trap material here due to proximity to the incubation.

After every deployment, sediment trap material was filtered onto pre-combusted GF-75 filters (0.3 μm nominal pore size). To conform to community standards, zooplankton carcasses were not included in the measurements of carbon and nitrogen flux. Filter samples (particles only) were wafted with HCl overnight to remove carbonate, dried at 40°C, packed into silver foil cups, and sent to the University of Washington Isolab facility in the Department of Earth and Space Sciences (Seattle, WA) for C and N analysis utilizing an Costech elemental analyzer attached to an isotope ratio mass spectrometer (ThermoFinnegan MAT 253).

Data Processing Description

ug C and N were converted to fluxes in Microsoft Excel using the area of the trap opening and the duration of the deployment.

BCO-DMO Processing Description

* Sheet 1 of submitted file "ETNP_2017_P2_sediment_traps.xlsx" was imported into the BCO-DMO data system.

Table will appear as Data File: 948735_v1_etnp-2017-sedtrap-fluxes.csv (along with other download format options).

** In the BCO-DMO data system missing data identifiers are displayed according to the format of data you access. For example, in csv files it will be blank (null) values. In Matlab .mat files it will be NaN values. When viewing data online at BCO-DMO, the missing value will be shown as blank (null) values.

* Date converted to ISO 8601 format

* Lat lon converted to decimal degrees (south and west are negative, degree symbols and directional NSEW removed)

* Supplemental reference table was attached without format changes.

Problem Description

Some sediment trap deployments were paired with in situ incubations with spiked 15N. While the material described here was not involved in those incubations, we choose not to report natural 15N for this material due to its proximity to spiked 15N.

[[table of contents](#) | [back to top](#)]

Data Files

File
Sediment trap fluxes and 13C from the offshore ETNP in January 2017 filename: 948735_v1_etnp-2017-sedtrap-fluxes.csv (Comma Separated Values (.csv), 1.00 KB) MD5:751db70f1bcd3c58956736926d74c587 Primary data file for dataset ID 948735, version 1. Sediment trap fluxes and 13C from St P2 in the offshore Eastern Tropical North Pacific Oxygen Deficient Zone in January 2017.

[[table of contents](#) | [back to top](#)]

Supplemental Files

File
Standard analysis filename: Standards_2017_sediment_traps_ETNP.xlsx (Microsoft Excel, 40.76 KB) MD5:fb5f43b9efd4d0f889ef4329e3bba63 A comparison of reference standards analyzed with this data to their official values.

[[table of contents](#) | [back to top](#)]

Related Publications

Cram, J. A., Fuchsman, C. A., Duffy, M. E., Pretty, J. L., Lekanoff, R. M., Neibauer, J. A., Leung, S. W., Huebert, K. B., Weber, T. S., Bianchi, D., Evans, N., Devol, A. H., Keil, R. G., & McDonnell, A. M. P. (2022). Slow Particle Remineralization, Rather Than Suppressed Disaggregation, Drives Efficient Flux Transfer Through the Eastern Tropical North Pacific Oxygen Deficient Zone. *Global Biogeochemical Cycles*, 36(1). Portico.

<https://doi.org/10.1029/2021gb007080> <https://doi.org/10.1029/2021GB007080>

Results

Fuchsman, C. A., Duffy, M. E., Cram, J. A., Huanca-Valenzuela, P., Gregory, B. P., Plough, L., Pierson, J. J., Fitzgerald, C. L., Devol, A. H., & Keil, R. G. (2024). Contributions of Vertically Migrating Metazoans to Sinking and Suspended Particulate Matter Fuel N₂ production in the Eastern Tropical North Pacific Oxygen Deficient Zone. <https://doi.org/10.22541/essoar.172745075.56787778/v1>

Results

Peterson, M. L., Wakeham, S. G., Lee, C., Askea, M. A., & Miquel, J. C. (2005). Novel techniques for collection of

sinking particles in the ocean and determining their settling rates. *Limnology and Oceanography: Methods*, 3(12), 520-532. doi:[10.4319/lom.2005.3.520](https://doi.org/10.4319/lom.2005.3.520)
Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Station	station sampled	unitless
latitude	location sampled (latitude)	decimal degrees
longitude	location sampled (longitude)	decimal degrees
Date_deployed	date deployed (ISO 8601 format)	unitless
Duration	Length of time for organic matter collection	hours (hr)
Depth	depth of trap	meters (m)
TrapID	Unique number for trap array	unitless
TrapType	trap type (cone or net trap)	unitless
Incubator	Did an insitu incubation occur as part of this deployment (Yes or No).	unitless
Trap_Area	The area of the trap opening	meters squared (m ²)
Flux_C	The sinking flux of organic C	micromols per meters squared per day umol/m ² -d
Flux_N	The sinking flux of organic N	micromols per meters squared per day umol/m ² -d
C_to_N	ratio of molar carbon to nitrogen concentrations (C:N)	unitless
d13C_VPDB	isotopic composition of C (d13C). Reference standard VPDB="Vienna Pee Dee Belemnite"	permil (0/00)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	Costech International Elemental Combustion System (ECS) 4010
Dataset-specific Description	Carbon and Nitrogen amounts and isotopic composition was measured with a Costech Elemental Analyzer attached to ThermoFinnigan MAT 253 isotope ratio mass spectrometer at the University of Washington Isolab facility in the Department of Earth and Space Sciences (Seattle, WA).
Generic Instrument Description	The ECS 4010 Nitrogen / Protein Analyzer is an elemental combustion analyser for CHNSO elemental analysis and Nitrogen / Protein determination. The GC oven and separation column have a temperature range of 30-110 degC, with control of +/- 0.1 degC.

Dataset-specific Instrument Name	ThermoFinnigan MAT 253
Generic Instrument Name	Isotope-ratio Mass Spectrometer
Dataset-specific Description	Carbon and Nitrogen amounts and isotopic composition was measured with a Costech Elemental Analyzer attached to ThermoFinnigan MAT 253 isotope ratio mass spectrometer at the University of Washington Isolab facility in the Department of Earth and Space Sciences (Seattle, WA).
Generic Instrument Description	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

[[table of contents](#) | [back to top](#)]

Deployments

SKQ201617S

Website	https://www.bco-dmo.org/deployment/828218
Platform	R/V Sikuliaq
Start Date	2016-12-20
End Date	2017-01-16
Description	Cruise DOI: 10.7284/907444 See more cruise information from the Rolling Deck to Repository (R2R): https://www.rvdata.us/search/cruise/SKQ201617S

[[table of contents](#) | [back to top](#)]

Project Information

Dimensions: Diversity, assembly and function of microbial communities on suspended and sinking particles in a marine Oxygen Deficient Zone (ETNP_ParticleOmics)

Coverage: Eastern Tropical North Pacific

Extracted from the NSF award abstract:

Marine oxygen deficient zones (ODZs) are waters that are functionally devoid of oxygen. Without oxygen, some microbes are capable of converting nitrogen in the water into N₂ gas, which then leaves the ocean and enters the atmosphere. This loss of an important nutrient from the ocean has impacts on phytoplankton growth and marine food webs. While oxygen deficient zones occupy a very small percentage of the ocean, they account for as much as half of the oceanic loss of N as N₂. Moreover, the size of these regions is predicted to expand during this century due to climate change. The microbes that are capable of producing N₂ gas are extremely diverse, and use several different biochemical pathways to carry out this process. They may occur both free-floating in the water and attached to small particles that are suspended or sinking from the surface waters and providing them a carbon source. However the importance of these two lifestyles (free-living vs particle attached) in terms of contributions to N loss from the oceans is not well understood. This project will identify the major organisms that result in N₂ gas production on both suspended and sinking particles, the chemical reactions they carry out, and the rates at which this occurs. This information will be used to improve global climate models to better predict rates of N loss in a future ocean. Elementary and middle school teachers enrolled in a Masters in Science for Science Teachers program will be involved in the project and the graduate students and post-doctoral researchers supported by the project will have opportunities to participate in their classrooms. Underserved populations will also be integrated into the research at the undergraduate and middle school level through a series of summer internships.

ODZs have very complex elemental cycles, implying great microbial diversity. Intertwined with the microbial complexity of ODZ regions is the relatively unexplored interplay between free-living bacteria and those living on either suspended or sinking particles. Determining how these communities and niches interact and relate is one of the most challenging components of ODZ system studies today. Current climate models portray the dynamics of particles in the ODZs and throughout the deep ocean through prescribed functions based on sparse data from the oxic ocean with microbes represented only by the net chemical reactions of the community. However, in reality a phylogenetically and metabolically diverse group of microbes, likely acting in consortia, are responsible for the nitrogen transformations that ultimately result in the production of N₂. To explore the processes maintaining the genetic diversity and functional redundancy in N loss processes, four research areas will be integrated: the community phylogenetic diversity (both taxonomic and genomic diversity) the genetic diversity of the proteins that carry out key N transformation processes (as seen through quantitative proteomics), the resulting biogeochemical functions (15N labeled nitrogen transformation rate measurements) and predictions about how this diversity and corresponding function may change in response to climate change (biogeochemical modeling). The approach will be to assay both phylogenetic (16S rRNA tag sequencing) and functional genetic diversity (genomics) on sinking particles collected using large-volume sediment traps. Phylogenetic and genomic studies will be intimately tied to measurements of activity - who is doing key biogeochemical transformations (proteomics) and what are the in situ rates at which they are doing them (using novel incubation systems). Data will then be used to model how diversity and corresponding function change on a range of time and space scales, from the sinking of a single particle to seasonal cycles. To understand the relationship of community diversity and function on suspended and sinking particles, a series of three cruises will be conducted in the Eastern Tropical North Pacific ODZ.

[[table of contents](#) | [back to top](#)]

Program Information

Dimensions of Biodiversity (Dimensions of Biodiversity)

Website: http://www.nsf.gov/funding/pgm_summ.jsp?pims_id=503446

Coverage: global

(adapted from the NSF Synopsis of Program)

Dimensions of Biodiversity is a program solicitation from the NSF Directorate for Biological Sciences. FY 2010 was year one of the program. [\[MORE from NSF\]](#)

The NSF Dimensions of Biodiversity program seeks to characterize biodiversity on Earth by using integrative, innovative approaches to fill rapidly the most substantial gaps in our understanding. The program will take a broad view of biodiversity, and in its initial phase will focus on the integration of genetic, taxonomic, and functional dimensions of biodiversity. Project investigators are encouraged to integrate these three dimensions to understand the interactions and feedbacks among them. While this focus complements several core NSF programs, it differs by requiring that multiple dimensions of biodiversity be addressed simultaneously, to understand the roles of biodiversity in critical ecological and evolutionary processes.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Environmental Biology (NSF DEB)	DEB-1542240

[[table of contents](#) | [back to top](#)]